

Validation documentation
APIs & excipients, liquid/semi-solid (other)

HiperScan GmbH

July 18, 2018

Contents

Introduction	7
Context of this document	7
Criteria for the inclusion of substances	8
Validation concept	9
Model creation procedure and validation runs	9
Particularities of individual substance classes	11
Significance of testing with <i>Apo-Ident</i>	12
Conclusion	13
Explanation of terminology	14
Summary	19
Validation samples	19
Validation results	19
Validation reports	21
Alfason Basis Cresa®	21
Allergika evening primrose oil cream 20% / Dermifant® kids cream	25
alpha-bisabolol (racemic) at least 85%	29
Angelica root oil	33
Anise oil, organic	37
Aqueous ointments	41
Argan oil	47
Balm Bio Nature	51
Banana aroma	55
Basil oil	59
Benzoin siam oil 20%	63
Bepanthen® Wund- und Heilsalbe	67
Bergamot oil	71
Cajeput / Niaouli oil	75
Camphor oil	79
Carrot seed oil	83
Cedar wood oil	87
Cinnamon bark oil, Ceylon	91
Citric oil	95
Citronella oil	99
Citus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil	103
Clove oil	109
Contramarum aroma	113
Cumin oil	117
Cypress oil	121
Dermatop® base ointment	125
Dexeryl®	129
Dichloroacetic acid	133
Dimeticone ointment 10% SR / Optiderm® lotion	137
Dwarf pine oil	141
Ethanol 70% pure / denatured	145
Eucalyptus oil / Rosemary oil	153
Excipial® almond oil ointment	157
Fat ointments	161
Fenistil® Gel	169
Fennel oil	173
Ginger oil	177
Grapefruit oil organic / Orange oil	181
Hydrophilic ointments	185
Immortelle oil	201

Juniper oil	205
La Roche-Posay Cold Cream Naturel	209
La Roche-Posay Toleriane	213
Lemon grass oil	217
Lime oil	221
Liniment ointments	225
Linola® fat creme / Linola® H fat N / Alfason® Repair	231
Lipolotio urea 5% F body lotion	235
Lygal® head ointment N 3%	239
Lygal® ointment base	243
Mandarin oil, green	247
Manuka oil	251
Marjoram oil	255
Matricaria chamomilla oil	259
Matricaria oil, morrocan	263
Matricaria oil, roman	267
Myrrh oil	271
Myrtle oil	275
Neroli oil	279
Neuroderm® care lotion / moisturising cream	283
Neuroderm® moisturising cream lipo	287
Nicotine	291
Ointment ash basis / Neribas®	295
Orange aroma	301
Oregano oil	305
Palmarosa oil, organic	309
Panthenol ointment Lichtenstein	313
Patchouli oil	317
Peppermint oil	321
Perfume oil vanilla	325
Raspberry aroma	329
Ravensara	333
Rose-geranium oil, organic	337
Rosewood oil	341
Sage oil	345
Salicylic white paraffin 10%	349
Sandalwood oil	353
Sebexol® formula basis / Sebexol® cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver	357
Siam benzoin	365
Solutio Cordes®	369
Spearmint oil	373
Spruce needle oil / Frankincense oil / Silvire fir oil	377
Stomahesive® adhesive paste	383
Strawberry aroma	387
Swiss pine oil	391
Tea tree oil	395
Thyme oil, red	399
Thyme oil, white / Coriander oil	403
Vetiver bourbon oil	407
Wart ointment InfectoPharm® (NRF 11.31)	411
Water and aqueous solutions	415
Wax ointment (stabilised)	421
White paraffin oil	427
Wintergreen oil	431
Yarrow oil	435
Ylang-ylang oil	439

Appendix	443
Additional calibration samples (<i>Type A</i>)	443
Additional validation samples (<i>Type B</i>)	445
Additional validation samples (<i>Type C</i>)	447
Requirements of validation	449
Conformity of <i>Apo-Ident</i> with the <i>European pharmacopeia</i>	450
Addition to the two-stage models	451
References	453
Index	455

Introduction

The unequivocal proof of the identity of pharmaceutical raw materials based on a monograph or traditional alternative methods is work-intensive, time-intensive and economically often no longer makes sense. Near-infrared spectroscopy (NIR) provides a new means here. It enables relatively easy, fast and nonetheless safe identity testing via the preparation and evaluation of spectra.

The analysis system *Apo-Ident* was developed specially for pharmacy use. Pharmacists have the duty to check the identity of all raw materials for extemporaneous products in their pharmacies. This normally takes place based on the monographs for the respective substances in the European Pharmacopoeia. But also NIR spectroscopy is described in the European Pharmacopoeia as an identification method which, as opposed to the methods incorporated in the respective monographs, is approved for testing [1] (quotation translated)

with the prerequisite that the same results (“namely the determination of the identity” [2]), (quotation translated) are achieved as with the described methods and instruments.

The *Apo-Ident* analysis system serves to identify raw materials for prescriptions in the defined manner according to *ApBetrO* [Pharmacies Rules and Regulations] §§ 6 and 11 at pharmacies (NIR spectroscopy as an alternative testing method). *Apo-Ident* consists of three components:

- An *NIR spectrometer*, which records the spectra of non-preprocessed raw materials in a measuring glass in diffuse reflection or transfection.
- The *QuickStep* spectroscopy software controls the instrument and records the spectra and user inputs via a pharmacy-specific software plug-in. It also generates the test protocol for documentation of the testing and storage of the printout to be signed in the pharmacy.
- The software module *IdentModule* incorporates *reference databases*. The spectra from the *QuickStep* software are presented to it for evaluation.

NIR spectroscopy is a very powerful analytical method. It is also able to establish the identity of several chemical compounds and mixes in as far as an appropriate database (technically correct: a [chemometric model](#)) was created. Identity testing with *Apo-Ident* is a very safe, very fast and easy to operate analytical method for testing a large number of raw materials.

Context of this document

The suitability of the instrument, method and database is proven as follows:

- *NIR spectroscopy as a method for identity testing*: The *Ph. Eur.* [3] describes NIR spectroscopy in *Section 2.2.40* as an analytical method which is also suitable for the identification of raw materials. Therefore, validation of the method as such is not necessary.
- *Performance of the instrument*: The *Ph. Eur.* [3] furthermore describes the apparatus and the testing of its performance in *Section 2.2.40*. The document *Erfüllung von 2.2.40 Ph. Eur. durch Apo-Ident* [4] compares the implementation by *Apo-Ident* with this monograph in order to prove that *Apo-Ident* meets the specifications of the Pharmacopoeia. Each individual instrument delivered to a pharmacy is qualified in accordance with the tests described in “*Control of Instrument Performance*”. In this test, the unit consisting of analysis instrument hardware and the *QuickStep* spectroscopy software is assessed. The result is documented in a test protocol which is kept at the pharmacy.
- *Validation of the database* is documented separately for each substance class. The report at hand documents the substance class *APIs & excipients, liquid/semi-solid (other)*.

The *Arbeitsgemeinschaft der Pharmazieräte Deutschlands (APD)* [Working Group of German Pharmacy Inspectors] has clarified the following in its resolution dated October 16, 2013 ([5], quotation translated):

NIR is a testing method incorporated in the Pharmacopoeia. The testing quality depends on the quality of the database stored. The APD views the use of NIR instruments in case of ensured validation of the databases used in conjunction with it as one of several options for identity testing.

The APD defined more precisely ([6], quotation translated) on October 1, 2014:

The use of near-infrared is a recognised testing method according to Ph. Eur. 8. For the use of NIR instruments in pharmacies for testing the identity of raw materials, sufficient and verifiable validation of the instrument used is required. The quality of the database stored by the instrument manufacturer is decisive for quality. Batch-specific differences with the same original substances must be taken into account if present.

So NIR is basically suitable. The validity of the reference database is proven with the existing validation documentation.

Criteria for the inclusion of substances

This validation documentation describes the results of the validation of the reference database for the substance class *APIs & excipients, liquid/semi-solid (other)*. Validation documentation is created for each published version of the reference database for all substance classes incorporated.

The reference database is incorporated in the software module *IdentModule*. During identification testing with *Apo-Ident*, spectra which are used for evaluation purposes are presented to it by the *QuickStep* software. In the same manner, the *IdentModule* is presented all validation spectra successively during the validation runs for evaluation purposes. The *IdentModule* responds respectively (without taking the initial assumption into account) with the identified substance or rejects it as unknown. The correctness of this response is checked for each possible initial assumption and counted.

The results are summarised for each substance and reproduced in this document. The core statement of this validation report is that the following criteria must be fulfilled for each database entry, so that *Apo-Ident* offers verification of identity by means of NIR for the relevant substance/substance group:

- The database is exclusively generated from spectra which have been recorded by *HiperScan GmbH* on traceable samples in pharmaceutical quality.
 - The samples are procured via typical pharmacy sources (*DAC III.2.: Bezugsquellennachweis für Rezepturbestandteile [reference source for prescription components]* [7]).
 - A valid manufacturer's certificate exists (content, purity and identity of the batch).
 - The identity was confirmed by a certified test laboratory or *HiperScan GmbH*.
- Each version of the reference database (every update) is validated in-full.
 - Calibration spectra (*Type A*), other spectra recorded under the control of *HiperScan GmbH* (*Type B*), and spectra from the field (*Type C*) are presented to the *IdentModule* for evaluation in three separately evaluated validation runs.
 - Here, no single *false positive* result may arise.
 - Here, the various substance classes are also tested for reciprocal rejection, where this is objectively justified (see *Summary* section).
- In the validation with spectra recorded under the control of *HiperScan GmbH*, spectra of at least one independent sample must be considered, i.e. spectra from a batch whose spectra have not been used for the generation of the database. In addition, the set of *Type A* and *Type B* spectra must originate from at least three different batches.
- Spectra of additional substances may be used for the generation of the database even though they will not be offered for identification by this database. The purpose is the reliable distinction from these substances.

- Any positive result of *Apo-Ident* confirms the identity of the substance/substance group and distinguishes it from all other substances of the database. In the case of substance groups, the result is ambiguous: Distinction from all substances not belonging to the group is proven. The substance is identified as a member of this group. However, within the substance group, it is not possible to reliably classify which substance has been tested.
- The criteria for clear identifiability are a [specificity](#) of 100 % ([true negative rate](#)) and a minimum distance in the distance matrix. See 2. d) under [Model creation procedure and validation runs](#).

Validation concept

[Chemometrics](#) is a statistical technique for the extraction of relevant chemical information from spectra. In mathematics, this method is described as *multivariate data analysis*. Chemometrics proceeds here as follows:

1. Collection of spectra for the *calibration sample*. The results (identities) of the calibration sample must be known. The calibration samples must be representative for the samples which are to be evaluated later. Therefore, they must take the various possible (physical) compositions into account. (Therefore, sourcing calibration samples for NIR from the specialist trade is superior to the use of CRS reference substances.)
2. The first mathematical step is *calibration*. Here, the [chemometric model](#) is calculated from the *calibration sample spectra* ([reference spectra](#)) and limits as well as some parameters are stipulated. The chemometric model is used later to calculate the analysis result (*prediction*).
3. Collection of further spectra for the *validation sample* which should be independent of the *calibration sample*. The results (identities) of the *validation sample* must also be known. The textbook suggests a random sample with a normal scope of 25 % to 50 % of the *calibration sample* [8].
4. The second technical data step is *validation*. Here, the [chemometric model](#) created is evaluated based on the spectra of the *validation samples*. As validation parameters for the identification, the *Ph. Eur. Section 2.2.40* [3] specifies the [specificity](#) and [robustness](#).

The validation step according to the textbook has the target of estimating the performance capability of the model created based on a random sample. In order to achieve the best possible precision, attention is paid to the calibration sample. In the field of pharmaceuticals, the safety of the method has priority. In order to be able to [validate](#) the model within the regulatory scope, the validation step must include probative force. For this purpose, the validation sample must be *representative and complete* in order to enable the testing of all cases.

A *sufficient number of batches* must be secured for validation because validation finally proves whether the number of batches in calibration suffices.

Each substance is validated individually. The validation results are documented per substance in this document. Moreover, the documents show how many and which batches have been used for creating the model or model validation.

At least one certificate is taken in for each substance from an accredited test laboratory for the independent testing of identity of the sample. The identification number of the corresponding test certificate is listed in the report, enabling traceability of a substance tested according to the monographs in the Pharmacopoeia.

Model creation procedure and validation runs

The safety of the [chemometric models](#) is guaranteed by several measures during model creation, of which the validation step is the final one. Normally, the procedure is as follows. It is in particular valid for the active pharmaceutical ingredients (APIs) *solid API excipients, liquid/semi-solid API excipients (with a test certificate), narcotics - solid medicinal substances and drugs*. If, for individual substance classes, variations are required, they are depicted in the section [Particularities of individual substance classes](#).

1. Collecting the reference spectra (calibration sample)

- a) Procurement of the samples from the same sources from which pharmacies source their raw material for compounding (Caelo, Fagron, Euro-OTC, . . . , see also *DAC III.2. Bezugsquellen-nachweis für Rezepturbestandteile* [Sources of supply for compounding] [7]).
- b) Testing the suitability according to *ApBetrO* [Pharmacies Rules and Regulations] §§ 6, 11, that is to say the availability of a valid manufacturer certificates via identity, purity and contents of the batch.
- c) Recording standard 40 spectra of the sample in different positions, as a standard on four instruments. Here, handling and presentation of the samples as later in the pharmacy.
- d) Visual checking for anomalies in the spectra. In case of indications of measurement errors, measurement must be repeated. If a signature is missing in the spectrum, the substance may be excluded from the start as not promising (the spectra are nonetheless entered in the database validation as independent *Type B* spectra).
- e) Testing identity. For each substance, a certificate of correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the respective following substance page of this validation documentation the *Mahalanobis distance* to this reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance. Such samples underline the statistical spread of the original reference substance, but cannot add any new characteristics of the substance.

HiperScan GmbH cooperates with some suppliers as follows: the raw materials supplier takes a sufficiently large sample in his incoming goods area so that part of it can be used for recording the NIR spectra. The remainder of the sample goes to analytics for market approval. The manufacturer's batch certificate emerges from these identity, contents and purity tests, which consequently also prove the correct identity of the NIR reference sample. Therefore, the NIR spectra are suitable for structuring the database (*Type A*) and can also optionally be used for validation (*Type B*). The samples which this applies to are marked in the validation report with a footnote.

- f) If the identity of the new sample is proven, it is declared as a reference sample and the spectra are approved for structuring the database.

2. Generating the chemometric models (calibration)

- a) Determination of the transformation matrix from the reference spectra using variance maximisation [8, 9]. (*All* reference spectra are always included, even if only a few spectra are added for an update.) All reference spectra receive the same data pre-treatment, which is also later applied to all measurement spectra in the field (in the pharmacies).
- b) Checking that the number of principal components used is still sufficient.
- c) Calculating the limits for each substance from the spread of the reference spectra. The calculation regulation is identical for each substance in a substance class.
- d) Checking the distances between the limits of the separable substances: the distance matrix incorporates the *Mahalanobis distances* from each substance to all the others. If a distance is less than 10, both two substances must be declared as inseparable or *both two* must be removed from the database. (The spectra of removed substances remain as independent *Type B* spectra in the validation.)
- e) Testing the model based on the reference spectra. If *false positive* results arise, you must proceed as in the case of *Mahalanobis distances* which are too small.
- f) If chemometric models are available for all substance classes which meet both criteria (distance matrix and no *false positives*), they are joined together with the evaluation algorithms as an *IdentModule* and encrypted. This unit can no longer be changed. Its overall function is tested by validation.

3. Set of validation spectra (validation samples)

The following is provided for validation:

- a) *Type A*: The reference spectra = calibration spectra from which the database was generated. These also include spectra from substances which the *chemometric model* should not identify, but were also recorded during generation in order to increase selectivity. (As a result, the model “learns” to differentiate from other substances which are actually unknown to it.)
- b) *Type B*: Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database. These also include reference spectra of other substance classes, and spectra that are not used as reference spectra. Samples are considered to be independent, if they originate from a batch, of which no spectra have been used for the generation of the database. (Up to *IdentModule 2018-01*, samples were still considered independent if the sampling was done independently, i.e. if they originate from another sales container.)
- c) *Type C*: Spectra from the field, which have not been recorded under the control of *HiperScan GmbH* and have not been used for the generation of the database. The spectra include not only substances of the substance class to be tested, but also substances from other classes.

All manufacturers’ batches from which spectra are used for the validation are listed by substance in this document: for substances included in the substance class *APIs & excipients, liquid/semi-solid (other)* in the respective validation reports; otherwise in attachments *A, B* and *C*.

Furthermore remains valid: validation spectra may only be removed if a error in the spectrum can be proven. Here, the spectra are not deleted, but instead placed on a *blacklist* incorporating the reason, date and initials in the commentary.

The section *Particularities of individual substance classes* treats the other substance classes from which *Type B* and *Type C* spectra are cited for validation purposes.

4. Validation runs and approval

- a) Validation spectra are transferred holistically to the *IdentModule* for evaluation in the same way as the spectroscopy software *QuickStep* transfers measured spectra.
- b) Following the provision of each spectrum, the *IdentModule* responds as to whether it has recognised a substance and which substance was recognised.
- c) The correctness of this response is checked for each possible initial assumption (each measurable substance with the substance class) and counted according to *true negative*, *false negative*, *true positive* and *false positive*. These figures are provided for each substance and additionally in the section *Summary*, separated according to types *A, B* and *C*.
- d) No *false positive* results whatsoever are permissible.
- e) If the criterion is also met for all substance classes, the *IdentModule* is approved.

Particularities of individual substance classes

Basically, *HiperScan GmbH* procures and tests the manufacturer’s certificate for the batch, commissions external testing of the identity of the sample or carries it out independently and stores the certificates. As described, this process is established for the Pharmacopoeia substances, that is to say for substance classes **APIs & excipients, solid**, **APIs & excipients, liquid/semi-solid (with analysis certificate)**, **Narcotic substances, liquid/semi-solid** and **Drugs**. Therefore, *HiperScan GmbH* is able to furnish proof of the identity of the reference samples. In case of manufacturer-specific substance classes and others, individual steps are organised differently in-part:

The substance class **APIs & excipients, liquid/semi-solid (other)** (often described as cosmetics) incorporates substances for which no specification of the requirements of the pharmaceutical quality is determined, neither in a Pharmacopoeia monograph, a DAC/NRF monograph nor via a manufacturer’s specification. Consequently, neither the identity nor contents can be tested independently. No certificates whatsoever exist for the reference samples. So here, merely the matching of the sample with former samples of this product is established and confusion with the other substances

is ruled-out. (If the manufacturer of such a substance prepares a specification, determines testing methods and provides manufacturer's certificates in accordance with *ApBetrO* [Pharmacies Rules and Regulations] §§ 6, 11, *HiperScan GmbH* can assign the substance to the substance class *APIs & excipients, liquid/semi-solid (with analysis certificate)* again in the future).

Substance class **HCK – nutritional supplements (Hepart)** contains the HCK micro-nutrients from the Swiss company *Hepart AG*. *HiperScan GmbH* receives the reference samples directly from the manufacturer. For each reference sample, *HiperScan GmbH* also receives manufacturer's certificates and keeps these. New checking of the identity of the reference sample is not carried out by *HiperScan GmbH*. The identity of the reference samples is therefore documented by *Hepart AG*. The spectra of all batches provided by *Hepart AG* are recorded by *HiperScan GmbH* and entered in the database.

All the manufacturer's batches are used for the generation and validation of the substance class *HCK*. The expected variation is also represented in the generation and validation if there are less than three batches.

Also, for the substance class **PhytoComm** (TCM-Granulated herbal extracts of the manufacturer *PhytoComm*) spectra for all useable batches are recorded by *HiperScan GmbH* and entered in the database. The supplier organises the respective tests themselves and keeps the test certificates.

A new evaluation option was created for the class *PhytoComm* with the update 2016-01. As the risks are considerably fewer than those from chemical agents, the pharmacist can specify a reasonable criterion for the *specificity* in accordance with internal risk estimation. The database for this is created without taking safety distances into account and no criterion is determined in advance for the *specificity*. Instead, the *specificity* for testing the identity with this concrete substance is calculated in the validation for each substance and provided with the measurement result. The pharmacist then judges himself whether this safety is reasonable with regard to the risk of the substance.

Additionally, a statistical forecast is provided for the *specificity* which is determined according to the *Rule of Three* [10, 11]. For this forecast, it is assumed that there would have been three wrong results more and is provided with a lower limit for *specificity*. This value has a special meaning if a *specificity* of 100 % is achieved for a substance during validation. In this case, the lower limit allows conclusions regarding the scale of existing safety for which with an endless number of validation spectra a value of less than 100 % is to be assumed.

If, for example 14000 spectra not belonging to the substance are presented and no *false positive* classification is made, a hypothetical number of three *false positive* results is assumed (*Rule of Three* [10, 11]) and the *specificity* is defined with 100.0000 % (> 99.9786 %). Here, it applies that the higher the number of validation spectra which form the statistical basis, the better the *specificity* calculated via the lower *specificity* limit will be approximated.

The positive result of the identity test using *Apo-Ident* establishes that the sample spectrum is in accordance with a batch of the specified granulate from the supplier *PhytoComm*, whereby all useable batches from the supplier are known.

The *PhytoComm* class can only confirm the identity of batches that have been used for the generation of the database. As a consequence there cannot be any validation spectra of other batches. Therefore, the criterion reads that two samples (from different sales containers) from each batch must exist, one for the structure of the database (*Type A*) and one for the validation (*Type B*).

Significance of testing with *Apo-Ident*

The analysis result is determined using sophisticated statistical methods according to state-of-the-art science and technology. Chemical and pharmaceutical knowledge is applied for the selection of the samples from which the calibration spectra and validation spectra are recorded. Otherwise it does not influence the further steps of model creation.

Verbally, the statement of the analysis result can be expressed as follows. Here "*the spectra match*" means that the criteria *Mahalanobis distance*, *outlier analysis* and *correlation* are met as shown in *Erfüllung von 2.2.40 Ph. Eur. durch Apo-Ident* [4]. "The spectra do not match", on the other hand, means that at least the criterion *Mahalanobis distance* is not met.

The positive analysis result "*was identified as ...*" is very meaningful because both the quantity of substances to be taken into account and the number of underlying samples is very comprehensive.

1. The spectrum of the sample measured matches spectra of the defined substance.

2. The spectrum of the sample measured does not match any spectrum of any other substance in this substance class. Therefore, all other substances can be clearly ruled-out.
3. As the spectra from other substance classes were used for validation, it is proven that no spectrum of one of these other substances matches the defined substance. (All substance classes with which a spectrum comparison is possible and makes sense are used for validation. This is documented for each substance class in the section [Summary](#).)
4. If the defined substance belongs to a [substance group](#) which in itself is not clearly separable with *Apo-Ident*, matching with the spectra of one or several substances in this group is confirmed. Which of these substances it actually is cannot be determined clearly. All other substances are excluded analogous to [2](#) and [3](#).

On the other hand, a negative analysis result “*was not identified as ...*” means:

1. The substance offered could not be recognised based on the spectrum of this sample.
2. The identity of this sample is not confirmed.
3. Testing must be repeated in accordance with the specifications of the Pharmacopoeia.

Conclusion

NIR spectroscopy is a testing method incorporated in the Pharmacopoeia. In case of successful database validation, it is a possible method for identity testing [\[5\]](#). *Apo-Ident* meets the criteria of the *European Pharmacopoeia* as a near-infrared spectrometer and proves the validity of the reference database with the existing validation documentation. This means that *Apo-Ident* can be used as an alternative testing method for testing raw materials at pharmacies.

Explanation of terminology

The following section serves to explain or define specialist terminology which is required in order to understand this document. If necessary, definitions for the analysis system *Apo-Ident* are defined more precisely.

The term database is used in this document exactly as in the *Ph. Eur. Section 2.2.40* [3] synonymous with [chemometric model](#). In order to differentiate the databases which are relatively independent of each other, *HiperScan GmbH* frequently also uses the term [substance class](#) (primarily in the plural). On the other hand, the spectra used to structure the database are termed spectrum collection and not database.

Substance classes are units of the organisational structure of the *IdentModule*. The substance classes are substance [databases](#) which are also broadly independently subscribable. On the one hand, the liquid and semi-solid substances are separated from the solid powders because they are measured against different references and therefore the spectra cannot be compared. On the other hand, for example the Pharmacopoeia substances are kept separated from the manufacturer-specific database *PhytoComm* for TCM (traditional Chinese medicine) raw materials.

The individual substance classes need only be limited against each other in-part. Often, no risk of confusion exists because they can only be procured from different sources. On the other hand, in several cases we handle substances which need not be distinguished. For example, em Huang Qi granulate from the company *PhytoComm* neither needs to be delimited from *Huang Qi* granulate from the company *HerbaSinica* nor is matching required. Respectively one single [chemometric model](#) is behind a substance class. (Even if several reciprocally secured chemometric models would be permissible.) The terms *substance class*, *chemometric model* and *databases* are mostly used here as synonyms.

A substance group respectively summarises all the substances within a [substance class](#) which cannot safely be distinguished from one another based on their NIR spectra. However, all the other substances in the database can be excluded.

The formation of subgroups is mentioned in the *Ph. Eur. Section 2.2.40* [3]. In this manner, technical restrictions in case of extensive databases can be avoided and it is possible to prepare individual subgroups with different spectrum pre-treatment. Validation of the subgroups against each other is required. *HiperScan GmbH* has solved these technical restrictions and doesn't use any subgroups within a substance class any longer.

Principal component analysis (PCA) [8, 9] is a multivariate statistics process or multivariate data analysis. It serves to structure, simplify and illustrate comprehensive data records by describing a large number of statistical variables by describing a lower number of linear combinations (the *principal components*) which are as significant as possible. In the *Apo-Ident IdentModule*, *PCA* is used to evaluate the recorded spectrum data (corresponding with *Ph. Eur. 2.2.40* [3]).

The term validation is defined in both relevant contexts here with different (even if related) meanings.

Within the sense of the expert discipline of *chemometrics*, validation is a process step when creating a [chemometric model](#): after a transformation matrix, limits and various parameters have been calculated or determined from a set of reference spectra during the course of the calibration step [8, 9], the validation step determines the performance capability of the model (selectivity, precision, ...) based on the validation spectra. Normally, random sampling is planned here. In order for the validation to gain strength of proof, the validation spectrum set must be selected with an appropriately wide scope (*representative* and *complete*). The terms *validation run* and *validation step* always actually mean the process step in this sense.

In the regulatory sense (of pharmaceutical production), validation is the documented proof that a process or system meets the previously specified requirements reproducibly when applied practi-

cally. In this sense, the *Apo-Ident* databases only become validated databases with the validation documentation, which this document is part of.

The *European Pharmacopeia* uses the term validation in *Section 2.2.40* within the sense of the specialist discipline of *chemometrics* [3].

The robustness of a process is the property of only being influenced by environmental fluctuations (e.g. temperature or humidity) a little. A method is robust if the environmental conditions do not or hardly falsify the final result.

The specificity of a classification (of a [chemometric model](#)) is the [true negative rate](#).

The recognition rate (also sensitivity) is the [true positive rate](#). It defines in how many percent of cases a correctly set up substance is actually confirmed.

The true negative rate describes the share of spectra correctly classified as non-identity during validation. This is equivalent to correct classification. It means that a substance *A* within identity checking as substance *B* is judged as “*not identified*”. The *true negative rate* is equivalent to the conditional frequency

$$h(\text{rejected}|\text{genuinely no identity}) = \frac{r_n}{r_n + f_p}$$

with r_n as the total number of *true negative* classifications and f_p as the total number of *false positive* classifications. For successful validation of an *IdentModule*, all spectra presented belonging to this category must be classified as *not in accordance*.

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of this number. The weight of each spectrum of a substance/substance group *i* therefore results as

$$w_i = \frac{1}{n_i}$$

with n_i number of spectra in this substance/substance group. This weighting ensures that the overall result cannot be enhanced by adding especially large numbers of spectra from easily separable substances.

The true positive rate describes the share of spectra correctly classified as identity during validation. This is equivalent to correct classification. It means that a substance *A* within identity checking as substance *A* is judged as “*identified*”. The *true positive rate* is equivalent to the conditional frequency

$$h(\text{identified}|\text{genuine identity}) = \frac{r_p}{r_p + f_n}$$

with r_p as the total number of *true positive* classifications and f_n as the total number of *false negative* classifications. The *true positive rate* is a measure for the recognition rate of the validated *Apo-Ident IdentModule*.

In order to ensure that each substance is received with the same weight, the spectra are weighted as described for the [true negative rate](#).

The true negative result describes a spectrum correctly classified as non-identity during validation. It is equivalent to correct classification. It means that a substance *A* within identity checking as substance *B* is judged as “*not identified*”.

The false positive result describes a spectrum falsely classified as non-identity during validation. This is the most critical type of possible false classification. It means that a substance *A* within identity checking as substance *B* is judged as “*identified*”. For successful validation of an *IdentModule*, a number of false positive events of zero are demanded for all spectra entering the validation. The exception to this restriction is the class of TCM granulates from the company *PhytoComm* as described under [Particularities of individual substance classes](#).

The true positive result describes a spectrum correctly classified as identity during validation. It is equivalent to correct classification. It means that a substance *A* within identity checking as substance *A* is judged as “*identified*”.

The false negative result describes a spectrum falsely classified as non-identity during validation. It is equivalent to false classification. It means that a substance *A* within identity checking as substance *A* is judged as “*not identified*”.

The ‘Rule of Three’ says that with a probability of 95 % the next random sample of the same size no more than three false results are to be expected if no false result existed in the existing random sample [10, 11].

The *specificity* and *recognition rate* are determined both globally and from the validation runs for all substances. The information is supplemented with the hypothetical value if there had been three false results more. The percent information is provided in parentheses with the “greater than” symbol ‘>’, e.g. *specificity* 100.000 % (>99.983 %) if 17 567 false spectra have been presented without one single *false positive* result. The larger the statistical basis, the lower the influence of the hypothetical false results.

The Mahalanobis distance is a distance measure between two points in *n*-dimensional vector space. Here, the respective direction component of the distance to *standard deviation* [12] of an *n*-dimensional distribution is standardised. In case of the *principal component analysis* [8, 9] this standardisation relates to the distribution of the respective calibration data set for a classification (substance/substance group) in the *principal component space* [8]. The *Mahalanobis distance* of a point (mapping of a spectrum) $\vec{y}$ in the *n*-dimensional principal component space to the expected value of an *n*-dimensional distribution $\mathbf{X}$ then results as

$$d(\mathbf{X}, \vec{y}) = \sqrt{(\bar{\mathbf{X}} - \vec{y})^T \mathbf{S}^{-1} (\bar{\mathbf{X}} - \vec{y})} \quad \text{with} \quad \mathbf{X} \in \mathbb{R}^{m \times n}, \vec{y} \in \mathbb{R}^m$$

[13]. Here, *m* is equivalent to the number of principal components used (dimension of the principal component space) and *n* the number of measurements existing in the calibration data set (spectra). $\bar{\mathbf{X}}$ is the expected value of the resulting distribution for the calibration data set (the average value of *n* measurements received). $\mathbf{S}^{-1}$ is the inverse covariance matrix [12] for distribution $\mathbf{X}$.




Figure 1: (a) The *Mahalanobis distance* from *A* to *B* is less than from *A* to *C*. However, the Euclidean distances are identical. (b) The *Mahalanobis distance* between the two measurement series *S4* and *S5* is smaller than between *S1* and *S2*. However, the Euclidean distances are identical.

The *Mahalanobis distance* offers advantages compared to the *Euclidean distance*: For the calculation of the distance it takes the statistical properties of a data point distribution (measurement series), i.e. average value, variance and covariance of the data points [14] into account. The *Mahalanobis distance* is applied while creating the reference database for evaluating the spectra from different samples of a substance.

A chemometric model is a classifier based on statistical methods [8, 9]. Through the respective algorithm used (e.g. *Principal Component Analysis*, *Cluster Analysis*), a maximum of chemical information is extracted from measurement data. Here, systematical or physical disturbances are eliminated using appropriate data pre-processing [15, 16].

At several places in this document, in order to simplify understanding, the term **database** is used instead of *chemometric model* – in the same manner as in the *Ph. Eur. Section 2.2.40* [3].

A sample (with its own sample ID) refers to substance in a sales container. Repeated sampling from the same sales container is listed under the same sample ID. (The suffix “SI” is not part of the sample ID.) Several samples may originate from the same batch. Samples are called “independent”, if they originate from a batch, of which no spectra have been used for the generation of the database. (Up to *IdentModule 2018-01*, samples from different sales packages were considered to be independent.) The information above the list of validation spectra now includes also the number of batches that deliver independent samples for the validation (for both *Type B* and *Type C*).

In case a supplier takes a sample for testing from its incoming goods and splits it to multiple laboratory containers, the substance in all laboratory containers will still be ascribed to the same sample. *HiperScan GmbH* only uses one of the subsamples.

Reference samples are used to structure the database. The *reference spectra* originate from these samples. In chemometric technical jargon you would normally say: For *calibration*, a **chemometric model** is generated from the *calibration spectra* recorded from the *calibration samples*, whose quality is subsequently assessed in **validation**.

Reference samples are procured via typical pharmacy sources. Their identity is tested. The *reference spectra* are recorded by *HiperScan GmbH*. The documentation also includes the manufacturer’s name and batch number.

Reference samples are clearly identified by a sample ID. Samples without sample ID may not be used as *reference samples*.

Summary

A total of 44 773 spectra from 2410 different batches for a total of 203 substances were used to validate the substance class *APIs & excipients, liquid/semi-solid (other)*.

Validation samples

The validation samples can be categorised as follows:

Type A Calibration spectra. These are the spectra used to generate the chemometric model. They were recorded by *HiperScan GmbH*. Detailed information regarding the batches or samples can be found in the following validation reports under *calibration samples* and under *Type A*. Further information is listed in [Appendix A](#).

Substance class	Substances	Batches	Spectra
APIs & excipients, liquid/semi-solid (other)	151	441	18 749

From category *A* a total of 18 749 spectra from 441 batches for a total of 151 substances were taken into account for validation.

Type B Spectra from independent samples which are not included in database generation. These spectra were recorded by *HiperScan GmbH*. Detailed information regarding the batches or samples can be found in the following validation reports in the section *Type B* or in [Appendix B](#).

Substance class	Substances	Batches	Spectra
APIs & excipients, liquid/semi-solid (other)	203	561	22 831

From category *B* a total of 22 831 spectra from 561 batches for a total of 203 substances were taken into account for validation.

Type C Spectra from independent samples which are not included in database generation. *Apo-Ident* customers carried out the measurements. Detailed information regarding the batches or samples can be found in the following validation reports in the section *Type C* or in [Appendix C](#).

Substance class	Substances	Batches	Spectra
APIs & excipients, liquid/semi-solid (other)	122	1696	3193

From category *C* a total of 3193 spectra from 1696 batches for a total of 122 substances were taken into account for validation.

Validation results

The validation runs checked whether all substances/substance groups in the substance class *APIs & excipients, liquid/semi-solid (other)* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, the matching of all relevant spectra of various substances with the substances/substance groups in the substance class *APIs & excipients, liquid/semi-solid (other)* was checked and the correctness of the results was evaluated. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	18 749	0	1 781 155
Type B	0	16 870	1630	2 172 040
Type C	0	2803	228	303 455

All substances/substance groups in the substance class *APIs & excipients, liquid/semi-solid (other)* can be clearly distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.000 00 % (> 99.999 52 %)	100.000 00 % (> 99.953 51 %)
Type B	100.000 00 % (> 99.999 51 %)	90.904 89 % (> 90.877 63 %)
Type C	100.000 00 % (> 99.983 02 %)	88.040 07 % (> 87.399 96 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Alfason Basis Cresa®**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30723-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Alfason Basis Cresa®

Special notes

When selecting the *Alfason Basis Cresa®* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Alfason Basis Cresa®*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Astellas Pharma	Alfason Basis Cr...	12B11/77	30723	50	not required
Astellas Pharma	Alfason Basis Cr...	13D06/75	31071	40	not required
Astellas Pharma	Alfason Basis Cr...	15I46/79	32064	40	not required
Astellas Pharma	Alfason Basis Cr...	16D35/77	32815	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 170 spectra of 4 reference samples from the substance/substance group *Alfason Basis Cresa*®. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 579 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 104 spectra from 4 reference samples of the substance/substance group *Alfason Basis Cresa*®.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Astellas Pharma	Alfason Basis Cresa®	12B11/77	30723†	4
Astellas Pharma	Alfason Basis Cresa®	15I46/79	32064†	20
Astellas Pharma	Alfason Basis Cresa®	16D35/77	32840	40
LEO Pharma GmbH	Alfason Basis Cresa®	17B27/75	33487	40

- 22 727 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

†These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 30 spectra from 9 *Apo-Ident* customers from 16 batches from the substance/substance group *Alfason Basis Cresa*[®].
- These include spectra of independent samples from 16 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Astellas Pharma	Alfason Basis Cresa [®]	13K15/75	2
Astellas	Alfason Basis Cresa [®]	13A16/76	1
Astellas Pharma	Alfason Basis Cresa [®]	14C32/78	1
Astellas	Alfason Basis Cresa [®]	14J02/77	1
Astellas	Alfason Basis Cresa [®]	14c32-78	1
Astellas Pharma	Alfason Basis Cresa [®]	14J02/77	2
Astellas	Alfason Basis Cresa [®]	14J02/77	1
Astellas / Noweda	Alfason Basis Cresa [®]	14J02/77	1
Noweda-Astella	Alfason Basis Cresa [®]	15A01/79	1
Astellas Pharma	Alfason Basis Cresa [®]	15D11/79	3
Astellas / Noweda	Alfason Basis Cresa [®]	15D11/79	2
Caeo/Noweda	Alfason Basis Cresa [®]	15H06/80	1
Astellas / Noweda	Alfason Basis Cresa [®]	15D1179	1
Astellas Pharma	Alfason Basis Cresa [®]	15H06/80	1
Astellas / Noweda	Alfason Basis Cresa [®]	15J35/76	1
Astellas Pharma/Noweda	Alfason Basis Cresa [®]	15J38/79	1
Astellas Pharma	Alfason Basis Cresa [®]	16A06/76	1
A.H.D.	Alfason Basis Cresa [®]	12/05/75	1
Sanacorp	Alfason Basis Cresa [®]	13A16/76	1
Sanacorp	Alfason Basis Cresa [®]	12F23/76	1
Lamotte/ Noweda	Alfason Basis Cresa [®]	16A06/76	2
AHD	Alfason Basis Cresa [®]	13A18/76	1
Sanacorp	Alfason Basis Cresa [®]	13E08/75	1
Ichtyol/Phönix	Alfason Basis Cresa [®]	13E08/75	1

- 3163 spectra from 303 *Apo-Ident* customers from a total of 1643 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Alfason Basis Cresa*[®] can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Alfason Basis Cresa*[®] and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	170	0	18 579
Type B	0	104	0	22 727
Type C	0	30	0	3163

The substance/substance group *Alfason Basis Cresa*® can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.4706 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.2308 %)
Type C	100.0000 % (> 98.3595 %)	100.0000 % (> 80.0000 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30723	30723	0.00	21.38
31071	31071	0.00	19.91
32064	32064	0.00	24.37
32815	32815	0.00	19.73

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Allergika evening primrose oil cream 20% / Dermi...**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30970-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Allergika evening primrose oil cream 20% / Dermifant[®] kids cream; Allergika evening primrose oil cream 20%; Dermifant[®] kids cream

Special notes

When selecting the *Allergika evening primrose oil cream 20% / Dermifant[®] kids cream* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Allergika evening primrose oil cream 20% / Dermifant[®] kids cream*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Allergika	Dermifant [®] kids ...	1210082	30970	80	not required
Allergika	Dermifant [®] kids ...	1303033	31062	40	not required
Allergika	Allergika evenin...	1504021	31725	40	not required
Allergika	Allergika evenin...	1511051	31871	40	not required
Allergika	Allergika evenin...	1609043	33486	40	not required
Allergika	Dermifant [®] kids ...	1705043	33639	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 280 spectra of 6 reference samples from the substance/substance group *Allergika evening primrose oil cream 20% / Dermifant[®] kids cream*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 6 different batches.
- 18 469 spectra from a total of 433 batches from further 149 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 200 spectra from 6 reference samples of the substance/substance group *Allergika evening primrose oil cream 20% / Dermifant[®] kids cream*.
- These include spectra of independent samples from 4 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Allergika	Allergika evening primrose o...	1504021	31725 [†]	20
Allergika	Allergika evening primrose o...	1511051	31871 [†]	20
Allergika	Dermifant [®] kids cream	1607032	32754	40
Allergika	Allergika evening primrose o...	1607031	32756	40
Allergika	Allergika evening primrose o...	1410062	32909	40
Allergika	Dermifant [®] kids cream	1801010	34033	40

- 22 631 spectra from a total of 543 batches from further 201 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 4 spectra from 1 *Apo-Ident* customers from 3 batches from the substance/substance group *Allergika evening primrose oil cream 20% / Dermifant® kids cream*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Gehe	Dermifant® kids cream	1312102	2
Gehe	Dermifant® kids cream	1407053	1
Allergika Pharma GmbH	Dermifant® kids cream	1410072	1

- 3189 spectra from 304 *Apo-Ident* customers from a total of 1656 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Allergika evening primrose oil cream 20% / Dermifant® kids cream* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Allergika evening primrose oil cream 20% / Dermifant® kids cream* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	280	0	18 469
Type B	0	200	0	22 631
Type C	0	4	0	3189

The substance/substance group *Allergika evening primrose oil cream 20% / Dermifant® kids cream* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.8571 %)
Type B	100.0000 % (> 99.9529 %)	100.0000 % (> 97.0000 %)
Type C	100.0000 % (> 98.3777 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30970	30970	0.00	18.89
31062	31062	0.00	24.18
31725	31725	0.00	21.29
31871	31871	0.00	18.56
33486	33486	0.00	18.38
33639	33639	0.00	18.98

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **alpha-bisabolol (racemic) at least 85%**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31464-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

alpha-bisabolol (racemic) at least 85%

Special notes

When selecting the *alpha-bisabolol (racemic) at least 85%* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *alpha-bisabolol (racemic) at least 85%*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	alpha-bisabolol ...	13360114	31464	40	not required
Caelo	alpha-bisabolol ...	13360114	31465	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *alpha-bisabolol (racemic)* at least 85%. These samples are listed above in the section [calibration samples](#). The reference samples originate from 1 different batches.
- 18 669 spectra from a total of 438 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *alpha-bisabolol (racemic)* at least 85%.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	alpha-bisabolol (racemic) at...	13360114	31464 [†]	20
Caelo	alpha-bisabolol (racemic) at...	13360114	31465 [†]	20
Caelo	alpha-bisabolol (racemic) at...	16271902	32782	40
Caelo	alpha-bisabolol (racemic) at...	17065801	33417	40

- 22 711 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *alpha-bisabolol (racemic)* at least 85%.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *alpha-bisabolol (racemic)* at least 85% can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *alpha-bisabolol (racemic)* at least 85% and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	118	2	22 711
Type C	0	0	0	3193

The substance/substance group *alpha-bisabolol (racemic)* at least 85% can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9530 %)	98.3333 % (> 95.8333 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31464	31464	0.00	54.86
31465	31465	0.00	57.54

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Angelica root oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30535-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Angelica root oil; Angelica archangelica oleum; Oleum angelicae (e radice)

Special notes

When selecting the *Angelica root oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Angelica root oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Angelica root oil	33019-109871BAG90451	30964	40	not required
Taoasis	Angelica root oil	4053-116078	31682	40	not required
Taoasis	Angelica root oil	2829199-122587	31780	40	not required
Taoasis	Angelica root oil	3029209-123992	32068	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Angelica root oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 589 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 180 spectra from 6 reference samples of the substance/substance group *Angelica root oil*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Angelica root oil	zu erfassen	30535	40
Taoasis	Angelica root oil	4053-116078	31682 [†]	20
Taoasis	Angelica root oil	2829199-122587	31780 [†]	20
Taoasis	Angelica root oil	3029209-123992	32068 [†]	20
Taoasis	Angelica root oil	3029209-125059	32477	40
Taoasis	Angelica root oil	1839219-127663	33644	40

- 22 651 spectra from a total of 544 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B*

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 4 spectra from 4 *Apo-Ident* customers from 4 batches from the substance/substance group *Angelica root oil*.
- These include spectra of independent samples from 4 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Angelica root oil	33019-110236BAG904	1
Caelo	Angelica root oil	11352704	1
Caelo	Angelica root oil	11352706	1
Taoasis	Angelica root oil	4053-117762	1

- 3189 spectra from 304 *Apo-Ident* customers from a total of 1655 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Angelica root oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Angelica root oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	140	40	22 651
Type C	0	2	2	3189

The substance/substance group *Angelica root oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9529 %)	77.7778 % (> 76.1111 %)
Type C	100.0000 % (> 98.3777 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to

exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30964	30964	0.00	30.28
31682	31682	0.00	31.41
31780	31780	0.00	29.80
32068	32068	0.00	32.43

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Anise oil, organic**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30670-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Anise oil, organic

Special notes

When selecting the *Anise oil, organic* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Anise oil, organic*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Anise oil, organic	2241027-120475	31681	40	not required
Taoasis	Anise oil, organic	1492-125379	32794	40	not required
Taoasis	Anise oil, organic	1492-125379	32935	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Anise oil, organic*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 629 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 4 reference samples of the substance/substance group *Anise oil, organic*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Anise oil, organic	B080009-99431BAG90451	30670	40
Taoasis	Anise oil, organic	2241027-120475	31681 [†]	20
Taoasis	Anise oil, organic	42803-290	33622	40
Taoasis	Anise oil, organic	42803-290	33875	40

- 22 691 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 25 spectra from 13 *Apo-Ident* customers from 19 batches from the substance/substance group *Anise oil, organic*.
- These include spectra of independent samples from 18 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Anise oil, organic	2241027-120196	1
Taoasis	Anise oil, organic	1492-121627	1
Taoasis/Noweda	Anise oil, organic	1465-121083h08	1
Taoasis	Anise oil, organic	26569-115911	2
Caelo	Anise oil, organic	12322503	2
Caelo	Anise oil, organic	12322507	1
Schubert	Anise oil, organic	1236E0-1518	1
Caelo	Anise oil, organic	13056105	1
Taoasis	Anise oil, organic	130910-113040	1
Taoasis	Anise oil, organic	1492-123225	1
Taoasis	Anise oil, organic	1492-125315	1
Taoasis	Anise oil, organic	1492-125379	2
Caelo	Anise oil, organic	15148901	1
Taoasis	Anise oil, organic	26735-108555	1
Taoasis	Anise oil, organic	26735-110455	1
Taoasis	Anise oil, organic	26735-111137	2
Taoasis/Sanacorp	Anise oil, organic	26735-111137	2
Taoasis/Sanacorp	Anise oil, organic	26735-112250	1
Taoasis	Anise oil, organic	26735-113054	1
Apotheker S Bauer&Comp	Anise oil, organic		1

- 3168 spectra from 302 *Apo-Ident* customers from a total of 1640 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Anise oil, organic* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Anise oil, organic* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	139	1	22 691
Type C	0	22	3	3168

The substance/substance group *Anise oil, organic* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	99.2857 % (> 97.1429 %)
Type C	100.0000 % (> 98.3600 %)	88.0000 % (> 76.0000 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31681	31681	0.00	41.90
32794	32794	0.00	44.02
32935	32935	0.00	47.11

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Aqueous ointments
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30726-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Aqueous ointments; Dermatest base ointment; Excipial[®] lipo cream; Fabitop[®] base cream

Special notes

When selecting the *Aqueous ointments* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	<i>Zusatz zu den Modellen der zweiten Stufe</i>

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Aqueous ointments*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Spirig Pharma	Excipial [®] lipo c...	I011	30726	40	not required
Fontapharm	Fabitop [®] base cr...	6RWW2	31082	40	not required
Spirig Pharma	Excipial [®] lipo c...	N017	31088	40	not required
Spirig Pharma	Excipial [®] lipo c...	R011	31520	40	not required
Fontapharm	Fabitop [®] base cr...	GWWL3	31576	40	not required

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
P&M Cosmetics	Dermatest base o...	251115	31627	40	not required
P&M Cosmetics	Dermatest base o...	180317	31950	40	not required
P&M Cosmetics	Dermatest base o...	121217	33039	40	not required
Spirig Pharma	Excipial [®] lipo c...	T081	33207	40	not required
Fontapharm	Fabitop [®] base cr...	GLWD5	33221	40	not required
Galderma Labo...	Excipial [®] lipo c...	V051	33636	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 440 spectra of 11 reference samples from the substance/substance group *Aqueous ointments*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 11 different batches.
- 18 309 spectra from a total of 428 batches from further 148 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 420 spectra from 13 reference samples of the substance/substance group *Aqueous ointments*.
- These include spectra of independent samples from 6 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Fontapharm	Fabitop [®] base cream	GRWW1	30555	40
P&M Cosmetics	Dermatest base ointment	120911	30737	40
Fontapharm	Fabitop [®] base cream	GRWW1	30835	40
P&M Cosmetics	Dermatest base ointment	070615	31053	20
Spirig Pharma	Excipial [®] lipo cream	N017	31276	40
P&M Cosmetics	Dermatest base ointment	231115	31288	40
Spirig Pharma	Excipial [®] lipo cream	R011	31520 [†]	20
Fontapharm	Fabitop [®] base cream	GWWL3	31576 [†]	20

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra
P&M Cosmetics	Dermatest base ointment	251115	31627 [†]	20
P&M Cosmetics	Dermatest base ointment	180317	31950 [†]	20
Galderma Laboratorium G...	Excipial [®] lipo cream	V051	33637	40
Fontapharm	Fabitor [®] base cream	GLWD4	33869	40
P&M Cosmetics	Dermatest base ointment	260820	34032	40

- 22 411 spectra from a total of 537 batches from further 200 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 181 spectra from 31 *Apo-Ident* customers from 53 batches from the substance/substance group *Aqueous ointments*.
- These include spectra of independent samples from 51 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Sanacorp	Dermatest base ointment	70615	1
P&M Cosmetics	Dermatest base ointment	210915	1
P&M Cosmetics/Anzag	Dermatest base ointment	122631	1
P&M	Dermatest base ointment	241115	1
Caelo	Excipial [®] lipo cream	9031	1
Gehe/Galderma	Excipial [®] lipo cream	L0TS031	1
Caelo	Excipial [®] lipo cream	L0TS023	1
Spirig	Excipial [®] lipo cream	N061	2
Spirig/Anzag	Excipial [®] lipo cream	R011	1
Galderma	Excipial [®] lipo cream	R012	3
Galderma	Excipial [®] lipo cream	R026	2
Galderma/Phönix	Excipial [®] lipo cream	R026	2
Galderma	Excipial [®] lipo cream	R035	2
Galderma	Excipial [®] lipo cream	R042	1
Galderma/Galderma	Excipial [®] lipo cream	R042	9
Spirig/Phönix	Excipial [®] lipo cream	R042	1
Galderma/Gehe	Excipial [®] lipo cream	R042	2
spirig	Excipial [®] lipo cream	r042	1
Galderma	Excipial [®] lipo cream	R012	1
Galderma/Phönix	Excipial [®] lipo cream	R012	1
Galderma/Galderma	Excipial [®] lipo cream	R026	1
Galderma/Galderma	Excipial [®] lipo cream	ro26	1
Galderma/Phönix	Excipial [®] lipo cream	R042	1
Galderma/Anzag	Excipial [®] lipo cream	S011	1
Galderma/Galderma	Excipial [®] lipo cream	S023	8
Spirig/Anzag	Excipial [®] lipo cream	S023	1

continued on the next page

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

continued from previous page

Supplier	Substance	Batch	Spectra
Galderma/ Noweda	Excipial [®] lipo cream	s023	6
Galderma/Phönix	Excipial [®] lipo cream	S023	1
Wolff	Excipial [®] lipo cream	S031	1
Spirig/Anzag	Excipial [®] lipo cream	S031	1
Galderma/ Noweda	Excipial [®] lipo cream	S031	9
Galderma/Phönix	Excipial [®] lipo cream	S031	1
Phönix/Galderma	Excipial [®] lipo cream	S023	1
Galderma/Galderma	Excipial [®] lipo cream	T021	8
Galderma/Phönix	Excipial [®] lipo cream	T021	1
Galderma	Excipial [®] lipo cream	T021	2
Galderma/ Noweda	Excipial [®] lipo cream	T021	5
Galderma/Noweda	Excipial [®] lipo cream	T021	1
Spirig Pharma	Excipial [®] lipo cream	T021	1
Galderma/ Noweda	Excipial [®] lipo cream	t022	5
Galderma/ Noweda	Excipial [®] lipo cream	T022	4
Galderma	Excipial [®] lipo cream	T061	2
Spirig (Galderma)	Excipial [®] lipo cream	T061	1
Galderma	Excipial [®] lipo cream	V051	1
Galderma	Excipial [®] lipo cream	v042	1
spirig	Excipial [®] lipo cream		1
Spirig /Gehe	Excipial [®] lipo cream	LOTN021	2
Spirig Pharma AG/ Sana...	Excipial [®] lipo cream	LOTN021	2
Spirig Pharma AG/ Anzag	Excipial [®] lipo cream	LOTR012	2
Spirig /Gehe	Excipial [®] lipo cream	LOTR012	1
spirig	Excipial [®] lipo cream	N018	4
Spirig Pharma AG/Allian...	Excipial [®] lipo cream	N018	1
spirig	Excipial [®] lipo cream	N021	5
Spirig	Excipial [®] lipo cream	N021	1
Gehe	Excipial [®] lipo cream	N061	1
Galderma	Excipial [®] lipo cream	N061	2
Spirig / Anzag	Excipial [®] lipo cream	N018	1
Spirig /Gehe	Excipial [®] lipo cream	R012	1
spirig	Excipial [®] lipo cream	R012	3
Spirig	Excipial [®] lipo cream	R012	1
Spirig/Gehe	Excipial [®] lipo cream	R012	1
Spirig Pharma Europe GmbH	Excipial [®] lipo cream	R026	1
Spirig/Phönix	Excipial [®] lipo cream	4622	1
Spirig Pharma	Excipial [®] lipo cream	813994/09	2
Spirig / Anzag	Excipial [®] lipo cream	813994/09	1
spirig	Excipial [®] lipo cream	813994/09	1
Spirig	Excipial [®] lipo cream	K021	1
Sanacorp	Excipial [®] lipo cream	LOTN016	1
Alliance Healthcare	Excipial [®] lipo cream		1
spirig	Excipial [®] lipo cream	M016	4
spirig	Excipial [®] lipo cream	M017	1
Spirig/AHD	Excipial [®] lipo cream	M017	1
Spirig / Anzag	Excipial [®] lipo cream	M017	1
spirig	Excipial [®] lipo cream	M041	5
Spirig/AHD	Excipial [®] lipo cream	M041	2
Spirig / Anzag	Excipial [®] lipo cream	M042	3
Spirig/Phoenix	Excipial [®] lipo cream	M042	2
spirig	Excipial [®] lipo cream	M042	2
Spirig Pharma AG	Excipial [®] lipo cream	M042	1
Alliance Healthcare	Excipial [®] lipo cream	M045	1
Sanacorp	Excipial [®] lipo cream	M027	1
Spirig / Anzag	Excipial [®] lipo cream	N018	3
Spirig/AHD	Excipial [®] lipo cream	N018	2
Spirig Pharma AG/GEHE	Excipial [®] lipo cream	R012	1
Fontapharm	Fabitop [®] base cream	GLWD2	1
Fontapharm/Gehe	Fabitop [®] base cream	GLWD4	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Gehe	Fabitor [®] base cream	GRWW2	6
Fontapharm	Fabitor [®] base cream	GLWW4	1
Artesan	Fabitor [®] base cream	GLWW5	1
jenne artisan	Fabitor [®] base cream	GLWW5	1
Phönix/Fontapharm	Fabitor [®] base cream	GRWW2	1
Jenne	Fabitor [®] base cream	GWLL1	1

- 3012 spectra from 303 *Apo-Ident* customers from a total of 1614 batches from a further 119 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Aqueous ointments* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Aqueous ointments* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	440	0	18 309
Type B	0	345	75	22 321
Type C	0	149	32	3002

The substance/substance group *Aqueous ointments* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9532 %)	100.0000 % (> 98.6364 %)
Type B	100.0000 % (> 99.9531 %)	82.1429 % (> 81.4286 %)
Type C	100.0000 % (> 98.3655 %)	82.3204 % (> 80.6630 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30726	30726	0.00	57.18
31082	31082	0.00	23.31
31088	31088	0.00	50.33
31520	31520	0.00	50.17
31576	31576	0.00	22.15
31627	31627	0.00	19.49
31950	31950	0.00	21.07
33039	33039	0.00	20.36
33207	33207	0.00	44.72
33221	33221	0.00	24.59
33636	33636	0.00	57.14

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Argan oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 32168-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Argan oil; Oleum Arganiae

Special notes

When selecting the *Argan oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Argan oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Argan oil	161443	32168	40	-*
Caelo	Argan oil	16035101	32571	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Argan oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Argan oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	Argan oil	161443	32168 [†]	20
Caelo	Argan oil	16035101	32306	40
Caelo	Argan oil	162759	32679	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

*The spectra were taken from a part of the same sample which the supplier used for its analysis (identity, purity, content). In this case the manufacturer certificates proof the identity and quality of the sample from those the spectra descend.

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Argan oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Argan oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Argan oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	100	0	22 731
Type C	0	0	0	3193

The substance/substance group *Argan oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
32168	32168	0.00	19.54
32571	32571	0.00	16.33

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Balm Bio Nature**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31645-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Balm Bio Nature

Special notes

When selecting the *Balm Bio Nature* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Balm Bio Nature*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Balm Bio Nature	150422-5106	31645	40	not required
Taoasis	Balm Bio Nature	150422-5106	31724	40	not required
Taoasis	Balm Bio Nature	151210-5337	32994	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Balm Bio Nature*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 629 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 132 spectra from 5 reference samples of the substance/substance group *Balm Bio Nature*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Balm Bio Nature	150422-5106	31645 [†]	20
Taoasis	Balm Bio Nature	150422-5106	31724 [†]	20
Taoasis	Balm Bio Nature	160401-6083	32786	40
Taoasis	Balm Bio Nature	170403-7087	33436	40
Taoasis	Balm Bio Nature	151210-5337	not required	12

- 22 699 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Balm Bio Nature*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Balm Bio Nature* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Balm Bio Nature* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	127	5	22 699
Type C	0	0	0	3193

The substance/substance group *Balm Bio Nature* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	96.2121 % (> 93.9394 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31645	31645	0.00	18.38
31724	31724	0.00	22.50
32994	32994	0.00	10.28

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Banana aroma**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31105-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Banana aroma; Banana flavour

Special notes

When selecting the *Banana aroma* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Banana aroma*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Banana aroma	12302509	31105	40	1404166
Caelo	Banana aroma	14353702	31606	40	not required
Caelo	Banana aroma	171890	33687	40	20170707*

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Banana aroma*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Banana aroma*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	Banana aroma	14353702	31606 [†]	20
Caelo	Banana aroma	16089305	33496	40
Caelo	Banana aroma	16089305	33608	40

- 22 731 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

*The spectra were taken from a part of the same sample which the supplier used for its analysis (identity, purity, content). In this case the manufacturer certificates proof the identity and quality of the sample from those the spectra descend.

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 2 spectra from 2 *Apo-Ident* customers from 2 batches from the substance/substance group *Banana aroma*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Banana aroma	12302509	1
Caelo	Banana aroma	14353707	1

- 3191 spectra from 304 *Apo-Ident* customers from a total of 1657 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Banana aroma* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Banana aroma* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	100	0	22 731
Type C	0	2	0	3191

The substance/substance group *Banana aroma* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3987 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31105	31105	0.00	159.39
31606	31606	0.00	158.62
33687	33687	0.00	157.44

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Basil oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30751-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Basil oil; Ocimum basilicum oleum; Oleum ocimum basilicum

Special notes

When selecting the *Basil oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Basil oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Basil oil	121018-109872BAG90451	30963	40	not required
Taoasis	Basil oil	561-119057	31755	40	not required
Taoasis	Basil oil	1500312-128851	33725	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Basil oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 4 reference samples of the substance/substance group *Basil oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Basil oil	21022-108746BAG90451	30751	40
Taoasis	Basil oil	561-119057	31755 [†]	20
Taoasis	Basil oil	1500312-126393	32972	40
Taoasis	Basil oil	1500312-126393	32973	40

- 22 691 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Basil oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Basil oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Basil oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	140	0	22 691
Type C	0	0	0	3193

The substance/substance group *Basil oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 95.7143 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30963	30963	0.00	21.52
31755	31755	0.00	132.83
33725	33725	0.00	167.64

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Benzoin siam oil 20%**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30456-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Benzoin siam oil 20%; Oleum styrax tonkinensis

Special notes

When selecting the *Benzoin siam oil 20%* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Benzoin siam oil 20%*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Benzoin siam oil...	12990-114892	31256	40	not required
Taoasis	Benzoin siam oil...	51939-123084	31843	40	not required
Taoasis	Benzoin siam oil...	63287-126233	32974	40	not required
Taoasis	Benzoin siam oil...	2161-1395	33855	40	not required
Taoasis	Benzoin siam oil...	32105-119788BAG90451	33944	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 200 spectra of 5 reference samples from the substance/substance group *Benzoin siam oil 20%*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 5 different batches.
- 18 549 spectra from a total of 434 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 4 reference samples of the substance/substance group *Benzoin siam oil 20%*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Benzoin siam oil 20%	474034-96350	30456	40
Taoasis	Benzoin siam oil 20%	51939-123084	31843 [†]	20
Taoasis	Benzoin siam oil 20%	63287-126233	33028	40
Taoasis	Benzoin siam oil 20%	2010-836	33724	40

- 22 691 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 16 spectra from 7 *Apo-Ident* customers from 15 batches from the substance/substance group *Benzoin siam oil 20%*.
- These include spectra of independent samples from 13 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Benzoin siam oil 20%	17578-116819	1
Taoasis	Benzoin siam oil 20%	23478-116955	2
Taoasis	Benzoin siam oil 20%	41242-120661	1
Taoasis	Benzoin siam oil 20%	31343-118365	1
Taoasis	Benzoin siam oil 20%	41242-121512	1
Taoasis	Benzoin siam oil 20%	54125	1
Taoasis	Benzoin siam oil 20%	51939-123084	1
Taoasis	Benzoin siam oil 20%	59023-125420	1
Taoasis	Benzoin siam oil 20%	62330-124547	1
Taoasis	Benzoin siam oil 20%	63287-126233	1
Taoasis	Benzoin siam oil 20%	548481-111780	1
Taoasis	Benzoin siam oil 20%	541389-111081	1
Taoasis	Benzoin siam oil 20%	548442-113119	1
Taoasis	Benzoin siam oil 20%	474084-96350	1
Taoasis	Benzoin siam oil 20%	518434-108321	1

- 3177 spectra from 304 *Apo-Ident* customers from a total of 1644 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Benzoin siam oil 20%* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Benzoin siam oil 20%* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	200	0	18 549
Type B	0	120	20	22 691
Type C	0	15	1	3177

The substance/substance group *Benzoin siam oil 20%* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.0000 %)
Type B	100.0000 % (> 99.9530 %)	85.7143 % (> 83.5714 %)
Type C	100.0000 % (> 98.3619 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31256	31256	0.00	242.01
31843	31843	0.00	232.67
32974	32974	0.00	253.67
33855	33855	0.00	238.83
33944	33944	0.00	253.52

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Bepanthen® Wund- und Heilsalbe
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	31048-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Bepanthen® Wund- und Heilsalbe

Special notes

When selecting the *Bepanthen® Wund- und Heilsalbe* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Bepanthen® Wund- und Heilsalbe*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Bayer	Bepanthen® Wund-...	GP00LB9	31048	40	not required
Bayer	Bepanthen® Wund-...	GP012SS	31758	40	not required
Bayer	Bepanthen® Wund-...	GP01CTH	33613	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Bepanthen® Wund- und Heilsalbe*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Bepanthen® Wund- und Heilsalbe*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Bayer	Bepanthen® Wund- und Heilsalbe	GP012SS	31758 [†]	20
Bayer	Bepanthen® Wund- und Heilsalbe	GP018T5	32720	40
Bayer	Bepanthen® Wund- und Heilsalbe	GP019ZF	32948	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Bepanthen® Wund- und Heilsalbe*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Bepanthen® Wund- und Heilsalbe* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Bepanthen® Wund- und Heilsalbe* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	100	0	22 731
Type C	0	0	0	3193

The substance/substance group *Bepanthen® Wund- und Heilsalbe* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31048	31048	0.00	37.54
31758	31758	0.00	38.72
33613	33613	0.00	38.98

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Bergamot oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30307-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Bergamot oil; Oleum bergamottae; Oleum citrus bergamia

Special notes

When selecting the *Bergamot oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Bergamot oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Bombastus	Bergamot oil	290998	31641	40	not required
Bombastus	Bergamot oil	290998	31760	40	not required
Taoasis	Bergamot oil	3221028-121642	31784	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Bergamot oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 629 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 5 reference samples of the substance/substance group *Bergamot oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Bombastus	Bergamot oil	00237459	30307	40
Bombastus	Bergamot oil	290998	31641 [†]	20
Bombastus	Bergamot oil	290998	31760 [†]	20
Taoasis	Bergamot oil	3221028-121642	31784 [†]	20
Taoasis	Bergamot oil	1968-128263	33437	40

- 22 691 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 15 spectra from 12 *Apo-Ident* customers from 14 batches from the substance/substance group *Bergamot oil*.
- These include spectra of independent samples from 14 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Bombastus	Bergamot oil	53731-125890	1
Taoasis	Bergamot oil	d01be14m-121110bag90	1
Caelo	Bergamot oil	12020702	1
Kaders	Bergamot oil	130564	1
Caelo	Bergamot oil	13141501	2
Noweda	Bergamot oil	271858	1
Taoasis	Bergamot oil	32338-109135	1
Taoasis	Bergamot oil	32579-111866	1
Taoasis	Bergamot oil	33697-112408	1
Taoasis	Bergamot oil	33110-110701	1
Taoasis	Bergamot oil	B130301-110610	1
Taoasis	Bergamot oil	96243	1
Bombastus	Bergamot oil	264008	1
Bombastus	Bergamot oil	280260	1

- 3178 spectra from 303 *Apo-Ident* customers from a total of 1645 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Bergamot oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Bergamot oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	114	26	22 691
Type C	0	12	3	3178

The substance/substance group *Bergamot oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	81.4286 % (> 79.2857 %)
Type C	100.0000 % (> 98.3623 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31641	31641	0.00	38.36
31760	31760	0.00	38.50
31784	31784	0.00	33.61

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Cajeput / Niaouli oil
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	31136-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Cajeput / Niaouli oil; Cajeput; Niaouli oil; Oleum Cajeputi artificiale; Oleum melaleuka viridiflora; Oleum niaouli

Special notes

When selecting the *Cajeput / Niaouli oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Cajeput / Niaouli oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Cajeput	130829-112094BAG90451	31136	40	not required
Caelo	Niaouli oil	13099606	31420	40	AR-14-FG-012561-01

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Niaouli oil	34167-116962	31548	40	not required
Taoasis	Cajeput	1662-124326	32173	40	not required
Taoasis	Niaouli oil	40428-124242	32174	40	not required
Taoasis	Niaouli oil	1500450-125771	32970	40	not required
Taoasis	Cajeput	1724-127033	33657	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 280 spectra of 7 reference samples from the substance/substance group *Cajeput / Niaouli oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 7 different batches.
- 18 469 spectra from a total of 432 batches from further 149 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 240 spectra from 7 reference samples of the substance/substance group *Cajeput / Niaouli oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Cajeput	T149I11	30382	40
Caelo	Niaouli oil	13099606	31420 [†]	20
Taoasis	Niaouli oil	34167-116962	31548 [†]	20
Taoasis	Niaouli oil	1500450-125771	32971	40
Taoasis	Cajeput	1724-127033	33022	40
Taoasis	Cajeput	1724-127033	33023	40
Taoasis	Niaouli oil	1500450-1985	34040	40

- 22 591 spectra from a total of 543 batches from further 201 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 26 spectra from 6 *Apo-Ident* customers from 21 batches from the substance/substance group *Cajeput / Niaouli oil*.
- These include spectra of independent samples from 21 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
PVL	Cajeput	150712	1
Aurica	Cajeput	11024710	1
Taoasis	Cajeput	1301-121889	1
Taoasis	Cajeput	1301-32758	1
Taoasis	Cajeput	130829-113045	2
Taoasis	Cajeput	131010-115487	2
Taoasis	Cajeput	131010-117241	2
Taoasis	Cajeput	131010-117257	1
Taoasis	Cajeput	1518-122691	1
Taoasis	Cajeput	1518-122790	1
Taoasis	Cajeput	1662-124870	1
Taoasis	Cajeput	1724-125910	1
Taoasis	Cajeput	24077-106468	1
Taoasis	Cajeput	661027-117257	2
Taoasis	Cajeput	661027-118368	1
Taoasis	Cajeput	6781012-119058	1
Taoasis	Cajeput	L201323018-113057	1
Taoasis/Sanacorp	Cajeput	L201323018-113057	1
Taoasis	Niaouli oil	22553-120413	1
Taoasis	Niaouli oil	22553-120482	1
Taoasis	Niaouli oil	22553-121510	1
Taoasis	Niaouli oil	40428-123092	1

- 3167 spectra from 304 *Apo-Ident* customers from a total of 1638 batches from a further 120 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Validation results

The validation runs checked whether the substance/substance group *Cajeput / Niaouli oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Cajeput / Niaouli oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	280	0	18 469
Type B	0	211	29	22 591
Type C	0	15	11	3167

The substance/substance group *Cajeput / Niaouli oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.8571 %)
Type B	100.0000 % (> 99.9529 %)	87.9167 % (> 86.6667 %)
Type C	100.0000 % (> 98.3599 %)	57.6923 % (> 46.1538 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31136	31136	0.00	42.25
31420	31420	0.00	21.32
31548	31548	0.00	30.93
32173	32173	0.00	59.60
32174	32174	0.00	24.57
32970	32970	0.00	23.61
33657	33657	0.00	57.37

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Camphor oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 33826-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Camphor oil

Special notes

When selecting the *Camphor oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Camphor oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Camphor oil	74521-140	33826	40	not required
Taoasis	Camphor oil	944	33827	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Camphor oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 40 spectra from 1 reference samples of the substance/substance group *Camphor oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Camphor oil	78793-2563	34039	40

- 22 791 spectra from a total of 548 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Camphor oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Camphor oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Camphor oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	40	0	22 791
Type C	0	0	0	3193

The substance/substance group *Camphor oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9535 %)	100.0000 % (> 85.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
33826	33826	0.00	27.90
33827	33827	0.00	27.51

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Carrot seed oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31555-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Carrot seed oil; Oleum daucus carota

Special notes

When selecting the *Carrot seed oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Carrot seed oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Carrot seed oil	937-117443	31555	40	not required
Taoasis	Carrot seed oil	1775-125361BAG90451	32812	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Carrot seed oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 160 spectra from 4 reference samples of the substance/substance group *Carrot seed oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Carrot seed oil	937-117443	31555 [†]	20
Taoasis	Carrot seed oil	26058_119083BAG90451	31636	60
Taoasis	Carrot seed oil	1775-125361BAG90451	33203	40
Taoasis	Carrot seed oil	2037	33995	40

- 22 671 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Carrot seed oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Carrot seed oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Carrot seed oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	148	12	22 671
Type C	0	0	0	3193

The substance/substance group *Carrot seed oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9529 %)	92.5000 % (> 90.6250 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31555	31555	0.00	31.53
32812	32812	0.00	70.44

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Cedar wood oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30654-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Cedar wood oil; Oleum cedrus deodara

Special notes

When selecting the *Cedar wood oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Cedar wood oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Cedar wood oil	7891018-121034	31777	40	not required
Apotheker Bau...	Cedar wood oil	2.880926.1507	31851	40	not required
Taoasis	Cedar wood oil	1627-125061	33199	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Cedar wood oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 200 spectra from 6 reference samples of the substance/substance group *Cedar wood oil*.
- These include spectra of independent samples from 4 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Cedar wood oil	zu erfassen	30654	40
Taoasis	Cedar wood oil	26473-83914BAG90451	30750	40
Taoasis	Cedar wood oil	7891018-121034	31777 [†]	20
Apotheker Bauer & Cie	Cedar wood oil	2.880926.1507	31851 [†]	20
Taoasis	Cedar wood oil	1627-124623	32268	40
Taoasis	Cedar wood oil	1802-1975	34102	40

- 22 631 spectra from a total of 544 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B*

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 15 spectra from 4 *Apo-Ident* customers from 12 batches from the substance/substance group *Cedar wood oil*.
- These include spectra of independent samples from 11 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis/Phönix	Cedar wood oil	3292687	1
Taoasis	Cedar wood oil	831018-122540	1
Taoasis	Cedar wood oil	831018-123089	2
Taoasis	Cedar wood oil	1627-125061	1
Biofrid / Noweda	Cedar wood oil	17066	1
Taoasis GmbH	Cedar wood oil	32812	1
Taoasis	Cedar wood oil	32812-113008	3
Taoasis	Cedar wood oil	381008-116967	1
Taoasis	Cedar wood oil	7711023-120348	1
Taoasis	Cedar wood oil	7891018-120108	1
Taoasis	Cedar wood oil	7891019-121034	1
Taoasis	Cedar wood oil	831018-124172	1

- 3178 spectra from 304 *Apo-Ident* customers from a total of 1647 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Cedar wood oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Cedar wood oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	159	41	22 631
Type C	0	15	0	3178

The substance/substance group *Cedar wood oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9529 %)	79.5000 % (> 78.0000 %)
Type C	100.0000 % (> 98.3623 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31777	31777	0.00	79.52
31851	31851	0.00	20.95
33199	33199	0.00	81.71

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Cinnamon bark oil, Ceylon
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30957-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Cinnamon bark oil, Ceylon; Oleum cinnamomi ceylanici

Special notes

When selecting the *Cinnamon bark oil, Ceylon* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Cinnamon bark oil, Ceylon*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Cinnamon bark oi...	121108-109782BAG90451	30957	40	not required
Taoasis	Cinnamon bark oi...	1311130547-119819	31648	40	not required
Taoasis	Cinnamon bark oi...	415-770	34107	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Cinnamon bark oil, Ceylon*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Cinnamon bark oil, Ceylon*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Cinnamon bark oil, Ceylon	1311130547-119819	31648 [†]	20
Taoasis	Cinnamon bark oil, Ceylon	415-126411	32791	40
Taoasis	Cinnamon bark oil, Ceylon	415-126411	33202	40

- 22 731 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Cinnamon bark oil, Ceylon*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Cinnamon bark oil, Ceylon* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Cinnamon bark oil, Ceylon* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	100	0	22 731
Type C	0	0	0	3193

The substance/substance group *Cinnamon bark oil, Ceylon* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30957	30957	0.00	244.83
31648	31648	0.00	241.29
34107	34107	0.00	245.14

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Citric oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31558-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Citric oil; Limonis aetheroleum; Oleum citri

Special notes

When selecting the *Citric oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Citric oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Citric oil	D04LI14M-117513	31558	40	not required
Taoasis	Citric oil	3681027-122724	31838	40	not required
Taoasis	Citric oil	127939-ITJ04	33611	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Citric oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *Citric oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Citric oil	D04LI14M-117513	31558 [†]	20
Taoasis	Citric oil	3681027-122724	31838 [†]	20
Taoasis	Citric oil	03LI16M-125453I09	32771	40
Taoasis	Citric oil	03LI16M-125453I09	32941	40

- 22 711 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 17 spectra from 15 *Apo-Ident* customers from 14 batches from the substance/substance group *Citric oil*.
- These include spectra of independent samples from 14 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Fagron	Citric oil	14H06-B02-308273	1
Sanacorp	Citric oil	15182703	1
Caelo	Citric oil	16028702	1
Fagron	Citric oil	13K14-B03-294704	1
Fagron	Citric oil	14H06-B02-299949	1
Caelo	Citric oil	15182703	1
Fagron	Citric oil	15F29-B04	1
Fagron	Citric oil	15f29-b04-325354	1
Bombastus	Citric oil	15F29-B04-325354	1
Caelo	Citric oil	16028719	2
Caelo	Citric oil	16275202	1
Caelo	Citric oil	16275216	1
Bombastus	Citric oil	301912	1
Taoasis	Citric oil	3681003-124333	1
Caelo	Citric oil	14365212	2

- 3176 spectra from 301 *Apo-Ident* customers from a total of 1645 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Citric oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Citric oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	120	0	22 711
Type C	0	17	0	3176

The substance/substance group *Citric oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 95.0000 %)
Type C	100.0000 % (> 98.3616 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31558	31558	0.00	17.34
31838	31838	0.00	15.22
33611	33611	0.00	16.19

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Citronella oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31550-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Citronella oil; Oleum cymbopogon winterianus

Special notes

When selecting the *Citronella oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Citronella oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Citronella oil	140121-115623	31550	40	not required
Taoasis	Citronella oil	1291021-122850	31840	40	not required
Taoasis	Citronella oil	1794-373	33728	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Citronella oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *Citronella oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Citronella oil	140121-115623	31550 [†]	20
Taoasis	Citronella oil	1291021-122850	31840 [†]	20
Taoasis	Citronella oil	108494-126305	32798	40
Taoasis	Citronella oil	108494-126776	33047	40

- 22 711 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 1 spectra from 1 *Apo-Ident* customers from 1 batches from the substance/substance group *Citronella oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Citronella oil	5211028-120284	1

- 3192 spectra from 304 *Apo-Ident* customers from a total of 1658 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Citronella oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Citronella oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	120	0	22 711
Type C	0	1	0	3192

The substance/substance group *Citronella oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 95.0000 %)
Type C	100.0000 % (> 98.4407 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31550	31550	0.00	37.73
31840	31840	0.00	41.74
33728	33728	0.00	37.77

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Citrus oil / Lavender oil / Clary sage oil / Peti...
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30961-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil; Citrus oil; Clary sage oil; Hyssop oil; Lavender oil; Lavendula officinalis oleum; Oleum citrus aurantium; Oleum hyssopus officinalis; Oleum lavendula officinalis; Oleum salviae sclarea; Petitgrain oil

Special notes

When selecting the *Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Lavender oil	L16812DN-108661BAG90451	30961	40	not required

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Lavender oil	110186BAG90451	30962	40	not required
Taoasis	Hyssop oil	31823-118643	31649	40	not required
Taoasis	Clary sage oil	2098-115222	31669	40	not required
Taoasis	Lavender oil	22015-121843	31756	40	not required
Taoasis	Petitgrain oil	26056-120864	31778	40	not required
Taoasis	Hyssop oil	33065-122200	31782	40	not required
Apotheker Bau...	Hyssop oil	13.201345120.16.02	31850	40	not required
Apotheker Bau...	Petitgrain oil	6.56267.16.02	31852	40	not required
Taoasis	Hyssop oil	1616260-125049	32980	40	not required
Taoasis	Petitgrain oil	4031015-126048	33024	40	not required
Taoasis	Clary sage oil	1834-127059	33033	40	not required
Primavera	Citus oil	00438D25	33760	40	not required
Primavera	Lavender oil	00684K25	33804	40	not required
Taoasis	Citus oil	788	33816	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 600 spectra of 15 reference samples from the substance/substance group *Citus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 15 different batches.
- 18 149 spectra from a total of 424 batches from further 146 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 752 spectra from 23 reference samples of the substance/substance group *Citus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil*.
- These include spectra of independent samples from 11 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Hyssop oil	zu erfassen	30537	40
Taoasis	Petitgrain oil	zu erfassen	30653	40
Taoasis	Clary sage oil	487962-107805BAG90451	30749	40
Taoasis	Petitgrain oil	29551-105402BAG90451	30752	40
Taoasis	Hyssop oil	31823-118643	31649 [†]	20
Taoasis	Clary sage oil	2098-115222	31669 [†]	20
Taoasis	Lavender oil	22015-121843	31756 [†]	20
Taoasis	Petitgrain oil	26056-120864	31778 [†]	20
Taoasis	Hyssop oil	33065-122200	31782 [†]	20
Apotheker Bauer & Cie	Hyssop oil	13.201345120.16.02	31850 [†]	20
Apotheker Bauer & Cie	Petitgrain oil	6.56267.16.02	31852 [†]	20
Taoasis	Clary sage oil	2098-115222	32797	40
Apotheker Bauer & Cie	Hyssop oil	13.201604082.17.01	32923	40
Taoasis	Lavender oil	2/2015-125808I09	33015	40
Taoasis	Lavender oil	22015-125925I10	33016	40
Taoasis	Petitgrain oil	4031015-126048	33025	40
Taoasis	Citrus oil	1510-135	33817	40
Taoasis	Clary sage oil	10140-145	33871	40
Taoasis	Clary sage oil	10140-145	33943	40
Primavera	Lavender oil	00793A26	33953	40
Taoasis	Petitgrain oil	4031015-147	34108	40
Taoasis	Hyssop oil	1616260-1982	34110	40
Taoasis	Lavender oil	2/2015-125808I09	not required	12

- 22 079 spectra from a total of 531 batches from further 198 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 10 spectra from 6 *Apo-Ident* customers from 10 batches from the substance/substance group *Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil*.
- These include spectra of independent samples from 9 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Hyssop oil	31095-105392	1
Taoasis/Noweda	Lavender oil	16716-126380/11	1
Taoasis	Lavender oil	22015-125927	1
Bombastus	Lavender oil	300666	1
Taoasis	Lavender oil	34/151/16	1
Taoasis	Lavender oil	4651010	1
Bergland	Lavender oil	K502790	1
Taoasis	Clary sage oil	2098-115222	1
Taoasis	Clary sage oil	2098-119159	1
Taoasis	Clary sage oil	487962-106655	1

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

- 3183 spectra from 303 *Apo-Ident* customers from a total of 1649 batches from a further 119 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	600	0	18 149
Type B	0	742	10	22 079
Type C	0	10	0	3183

The substance/substance group *Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9531 %)	100.0000 % (> 99.0000 %)
Type B	100.0000 % (> 99.9528 %)	98.6702 % (> 98.2713 %)
Type C	100.0000 % (> 98.3651 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30961	30961	0.00	53.04
30962	30962	0.00	76.46
31649	31649	0.00	23.99
31669	31669	0.00	50.87

continued on the next page

continued from previous page

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31756	31756	0.00	78.86
31778	31778	0.00	72.39
31782	31782	0.00	25.64
31850	31850	0.00	29.03
31852	31852	0.00	46.10
32980	32980	0.00	18.91
33024	33024	0.00	60.95
33033	33033	0.00	51.93
33760	33760	0.00	23.85
33804	33804	0.00	81.13
33816	33816	0.00	29.27

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Clove oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31270-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Clove oil; Caryophylli floris aetheroleum; Oleum caryophylli

Special notes

When selecting the *Clove oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Clove oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Bombastus	Clove oil	279132	31035	40	not required
Caelo	Clove oil	12380605	31270	38	not required
Taoasis	Clove oil	2891012-116936	31557	40	not required
Taoasis	Clove oil	1426-120481	31683	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 158 spectra of 4 reference samples from the substance/substance group *Clove oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 591 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 122 spectra from 5 reference samples of the substance/substance group *Clove oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	Clove oil	12380605	31270 [†]	2
Taoasis	Clove oil	2891012-116936	31557 [†]	20
Taoasis	Clove oil	1426-120481	31683 [†]	20
Taoasis	Clove oil	1598-124689	32938	40
Taoasis	Clove oil	1598-124689	32940	40

- 22 709 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 3 spectra from 3 *Apo-Ident* customers from 3 batches from the substance/substance group *Clove oil*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Clove oil	15119304	1
Caelo	Clove oil	15316103	1
Primavera	Clove oil	47G24	1

- 3190 spectra from 304 *Apo-Ident* customers from a total of 1656 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Clove oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Clove oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	158	0	18 591
Type B	0	122	0	22 709
Type C	0	3	0	3190

The substance/substance group *Clove oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2025 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 95.0820 %)
Type C	100.0000 % (> 98.3847 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31035	31035	0.00	468.50
31270	31270	0.00	465.69
31557	31557	0.00	442.18
31683	31683	0.00	450.49

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Contramarum aroma**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31007-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Contramarum aroma; Contramarum flavour

Special notes

When selecting the *Contramarum aroma* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Contramarum aroma*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Contramarum aroma	13141605	31007	40	1403096
Caelo	Contramarum aroma	13411002	31289	40	not required
Caelo	Contramarum aroma	13411007	31398	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Contramarum aroma*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 159 spectra from 4 reference samples of the substance/substance group *Contramarum aroma*.
- These include spectra of independent samples from 4 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	Contramarum aroma	11073903	30342	39
Caelo	Contramarum aroma	162671	32695	40
Caelo	Contramarum aroma	172230	33701	40
Caelo	Contramarum aroma	181067	34066	40

- 22 672 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 61 spectra from 31 *Apo-Ident* customers from 38 batches from the substance/substance group *Contramarum aroma*.
- These include spectra of independent samples from 36 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Phoenix Goettin	Contramarum aroma	70401432001	1
Caelo	Contramarum aroma	2041515	1
Caelo	Contramarum aroma	1315Q-03148	1
Caelo	Contramarum aroma	13411007	4
Caelo	Contramarum aroma	13411002	1
Caelo	Contramarum aroma	13411011	1
Caelo	Contramarum aroma	14320005	1
Caelo	Contramarum aroma	134111002	1
Caelo	Contramarum aroma	143200009	1
Caelo	Contramarum aroma	14320006	3
VDL;02.01.15;6,42EUR	Contramarum aroma	14320001	1
Caelo	Contramarum aroma	14320001	1
VDL; 18.07.15; 6,56EUR	Contramarum aroma	14320005	1
Caelo	Contramarum aroma	14320009	1
Caelo	Contramarum aroma	14320015	3
Caelo	Contramarum aroma	15113003	2
Caelo	Contramarum aroma	15113002	2
Caelo	Contramarum aroma	15113011	1
Caelo	Contramarum aroma	16083801	6
Fagron	Contramarum aroma	16083801	1
Caelo	Contramarum aroma	16083802	2
Caelo	Contramarum aroma	16083804	1
Caelo	Contramarum aroma	16083806	1
Caelo	Contramarum aroma	16267101	1
Caelo	Contramarum aroma	16267102	1
Caelo	Contramarum aroma	16267103	1
Caelo	Contramarum aroma	169267103	1
Caelo	Contramarum aroma	17223002	1
Caelo	Contramarum aroma	21051508	1
Caelo	Contramarum aroma	5315A-03148	1
Caelo	Contramarum aroma	7071207	1
Caelo	Contramarum aroma	7091302	1
Caelo	Contramarum aroma	11378804	1
Fiebig	Contramarum aroma	11378810	1
Gehe 08.05.2013	Contramarum aroma	12201407	1
Caelo	Contramarum aroma	12201407	2
Gehe	Contramarum aroma	12201404	1
Gehe 09.04.2013	Contramarum aroma	12201410	1
Caelo	Contramarum aroma	13141602	1
Cael	Contramarum aroma	13141603	1
Caelo	Contramarum aroma	13141606	3
Caelo	Contramarum aroma	13141608	2

- 3132 spectra from 302 *Apo-Ident* customers from a total of 1621 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Contramarum aroma* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Contramarum aroma* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	151	8	22 672
Type C	0	47	14	3132

The substance/substance group *Contramarum aroma* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9529 %)	94.9686 % (> 93.0818 %)
Type C	100.0000 % (> 98.3580 %)	77.0492 % (> 72.1311 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31007	31007	0.00	57.26
31289	31289	0.00	54.21
31398	31398	0.00	57.06

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Cumin oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30449-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Cumin oil; Oleum cuminum cyminum

Special notes

When selecting the *Cumin oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Cumin oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Cumin oil	130910-113031BAG90451	31141	40	not required
Taoasis	Cumin oil	51764-124375	32267	40	not required
Taoasis	Cumin oil	51764-124375	32904	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Cumin oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 629 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 160 spectra from 4 reference samples of the substance/substance group *Cumin oil*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Cumin oil	348912-71369	30449	40
Taoasis	Cumin oil	51764-124375	32266	40
Taoasis	Cumin oil	2016109952-127656	33841	40
Taoasis	Cumin oil	2016109952-128855	33873	40

- 22 671 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 2 spectra from 2 *Apo-Ident* customers from 2 batches from the substance/substance group *Cumin oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Cumin oil	24081-106027	1
Taoasis	Cumin oil	121217-111221	1

- 3191 spectra from 304 *Apo-Ident* customers from a total of 1657 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Cumin oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Cumin oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	120	40	22 671
Type C	0	1	1	3191

The substance/substance group *Cumin oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9529 %)	75.0000 % (> 73.1250 %)
Type C	100.0000 % (> 98.3987 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31141	31141	0.00	162.59
32267	32267	0.00	164.56
32904	32904	0.00	178.99

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Cypress oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30457-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Cypress oil; Oleum cupressus sempervirens

Special notes

When selecting the *Cypress oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Cypress oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Cypress oil	732-119820	31656	40	not required
Taoasis	Cypress oil	43953-129229	33623	40	not required
Primavera	Cypress oil	00177G25	33802	40	not required
Primavera	Cypress oil	00403L25	33954	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Cypress oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 589 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 4 reference samples of the substance/substance group *Cypress oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Cypress oil	457749-105281	30457	40
Taoasis	Cypress oil	732-119820	31656 [†]	20
Taoasis	Cypress oil	107527-125443	32806	40
Taoasis	Cypress oil	107527-125443	33045	40

- 22 691 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 13 spectra from 2 *Apo-Ident* customers from 10 batches from the substance/substance group *Cypress oil*.
- These include spectra of independent samples from 10 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Cypress oil	487749-108547	1
Taoasis	Cypress oil	487749-107790	1
Taoasis	Cypress oil	514085-111139	1
Taoasis	Cypress oil	1039-121894	3
Taoasis	Cypress oil	341012-123298	1
Taoasis	Cypress oil	1039-122434	1
Taoasis	Cypress oil	33720-115882	1
Taoasis	Cypress oil	33720-118623	1
Taoasis	Cypress oil	33720-118895	2
Taoasis	Cypress oil	107527-125443	1

- 3180 spectra from 304 *Apo-Ident* customers from a total of 1649 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Cypress oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Cypress oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	123	17	22 691
Type C	0	12	1	3180

The substance/substance group *Cypress oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9530 %)	87.8571 % (> 85.7143 %)
Type C	100.0000 % (> 98.3631 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31656	31656	0.00	69.53
33623	33623	0.00	69.05
33802	33802	0.00	53.59
33954	33954	0.00	50.17

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Dermatop[®] base ointment**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31060-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Dermatop[®] base ointment

Special notes

When selecting the *Dermatop[®] base ointment* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Dermatop[®] base ointment*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Sanofi aventis	Dermatop [®] base o...	3F052A	31060	40	not required
Sanofi aventis	Dermatop [®] base o...	5F080A	31717	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Dermatop[®] base ointment*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Dermatop[®] base ointment*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Sanofi aventis	Dermatop [®] base ointment	5F080A	31717 [†]	20
Sanofi aventis	Dermatop [®] base ointment	5F089A	32718	40
Sanofi aventis	Dermatop [®] base ointment	7F106A	33965	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 14 spectra from 3 *Apo-Ident* customers from 12 batches from the substance/substance group *Dermatop[®] base ointment*.
- These include spectra of independent samples from 11 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Sanofi/Noweda	Dermatop [®] base ointment	4F075A	2
Sanofi/Gehe	Dermatop [®] base ointment	5F078A	1
Caelo	Dermatop [®] base ointment	5F080A	1
Sanofi/Noweda	Dermatop [®] base ointment	5F081A	1
Sanofi/Noweda	Dermatop [®] base ointment	5F075A	1
Sanofi/Noweda	Dermatop [®] base ointment	5F076A	1
Sanofi/Noweda	Dermatop [®] base ointment	5F083A	1
Sanofi/Noweda	Dermatop [®] base ointment	5F084A	1
Sanofi/Noweda	Dermatop [®] base ointment	5F086A	1
Sanofi/Noweda	Dermatop [®] base ointment	6F091A	1
Sanofi/Noweda	Dermatop [®] base ointment	3F059A	1
Sanofi/Noweda	Dermatop [®] base ointment	4F067A	2

- 3179 spectra from 304 *Apo-Ident* customers from a total of 1647 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Dermatop[®] base ointment* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Dermatop[®] base ointment* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	100	0	22 731
Type C	0	14	0	3179

The substance/substance group *Dermatop[®] base ointment* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3627 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several

new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31060	31060	0.00	16.94
31717	31717	0.00	23.60

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Dexeryl®**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31512-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Dexeryl®

Special notes

When selecting the *Dexeryl®* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Dexeryl®*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Pierre Fabre	Dexeryl®	G99016	31512	40	not required
Pierre Fabre	Dexeryl®	G00419	31630	40	not required
Pierre Fabre	Dexeryl®	G00650	33635	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Dexeryl*[®]. These samples are listed above in the section *calibration samples*. The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this *chemometric model*, the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in *Appendix A*. The samples in *Appendix A* were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *Dexeryl*[®].
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Pierre Fabre	<i>Dexeryl</i> [®]	G99016	31512 [†]	20
Pierre Fabre	<i>Dexeryl</i> [®]	G00419	31630 [†]	20
Pierre Fabre	<i>Dexeryl</i> [®]	G00623	32762	40
Pierre Fabre	<i>Dexeryl</i> [®]	G00658	33945	40

- 22 711 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this *chemometric model*, the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in *Appendix B*.

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 2 spectra from 1 *Apo-Ident* customers from 2 batches from the substance/substance group *Dexeryl*[®].
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
PiereFabre/Now	Dexeryl [®]	G00577	1
Pierre Fabre /Noweda	Dexeryl [®]	G00606	1

- 3191 spectra from 304 *Apo-Ident* customers from a total of 1657 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Dexeryl*[®] can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Dexeryl*[®] and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	120	0	22 708
Type C	0	2	0	3191

The substance/substance group *Dexeryl*[®] can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9535 %)	100.0000 % (> 95.0000 %)
Type C	100.0000 % (> 98.3987 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31512	31512	0.00	22.47
31630	31630	0.00	30.40
33635	33635	0.00	25.74

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Dichloroacetic acid**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30417-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Dichloroacetic acid; Acidum dichloroaceticum

Special notes

When selecting the *Dichloroacetic acid* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Dichloroacetic acid*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Merck KGaA	Dichloroacetic a...	S6674841451	31642	40	not required
Merck KGaA	Dichloroacetic a...	S6674841614	32750	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Dichloroacetic acid*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Dichloroacetic acid*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Sigma-Aldrich	Dichloroacetic acid	STBC3539V	30417	40
Merck	Dichloroacetic acid	S6674841451	31642 [†]	20
Merck	Dichloroacetic acid	S7209041737	33794	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Dichloroacetic acid*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Dichloroacetic acid* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Dichloroacetic acid* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	20	80	22 731
Type C	0	0	0	3193

The substance/substance group *Dichloroacetic acid* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9530 %)	20.0000 % (> 17.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31642	31642	0.00	189.27
32750	32750	0.00	188.05

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Dimeticone ointment 10% SR / Optiderm® lotion**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31243-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Dimeticone ointment 10% SR / Optiderm® lotion; Dimeticone ointment 10% SR; Optiderm® lotion;
Unguentum dimeticoni 10% SR

Special notes

When selecting the *Dimeticone ointment 10% SR / Optiderm® lotion* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Dimeticone ointment 10% SR / Optiderm® lotion*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
apomix	Dimeticone ointm...	08B041213	31243	30	not required
Almirall	Optiderm® lotion	341431	31266	40	not required
apomix	Dimeticone ointm...	10A181214	31638	40	not required
Almirall	Optiderm® lotion	511951	31759	40	not required
apomix	Dimeticone ointm...	11A271216	33041	40	not required
Almirall	Optiderm® lotion	706961	33268	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 230 spectra of 6 reference samples from the substance/substance group *Dimeticone ointment 10% SR / Optiderm[®] lotion*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 6 different batches.
- 18 519 spectra from a total of 433 batches from further 149 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 210 spectra from 7 reference samples of the substance/substance group *Dimeticone ointment 10% SR / Optiderm[®] lotion*.
- These include spectra of independent samples from 4 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
apomix	Dimeticone ointment 10% SR	07B140711	30356	30
apomix	Dimeticone ointment 10% SR	05B120412	30553	40
apomix	Dimeticone ointment 10% SR	10A181214	31638 [†]	20
Almirall	Optiderm [®] lotion	511951	31759 [†]	20
Almirall	Optiderm [®] lotion	633771	32757	20
Almirall	Optiderm [®] lotion	706961	33267	40
apomix	Dimeticone ointment 10% SR	10A021117	33870	40

- 22 621 spectra from a total of 542 batches from further 201 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 13 spectra from 3 *Apo-Ident* customers from 13 batches from the substance/substance group *Dimeticone ointment 10% SR / Optiderm® lotion*.
- These include spectra of independent samples from 12 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
apomix/Phönix	Dimeticone ointment 10% SR	1A070217	1
apomix/Holdermann	Dimeticone ointment 10% SR	1A120116	1
apomix/Holdermann	Dimeticone ointment 10% SR	1A150115	1
Apomix	Dimeticone ointment 10% SR	3B180214	1
apomix/Phönix	Dimeticone ointment 10% SR	4A010415	1
apomix/Holdermann	Dimeticone ointment 10% SR	4A080414	1
apomix/Phönix	Dimeticone ointment 10% SR	5A020616	1
apomix/Gehe	Dimeticone ointment 10% SR	6A190614	1
apomix/Phönix	Dimeticone ointment 10% SR	7A170816	1
apomix/Holdermann	Dimeticone ointment 10% SR	8A090915	1
apomix/Holdermann	Dimeticone ointment 10% SR	6A300616	1
apomix/Gehe	Dimeticone ointment 10% SR	10A181214	1
Apomix/Phönix	Dimeticone ointment 10% SR	8A041213	1

- 3180 spectra from 304 *Apo-Ident* customers from a total of 1646 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Dimeticone ointment 10% SR / Optiderm® lotion* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Dimeticone ointment 10% SR / Optiderm® lotion* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	230	0	18 519
Type B	0	210	0	22 619
Type C	0	13	0	3180

The substance/substance group *Dimeticone ointment 10% SR / Optiderm® lotion* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.3913 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 97.1429 %)
Type C	100.0000 % (> 98.3631 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31243	31243	0.00	18.41
31266	31266	0.00	17.65
31638	31638	0.00	20.88
31759	31759	0.00	14.92
33041	33041	0.00	25.66
33268	33268	0.00	13.63

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Dwarf pine oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31542-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Dwarf pine oil; Oleum pini pumilionis; Pini pumilionis aetherolum

Special notes

When selecting the *Dwarf pine oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Dwarf pine oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Dwarf pine oil	586-116867	31542	40	not required
Taoasis	Dwarf pine oil	25599-120877	31689	40	not required
Taoasis	Dwarf pine oil	8783-127273	32937	40	not required
Taoasis	Dwarf pine oil	11148-1980	34109	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Dwarf pine oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 589 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 85 spectra from 4 reference samples of the substance/substance group *Dwarf pine oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Dwarf pine oil	586-116867	31542 [†]	20
Taoasis	Dwarf pine oil	25599-120877	31689 [†]	20
Taoasis	Dwarf pine oil	8783-125163	32811	40
Taoasis	Dwarf pine oil	8783-125163	not required	5

- 22 746 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 3 spectra from 1 *Apo-Ident* customers from 3 batches from the substance/substance group *Dwarf pine oil*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Dwarf pine oil	6635-122703	1
Taoasis	Dwarf pine oil	8238-124968	1
Taoasis	Dwarf pine oil	8238-126318	1

- 3190 spectra from 304 *Apo-Ident* customers from a total of 1656 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Dwarf pine oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Dwarf pine oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	67	18	22 746
Type C	0	1	2	3190

The substance/substance group *Dwarf pine oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9531 %)	78.8235 % (> 75.2941 %)
Type C	100.0000 % (> 98.3847 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31542	31542	0.00	32.71
31689	31689	0.00	32.91
32937	32937	0.00	36.10
34109	34109	0.00	34.26

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Ethanol 70% pure / denatured**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30779-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Ethanol 70% pure / denatured; Ethanolum 70% rein / vergällt

Special notes

When selecting the *Ethanol 70% pure / denatured* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Ethanol 70% pure / denatured*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Hofmann's	Ethanol 70% pure...	271012	30779	80	1402563
apomix	Ethanol 70% pure...	54A170613	31055	40	not required
Kremer GmbH &...	Ethanol 70% pure...	171211	34101	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 3 reference samples from the substance/substance group *Ethanol 70% pure / denatured*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 589 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 80 spectra from 2 reference samples of the substance/substance group *Ethanol 70% pure / denatured*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Hofmann's	Ethanol 70% pure / denatured	570916	32591	40
apomix	Ethanol 70% pure / denatured	01A020117	33217	40

- 22 751 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 252 spectra from 80 *Apo-Ident* customers from 211 batches from the substance/substance group *Ethanol 70% pure / denatured*.
- These include spectra of independent samples from 208 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
eigen	Ethanol 70% pure / denatured		2
Apomix	Ethanol 70% pure / denatured	6A040213	1
Kremer/Gehe	Ethanol 70% pure / denatured	130116	1
Hirsch	Ethanol 70% pure / denatured	714E-02320	1
A.Pflüger/Noweda	Ethanol 70% pure / denatured	130116	2
Gutenberg-Apotheke	Ethanol 70% pure / denatured	130502003	1
Kremer	Ethanol 70% pure / denatured	130206	1
A. Pflüger/Phönix	Ethanol 70% pure / denatured	130607	1
Pflüger	Ethanol 70% pure / denatured	130607	1
Kemer Phönix	Ethanol 70% pure / denatured	130628	1
Stern-Defektur	Ethanol 70% pure / denatured	130705003	1
Kremer GmbH Wuppertal/G...	Ethanol 70% pure / denatured	130729	1
Noweda/KremerGmbH	Ethanol 70% pure / denatured	130821	1
A. Pflüger/Phönix	Ethanol 70% pure / denatured	130729	1
A. Pflüger/Jenne	Ethanol 70% pure / denatured	131108	1
Pflüger	Ethanol 70% pure / denatured	131108	2
Doppeleiche Apotheke	Ethanol 70% pure / denatured	131227003	1
Kremer/Gehe	Ethanol 70% pure / denatured	140121	1
Doppeleiche Apotheke	Ethanol 70% pure / denatured	140303003	1
Berkel AHK/HWA	Ethanol 70% pure / denatured	1410T31040613/70	1
Post-Apotheke 15.01.2014	Ethanol 70% pure / denatured	15012044	1
Caelo	Ethanol 70% pure / denatured	151008	2
Eigenherstellung	Ethanol 70% pure / denatured	1823	2
Defektur	Ethanol 70% pure / denatured	200114Et0H70	1
Berkel AHK	Ethanol 70% pure / denatured	20042012397	1
A.Pflüger	Ethanol 70% pure / denatured	21111301	1
Hohe-Wart-Apotheke	Ethanol 70% pure / denatured	23072013/1	1
Defektur	Ethanol 70% pure / denatured	270613Et0H	1
Eigenherstellung	Ethanol 70% pure / denatured	2753	2
frischhergestellt	Ethanol 70% pure / denatured		1
Doppeleiche Apotheke	Ethanol 70% pure / denatured	3013A-02320	1
Hofmann	Ethanol 70% pure / denatured	320713	1
Mühlen-Apotheke	Ethanol 70% pure / denatured	3713A-02320	1
Phönix	Ethanol 70% pure / denatured	351213	1
Römer-Apotheke	Ethanol 70% pure / denatured	4314A-03043	1
Hirsch	Ethanol 70% pure / denatured	4413I-02320	1
Belous	Ethanol 70% pure / denatured	4613E-02320	1
Fischar / Sanacorp	Ethanol 70% pure / denatured	7023091	1
Fischar / Noweda	Ethanol 70% pure / denatured	7023111	1
Martinus-Apotheke	Ethanol 70% pure / denatured	A-061213-3D	4
Martinus-Apotheke	Ethanol 70% pure / denatured	A-311013-1D	2
Martinus-Apotheke	Ethanol 70% pure / denatured	E-100114-1D	2
Martinus-Apotheke	Ethanol 70% pure / denatured	E-020314-1D	2
Fagron	Ethanol 70% pure / denatured	Probemessung	1
Eigene Herstellung	Ethanol 70% pure / denatured	SPV7025	1
Noweda	Ethanol 70% pure / denatured	120420	1
phönix,29.12.2012	Ethanol 70% pure / denatured	121130	1
Post-Apotheke	Ethanol 70% pure / denatured	250612	1
Defektur	Ethanol 70% pure / denatured	4712M-02320	1
Plüger/Anzag	Ethanol 70% pure / denatured	32016	1
Hersteller: selbst	Ethanol 70% pure / denatured	12203	1
Eigenherstellung	Ethanol 70% pure / denatured		1
A.Pflüger GmbH & Co.KG/...	Ethanol 70% pure / denatured	130514	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Kremer/Noweda	Ethanol 70% pure / denatured	131011	1
A.Pflüger GmbH & Co.KG/...	Ethanol 70% pure / denatured	131011	1
Heck/Wittich	Ethanol 70% pure / denatured	2113N-02320	1
Euro OTC	Ethanol 70% pure / denatured	220713	1
Hubertus Apotheke	Ethanol 70% pure / denatured	261113	1
Eigenherstellung	Ethanol 70% pure / denatured	4733	1
Fischar / Anzag	Ethanol 70% pure / denatured	7023091	1
Defektur	Ethanol 70% pure / denatured		2
Rondell Apotheke	Ethanol 70% pure / denatured	415M-03043	1
Rondell Apotheke	Ethanol 70% pure / denatured	416M-02320	1
Eigenherstellung	Ethanol 70% pure / denatured	517Q-02320	1
Rondell Apotheke	Ethanol 70% pure / denatured	615E-02320	1
Rondell Apotheke	Ethanol 70% pure / denatured	615E-07145	1
Eigenherstellung	Ethanol 70% pure / denatured	616Q-02320	1
eigene Herstellung	Ethanol 70% pure / denatured	6180115	1
HirschApotheke	Ethanol 70% pure / denatured	10151-02320	1
Eigen/Defektur	Ethanol 70% pure / denatured	1116E-03043	1
Gehe/Pflüger	Ethanol 70% pure / denatured	1204021712	1
Kastner (eigen)	Ethanol 70% pure / denatured	1215E-02320	1
Claudia Lange	Ethanol 70% pure / denatured	1234	1
Brinn- Holtz	Ethanol 70% pure / denatured	1234-04/18	1
Brinn- Holtz	Ethanol 70% pure / denatured	1257-06/17	1
Rosen Apotheke	Ethanol 70% pure / denatured	1317A-07095	1
apomix/Phoenix	Ethanol 70% pure / denatured	12A250116	1
Gehe 27.2.14	Ethanol 70% pure / denatured	140121	1
Gehe-Kremer	Ethanol 70% pure / denatured	140307	1
KremperGmbH & Co. KG	Ethanol 70% pure / denatured	140507	1
Kremer GmbH / Noweda	Ethanol 70% pure / denatured	140627	1
Pflüger/Phoenix	Ethanol 70% pure / denatured	140902	1
Kremer GmbH & Co.KG	Ethanol 70% pure / denatured	140902	2
Doppeleiche Apotheke	Ethanol 70% pure / denatured	140927002	1
Kremer	Ethanol 70% pure / denatured	141027	1
Pflüger	Ethanol 70% pure / denatured	141027	1
Doppeleiche Apotheke	Ethanol 70% pure / denatured	141029011	1
Kremer/Phönix	Ethanol 70% pure / denatured	141127	1
Kremer GmbH / Noweda	Ethanol 70% pure / denatured	141127	2
Doppeleiche Apotheke	Ethanol 70% pure / denatured	141208001	1
Doppeleiche Apotheke	Ethanol 70% pure / denatured	150114010	1
Gehe-Kremer	Ethanol 70% pure / denatured	150209	1
Doppeleiche Apotheke	Ethanol 70% pure / denatured	150219007	1
Kremer/Phönix	Ethanol 70% pure / denatured	150327	1
Stiftsapotheke	Ethanol 70% pure / denatured	150420005	1
Doppeleiche Apotheke	Ethanol 70% pure / denatured	150504003	1
Kremer GmbH & Co.KG	Ethanol 70% pure / denatured	150602	1
Apotheke im Forum	Ethanol 70% pure / denatured	150707	1
Klenk	Ethanol 70% pure / denatured	150730	1
Kremer/Phoenix	Ethanol 70% pure / denatured	150730	1
Caelo	Ethanol 70% pure / denatured	150730	1
Kremer/Phönix	Ethanol 70% pure / denatured	151123	1
Eigenherstellung	Ethanol 70% pure / denatured	1515I-02320	1
Libellen Apotheke	Ethanol 70% pure / denatured	1515Q-02320	1
Apomix/Alliance Healthc...	Ethanol 70% pure / denatured	15A170215	1
Caelo	Ethanol 70% pure / denatured	160330	1
Kremer/Gehe	Ethanol 70% pure / denatured	160414	2
Kremer	Ethanol 70% pure / denatured	160414	1
Dreikönigen Apotheke	Ethanol 70% pure / denatured	16049021	1
Kremer GmbH	Ethanol 70% pure / denatured	160512	1
Kremer	Ethanol 70% pure / denatured	161017	1
Pflüger GmbH & Co.KG	Ethanol 70% pure / denatured	170116	1
Kremer	Ethanol 70% pure / denatured	170208	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Pflüger/Gehe	Ethanol 70% pure / denatured	170619	1
n.a.	Ethanol 70% pure / denatured	1707918-a	1
eigene Herstellung	Ethanol 70% pure / denatured	1712904-A	1
Defektur	Ethanol 70% pure / denatured	1716M-03043	1
Rondell Apotheke	Ethanol 70% pure / denatured	1814Q-07145	1
Caelo	Ethanol 70% pure / denatured	20140403	1
Eigenherstellung	Ethanol 70% pure / denatured	2015E-02320	1
Apotärze	Ethanol 70% pure / denatured	2015I-02320	1
Defektur	Ethanol 70% pure / denatured	2015Q-03043	1
Rezeptur Apotheke	Ethanol 70% pure / denatured	2016Q-03043	1
Eigenherstellung	Ethanol 70% pure / denatured	2124	1
Eigenherstellung	Ethanol 70% pure / denatured	2216A-02320	1
Defektur	Ethanol 70% pure / denatured	23/14	1
kremer gmbh	Ethanol 70% pure / denatured	23081605	1
Eigenherstellung	Ethanol 70% pure / denatured	2314I-02320	1
HirschApotheke	Ethanol 70% pure / denatured	2315I-02320	1
Rosenapotheke	Ethanol 70% pure / denatured	2317E-07095	1
Eigen/Defektur	Ethanol 70% pure / denatured	2516M-03043	1
Defektur	Ethanol 70% pure / denatured	2517A-07097	1
Rondell Apotheke	Ethanol 70% pure / denatured	2615I-07145	1
Rondell Apotheke	Ethanol 70% pure / denatured	2616M-02320	1
Defektur	Ethanol 70% pure / denatured	2617I-07097	1
Fischar	Ethanol 70% pure / denatured	27031501	1
Eigen/Defektur	Ethanol 70% pure / denatured	2716M-03043	1
Apotheke im Forum	Ethanol 70% pure / denatured	300517	1
Kastner (eigen)	Ethanol 70% pure / denatured	3114E-03043	1
Rondell Apotheke	Ethanol 70% pure / denatured	3414M-02320	1
Rondell Apotheke	Ethanol 70% pure / denatured	3414M-07145	1
Eigenherstellung	Ethanol 70% pure / denatured	3416E-02320	1
Eigen/Defektur	Ethanol 70% pure / denatured	3416E-03043	1
Apomix / Alliance Healt...	Ethanol 70% pure / denatured	37A110416	1
Fagron	Ethanol 70% pure / denatured	390414	1
Met/ Noweda	Ethanol 70% pure / denatured	3815M-03043	1
Engel/Engel	Ethanol 70% pure / denatured	39-28715	1
Rondell Apotheke	Ethanol 70% pure / denatured	4015A-03043	1
Wolf-Apotheke / Defektur	Ethanol 70% pure / denatured	40722/51Hk	1
Fagron	Ethanol 70% pure / denatured	40929/52La	1
Apomix / Phönix	Ethanol 70% pure / denatured	41A180515	1
Eigenherstellung	Ethanol 70% pure / denatured	4215I-02320	1
Rezeptur	Ethanol 70% pure / denatured	4215Q-03043	1
eigene Herstellung	Ethanol 70% pure / denatured	431140218	1
eigene Herstellung	Ethanol 70% pure / denatured	4315Q-07016	1
Eigenherstellung	Ethanol 70% pure / denatured	4414A-02320	1
HirschApotheke	Ethanol 70% pure / denatured	4614A-02320	1
Defektur	Ethanol 70% pure / denatured	4714E-03043	1
Eigenherstellung	Ethanol 70% pure / denatured	4615	1
Eigenherstellung	Ethanol 70% pure / denatured	4715M	1
Defektur	Ethanol 70% pure / denatured	50427/52Wlo	1
Wolf-Apotheke / Defektur	Ethanol 70% pure / denatured	51019/51Wlo	1
Eigenherstellung	Ethanol 70% pure / denatured	5214A-02320	1
Apomix/Alliance Healthc...	Ethanol 70% pure / denatured	56a020715	1
Hoffmann & Sommer GmbH ...	Ethanol 70% pure / denatured	580916	1
Apomix/Noweda	Ethanol 70% pure / denatured	58A020715	1
Caelo	Ethanol 70% pure / denatured	60307/50La	1
Berkel AHK	Ethanol 70% pure / denatured		1
Claudia Lange	Ethanol 70% pure / denatured	61010/51La	1
Claudia Lange	Ethanol 70% pure / denatured	61107/50La	1
Apotheke	Ethanol 70% pure / denatured	70105/51Dre	1
Otto Fischar GmbH & Co...	Ethanol 70% pure / denatured	7024021	1
Otto Fischar	Ethanol 70% pure / denatured	7023111	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Caelo	Ethanol 70% pure / denatured	7023111	1
Fischar / Noweda	Ethanol 70% pure / denatured	7024021	1
Fischar	Ethanol 70% pure / denatured	7024051	1
otto Fischer/Noweda	Ethanol 70% pure / denatured	7024121	1
Anzag/ Otto Fischar Gmb...	Ethanol 70% pure / denatured	7024051	1
Otto Fischar GmbH & Co...	Ethanol 70% pure / denatured	7024071	1
Otto Fischar GmbH/ Amzag	Ethanol 70% pure / denatured	7024091	1
Otto Fischer / Noweda	Ethanol 70% pure / denatured	7025031	1
Otto Fischar GmbH & Co...	Ethanol 70% pure / denatured	7025031	2
Caelo	Ethanol 70% pure / denatured	7025081	1
Fischar/Phönix	Ethanol 70% pure / denatured	7026111	1
Fischar/Phönix	Ethanol 70% pure / denatured	7026031	1
Fischar / Alliance Hea...	Ethanol 70% pure / denatured	7026051	1
Fischar/Phönix	Ethanol 70% pure / denatured	7026092	1
Wolf-Apotheke / Defektur	Ethanol 70% pure / denatured	Bulk	1
Martinus-Apotheke	Ethanol 70% pure / denatured	E-050515-1D	1
Hustadt A	Ethanol 70% pure / denatured		1
Martinus-Apotheke	Ethanol 70% pure / denatured	E-080514-1D	1
Martinus-Apotheke	Ethanol 70% pure / denatured	E-110716-1D	1
Martinus-Apotheke	Ethanol 70% pure / denatured	E-021116-1D	1
	Ethanol 70% pure / denatured		1
Martinus-Apotheke	Ethanol 70% pure / denatured	E-121214-1D	1
Martinus-Apotheke	Ethanol 70% pure / denatured	E-180714-1D	1
Eigene Herstellung	Ethanol 70% pure / denatured	SP70140	1
eigene herstellung	Ethanol 70% pure / denatured	SPV7026	1
Eigene Herstellung	Ethanol 70% pure / denatured	SPV7029	1
Eigenherstellung	Ethanol 70% pure / denatured	5131213	1
Eigenproduktion	Ethanol 70% pure / denatured	714Q-02320	2
Sanacorp	Ethanol 70% pure / denatured	130124	1
Eigenherstellung	Ethanol 70% pure / denatured	1313I-02320	1
Gutenberg-Apotheke	Ethanol 70% pure / denatured	1813M-02320	1
Eigenherstellung	Ethanol 70% pure / denatured	2613I-03043	1
Eigenherstellung	Ethanol 70% pure / denatured	3413I-02320	1
Eigenherstellung	Ethanol 70% pure / denatured	4513I-02320	1
Caelo	Ethanol 70% pure / denatured		1
Caelo	Ethanol 70% pure / denatured	814E-02320	1
Defektur/Wolf-Apotheke	Ethanol 70% pure / denatured	1234567	1
Stiftsapotheke	Ethanol 70% pure / denatured	13052016	1
Doppeleiche Apotheke	Ethanol 70% pure / denatured	140630003	1
Hustadt Apotheke	Ethanol 70% pure / denatured	141212	1
Post-Apotheke	Ethanol 70% pure / denatured	1414A-02320	1
Doppeleiche Apotheke	Ethanol 70% pure / denatured	150126002	1
Hustadt apotheke	Ethanol 70% pure / denatured	150618	1
Caelo	Ethanol 70% pure / denatured	1715E-02320	1
Rondell	Ethanol 70% pure / denatured	1715E-07145	1
Caelo	Ethanol 70% pure / denatured	19151-02320	1
Caelo	Ethanol 70% pure / denatured	2614E-03043	1
sittig	Ethanol 70% pure / denatured	310316	1
Wolf-Apotheke	Ethanol 70% pure / denatured	40331/52La	1
Wolf-Apotheke / Defektur	Ethanol 70% pure / denatured	41125/52Hk	1
Defektur/Wolf-Apotheke	Ethanol 70% pure / denatured	41211/51Hk	1
Defektur	Ethanol 70% pure / denatured	50427/1La	1
Defektur/Wolf-Apotheke	Ethanol 70% pure / denatured	50714/51Hk	1
Defektur/Wolf-Apotheke	Ethanol 70% pure / denatured	50714/53Hk	1
Defektur	Ethanol 70% pure / denatured	60209/51Hk	1
Defektur / Wolf-Apotheke	Ethanol 70% pure / denatured	60902/50Hk	1
Claudia Lange	Ethanol 70% pure / denatured	60919/50La	1
Purren Apotheke	Ethanol 70% pure / denatured	17043	1
Hofmann Sommer	Ethanol 70% pure / denatured	320713	1
eigene Herstellung	Ethanol 70% pure / denatured	4411312106	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Gutenberg-Apotheke	Ethanol 70% pure / denatured	3412I-03043	1

- 2941 spectra from 293 *Apo-Ident* customers from a total of 1449 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Ethanol 70% pure / denatured* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Ethanol 70% pure / denatured* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	80	0	22 751
Type C	0	235	17	2941

The substance/substance group *Ethanol 70% pure / denatured* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9531 %)	100.0000 % (> 92.5000 %)
Type C	100.0000 % (> 98.3570 %)	93.2540 % (> 92.0635 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30779	30779	0.00	101.63
31055	31055	0.00	83.05
34101	34101	0.00	96.27

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Eucalyptus oil / Rosmary oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30966-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Eucalyptus oil / Rosmary oil; Eucalypti aetheroleum; Eucalyptus oil; Oleum eucalypti; Oleum rosmarini; Rosemary oil; Rosmarini aetheroleum

Special notes

When selecting the *Eucalyptus oil / Rosmary oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Eucalyptus oil / Rosmary oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Eucalyptus oil	527885-109705A	30966	40	not required
Taoasis	Rosemary oil	3231031-117639	31551	40	not required

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Rosemary oil	1921019-120666	31691	40	not required
Taoasis	Eucalyptus oil	7271001-121014	31694	40	not required
Taoasis	Eucalyptus oil	043127A-125381	32903	40	not required
Taoasis	Rosemary oil	1862-126630	32933	40	not required
Taoasis	Rosemary oil	2150-2819	34104	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 280 spectra of 7 reference samples from the substance/substance group *Eucalyptus oil* / *Rosemary oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 7 different batches.
- 18 469 spectra from a total of 432 batches from further 149 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 312 spectra from 10 reference samples of the substance/substance group *Eucalyptus oil* / *Rosemary oil*.
- These include spectra of independent samples from 6 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Eucalyptus oil	T150I11	30384	40
Taoasis	Eucalyptus oil	T51I10	30387	40
Taoasis	Rosemary oil	3231031-117639	31551 [†]	20
Taoasis	Rosemary oil	1921019-120666	31691 [†]	20
Taoasis	Eucalyptus oil	7271001-121014	31694 [†]	20
Taoasis	Rosemary oil	1614-125089	32802	40
Taoasis	Eucalyptus oil	41192-129138	33621	40
Primavera	Eucalyptus oil	00650K25	33808	40
Primavera	Eucalyptus oil	00256A26	33955	40
Taoasis	Rosemary oil	1614-125089	not required	12

- 22 519 spectra from a total of 540 batches from further 201 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 23 spectra from 14 *Apo-Ident* customers from 22 batches from the substance/substance group *Eucalyptus oil* / *Rosemary oil*.
- These include spectra of independent samples from 22 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Euro OTC	Eucalyptus oil	1412030-01	1
Caelo	Eucalyptus oil	15063309	1
Caesar & Loretz GmbH Hi...	Eucalyptus oil	15063312	1
Caesar & Loretz GmbH Hi...	Eucalyptus oil	15400203	1
Caelo	Eucalyptus oil	15400217	1
Bombastus	Eucalyptus oil	303336	1
Bombastus	Eucalyptus oil	305628	1
Taoasis	Eucalyptus oil	43703-122178	1
Taoasis	Eucalyptus oil	43703-122404	1
Taoasis	Eucalyptus oil	52046-124692	1
Caelo	Eucalyptus oil	11324112	1
Caelo	Eucalyptus oil	13106407LOT	1
Anzag, 10.12.2012	Eucalyptus oil	273979	1
Taoasis	Eucalyptus oil	33364-110871	1
Taoasis	Eucalyptus oil	543495-112704	1
Taoasis	Eucalyptus oil	130205-111868	1
Taoasis	Rosemary oil	1526-123355	1
Taoasis	Rosemary oil	26057-119164	1
Taoasis	Rosemary oil	1526-125088	1
Caelo	Rosemary oil	15442017	1
Taoasis	Rosemary oil	26057-122996	1
Bombastus	Rosemary oil	301449	2

- 3170 spectra from 302 *Apo-Ident* customers from a total of 1637 batches from a further 120 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Validation results

The validation runs checked whether the substance/substance group *Eucalyptus oil* / *Rosmary oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Eucalyptus oil* / *Rosmary oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	280	0	18 469
Type B	0	231	81	22 519
Type C	0	20	3	3170

The substance/substance group *Eucalyptus oil* / *Rosmary oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.8571 %)
Type B	100.0000 % (> 99.9529 %)	74.0385 % (> 73.0769 %)
Type C	100.0000 % (> 98.3603 %)	86.9565 % (> 73.9130 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30966	30966	0.00	53.92
31551	31551	0.00	32.08
31691	31691	0.00	27.84
31694	31694	0.00	44.62
32903	32903	0.00	30.74
32933	32933	0.00	31.92
34104	34104	0.00	31.91

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Excipial® almond oil ointment**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30720-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Excipial® almond oil ointment

Special notes

When selecting the *Excipial® almond oil ointment* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F *Zusatz zu den Modellen der zweiten Stufe*

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Excipial® almond oil ointment*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Spirig Pharma	Excipial® almond...	M023	30720	40	not required
Spirig Pharma	Excipial® almond...	M031	31090	40	not required
Spirig Pharma	Excipial® almond...	N012	31278	40	not required
Spirig Pharma	Excipial® almond...	R012	31757	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Excipial[®] almond oil ointment*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 589 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Excipial[®] almond oil ointment*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Spirig Pharma	Excipial [®] almond oil ointment	R012	31757 [†]	20
Spirig Pharma	Excipial [®] almond oil ointment	S021	33209	40
Spirig Pharma	Excipial [®] almond oil ointment	T052	33210	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 22 spectra from 9 *Apo-Ident* customers from 18 batches from the substance/substance group *Excipial[®] almond oil ointment*.
- These include spectra of independent samples from 17 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Galderma/Now	Excipial [®] almond oil ointment	LOTS021	1
Spirig/Noweda	Excipial [®] almond oil ointment	R043	1
Galderma (Spirig Pharma...	Excipial [®] almond oil ointment	R043	1
Galderma	Excipial [®] almond oil ointment	R043	1
Spirig	Excipial [®] almond oil ointment	S011	1
Galderma/Noweda	Excipial [®] almond oil ointment	S012	1
Spirig	Excipial [®] almond oil ointment	S032	1
Spirig	Excipial [®] almond oil ointment	LOTM023	1
Spirig / Gehe	Excipial [®] almond oil ointment	LOTN022	1
Caelo	Excipial [®] almond oil ointment	lotno32	1
Spirig	Excipial [®] almond oil ointment	M023	1
Spirig	Excipial [®] almond oil ointment	N031	1
Spirig	Excipial [®] almond oil ointment	N032	1
Spirig Pharma/Noweda	Excipial [®] almond oil ointment	N022	1
Spirig	Excipial [®] almond oil ointment	R012	1
Spirig /Noweda	Excipial [®] almond oil ointment	R012	1
Spirig /Noweda	Excipial [®] almond oil ointment	r021	1
Spirig/Sanacorp	Excipial [®] almond oil ointment	LOTN011	1
Phoenix	Excipial [®] almond oil ointment	LOTN022	2
Spirig	Excipial [®] almond oil ointment	M032	1
Spirig/AHD	Excipial [®] almond oil ointment	N011	1

- 3171 spectra from 304 *Apo-Ident* customers from a total of 1647 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Excipial[®] almond oil ointment* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Excipial[®] almond oil ointment* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	100	0	22 731
Type C	0	22	0	3171

The substance/substance group *Excipial[®] almond oil ointment* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3605 %)	100.0000 % (> 72.7273 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30720	30720	0.00	31.30
31090	31090	0.00	23.99
31278	31278	0.00	26.08
31757	31757	0.00	23.31

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Fat ointments
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30881-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Fat ointments; Asche Basis[®] fat ointment; Cottonwood ointment; Marjoram ointment; Neribas[®] fat ointment; Protegin[®] XN; Unguentum majoranae; Unguentum Populi

Special notes

When selecting the *Fat ointments* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Fat ointments*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Marjoram ointment	12123001	30881	40	not required
Henry Lamotte	Marjoram ointment	5322500	30926	40	not required
Chiesi	Asche Basis [®] fat...	23014A	31013	40	not required
Caelo	Protegin [®] XN	13081901	31112	40	not required

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Marjoram ointment	14089901	31443	40	not required
Caelo	Protegin [®] XN	13344402	31475	40	not required
Jenapharm	Neribas [®] fat oin...	32032C	31489	40	not required
Chiesi	Asche Basis [®] fat...	41017A	31858	40	not required
Jenapharm	Neribas [®] fat oin...	YY00K7C	31872	40	not required
Caelo	Cottonwood ointm...	161175	32473	40	20160607*
Caelo	Cottonwood ointm...	170572	33130	40	20170307*
Caelo	Cottonwood ointm...	170573	33131	40	20170308*
Chiesi	Asche Basis [®] fat...	53021A	33634	40	not required
Caelo	Protegin [®] XN	17258703	33921	40	not required
Jenapharm	Neribas [®] fat oin...	YY01TF2	33986	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 600 spectra of 15 reference samples from the substance/substance group *Fat ointments*. These samples are listed above in the section *calibration samples*. The reference samples originate from 15 different batches.
- 18 149 spectra from a total of 424 batches from further 146 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this *chemometric model*, the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in *Appendix A*. The samples in *Appendix A* were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 870 spectra from 21 reference samples of the substance/substance group *Fat ointments*.
- These include spectra of independent samples from 15 batches, of which no spectra have been used for the generation of the database.

*The spectra were taken from a part of the same sample which the supplier used for its analysis (identity, purity, content). In this case the manufacturer certificates proof the identity and quality of the sample from those the spectra descend.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	Protegin [®] XN	11022410	30422	40
Caelo	Protegin [®] XN	10079803	30435	40
Caelo	Protegin [®] XN	11196702	30469	40
Caelo	Protegin [®] XN	11196801	30471	40
Caelo	Protegin [®] XN	11196801	30472	70
Caelo	Protegin [®] XN	11196703	30500	40
Caelo	Protegin [®] XN	11196702	30511	40
Chiesi	Asche Basis [®] fat ointment	41017A	31858 [†]	20
Jenapharm	Neribas [®] fat ointment	YY00K7C	31872 [†]	20
Caelo	Marjoram ointment	153533	31990	60
Caelo	Marjoram ointment	153540	31991	60
Caelo	Protegin [®] XN	161645	32340	40
Caelo	Marjoram ointment	160687	32516	40
Chiesi	Asche Basis [®] fat ointment	53021A	32716	40
Jenapharm	Neribas [®] fat ointment	YY011K0	32764	40
Caelo	Marjoram ointment	162575	32872	40
Caelo	Marjoram ointment	162576	32873	40
Chiesi	Asche Basis [®] fat ointment	53021A	32925	40
Caelo	Cottonwood ointment	170534	33162	40
Caelo	Cottonwood ointment	170574	33163	40
Jenapharm	Neribas [®] fat ointment	YY019KR	33290	40

- 21 961 spectra from a total of 531 batches from further 198 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 367 spectra from 120 *Apo-Ident* customers from 123 batches from the substance/substance group *Fat ointments*.
- These include spectra of independent samples from 116 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Chiese	Asche Basis [®] fat ointment	33016A	1
Chiesi	Asche Basis [®] fat ointment	41018A	3
Chiesi/Gehe	Asche Basis [®] fat ointment	41017A	1
Chiesi/Gehe	Asche Basis [®] fat ointment	51020A	1
Chiesi	Asche Basis [®] fat ointment	51020A	1
Chiese	Asche Basis [®] fat ointment	63023A	1
Caelo	Marjoram ointment	312115354001	1
Caelo	Marjoram ointment	2021506	1
Caelo	Marjoram ointment	10111402	1
Caelo	Marjoram ointment	13028401	9
Caelo	Marjoram ointment	13382903	3

continued on the next page

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

continued from previous page

Supplier	Substance	Batch	Spectra
Caelo	Marjoram ointment	13383001	1
Fagron	Marjoram ointment	13A16-T04-000208	1
Caelo	Marjoram ointment	13e10-t04-003550	1
Fagron	Marjoram ointment	13F27-T03-004962	3
Fagron	Marjoram ointment	13F27-To3-004962	1
Caelo	Marjoram ointment	140020101	1
Caelo	Marjoram ointment	14002101	15
Caelo	Marjoram ointment	14089901	8
Gehe	Marjoram ointment	14089901	1
Euro OTC	Marjoram ointment	14089901	1
Caelo	Marjoram ointment	14090002	9
Caelo	Marjoram ointment	14332101	8
Fagron	Marjoram ointment	14090002	1
Caelo	Marjoram ointment	14332202	1
Fagron	Marjoram ointment	14A07-T04-009079	3
Fagron	Marjoram ointment	14A07-T05-009082	2
Fagron	Marjoram ointment	14A07-T06-009085	3
Caelo	Marjoram ointment	15038303	1
Caelo	Marjoram ointment	15038401	2
Caelo	Marjoram ointment	15038501	3
Caelo	Marjoram ointment	15353303	1
Caelo	Marjoram ointment	15354001	5
Fagron	Marjoram ointment	15K10-T10-027644	1
Caelo	Marjoram ointment	15392402	2
Caelo	Marjoram ointment	16068801	1
Caelo	Marjoram ointment	16068901	3
Caelo	Marjoram ointment	16257602	1
Fagron	Marjoram ointment	16E02-T02-032846	1
Fagron	Marjoram ointment	16K07-T20-038496	2
Caelo	Marjoram ointment	16K07-T20-038496	1
Caelo	Marjoram ointment	27031505	1
Henry Lamotte/Sanacorp	Marjoram ointment	5563400	4
Henry Lamotte Oils GmbH	Marjoram ointment	5563400	1
Caelo	Marjoram ointment	2011405	1
Caelo	Marjoram ointment	7121203	1
Caelo	Marjoram ointment	8031304	1
Caelo	Marjoram ointment	11061201	1
Caelo	Marjoram ointment	11274603	2
Caelo	Marjoram ointment	11327501	2
Gehe	Marjoram ointment	11327501	1
Caelo	Marjoram ointment	12018001	7
Phönix 06.10.2012	Marjoram ointment	12018001	1
Phönix	Marjoram ointment	12018001	1
Caelo	Marjoram ointment	12018101	3
Caelo	Marjoram ointment	12081002	6
Fagron	Marjoram ointment	12081002	1
Sanacorp	Marjoram ointment	12081002	1
Caelo	Marjoram ointment	12123001	8
Euro OTC	Marjoram ointment	12123001	1
Caelo	Marjoram ointment	12123101	2
Caelo	Marjoram ointment	1225402	1
Caelo	Marjoram ointment	12295402	7
Fagron	Marjoram ointment	12D24-T01	2
Phönix 10.12.12	Marjoram ointment	12D24-T02	1
Caelo	Marjoram ointment	13028301	2
Caelo	Marjoram ointment	13028302	3
Jenne	Marjoram ointment	13028302	2
Phoenix	Marjoram ointment	13028401	1
Cealo	Marjoram ointment	13028401	2
Gehe	Marjoram ointment	13028401	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Caelo	Marjoram ointment	13051304	1
Caelo	Marjoram ointment	13095401	5
Caelo	Marjoram ointment	13307102	6
Sanacorp	Marjoram ointment	13095401	1
Gehe	Marjoram ointment	13307102	1
Caelo	Marjoram ointment	13382901	1
Caelo	Marjoram ointment	13382902	2
Fagron	Marjoram ointment	13A16-To4-000208	1
Fagron	Marjoram ointment	13A16-T02	1
Caelo	Marjoram ointment	1513E-02536	1
Caelo	Marjoram ointment	15101205	1
Caelo	Marjoram ointment	20290312-1	1
Erika apotheke	Marjoram ointment	5067400	1
Henry Lamotte Oils GmbH...	Marjoram ointment	5067400	1
Lamotte / Gehe	Marjoram ointment	5322500	1
Lamotte/ Noweda	Marjoram ointment	5322500	1
Noweda	Marjoram ointment	5322500	1
Henry Lamotte Oils GmbH	Marjoram ointment	5322500	1
Henry Lamotte Oils GmbH...	Marjoram ointment	5322500	1
Caelo	Neribas [®] fat ointment	yy00k7C	1
GP Grenzach Produktions...	Neribas [®] fat ointment	YY00K7C	1
/ AHD	Neribas [®] fat ointment	YY016LP	2
Jenapharm/Ahd	Neribas [®] fat ointment	YYo11Ko	1
Jenapharm/Phönix	Neribas [®] fat ointment	YY00K7C	1
Phoenix	Neribas [®] fat ointment	YY00K7C	1
Caelo	Protegin [®] XN	14220202	4
Caelo	Protegin [®] XN	14220205	6
Caelo	Protegin [®] XN	14220206	1
Caelo	Protegin [®] XN	14220208	4
Caelo	Protegin [®] XN	14220209	3
Caelo	Protegin [®] XN	14341301	3
Caelo	Protegin [®] XN	14341302	2
Caelo	Protegin [®] XN	15252702	1
Caelo	Protegin [®] XN	15252703	4
Caelo	Protegin [®] XN	16026701	5
Gehe	Protegin [®] XN	16026701	2
Caelo	Protegin [®] XN	16026702	3
Caelo	Protegin [®] XN	16026703	2
Caelo	Protegin [®] XN	16026704	1
Caelo	Protegin [®] XN	16026707	2
Caelo	Protegin [®] XN	16164502	1
Caelo	Protegin [®] XN	17258707	1
Caelo	Protegin [®] XN	13081905	12
Caelo	Protegin [®] XN	13081906	7
Caelo	Protegin [®] XN	13344401	8
Caelo	Protegin [®] XN	13344402	11
Caelo	Protegin [®] XN	14220201	1
Caelo	Protegin [®] XN	14220203	2
Caelo	Protegin [®] XN	14220204	3
Caelo	Protegin [®] XN	26031509	1
Caelo	Protegin [®] XN	6012014A	1
Caelo	Protegin [®] XN	3091208	1
Caelo	Protegin [®] XN	8031305	1
Caelo	Protegin [®] XN	11022403	1
Caelo	Protegin [®] XN	11196702	2
Caelo	Protegin [®] XN	11196703	1
Phönix	Protegin [®] XN	11196706	1
Caelo	Protegin [®] XN	11196707	1
Caelo	Protegin [®] XN	11196801	1
Großhandel	Protegin [®] XN	11196801	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Kehr	Protegin [®] XN	11196803	1
Caelo	Protegin [®] XN	11196805	1
Caelo	Protegin [®] XN	12229201	5
Phönix	Protegin [®] XN	12229201	1
Gehe	Protegin [®] XN	12229201	2
Anzag	Protegin [®] XN	12229205	1
Phönix 12.03.2013	Protegin [®] XN	12229205	1
Caelo	Protegin [®] XN	12229209	1
Caelo	Protegin [®] XN	12229210	1
Caelo	Protegin [®] XN	12229211	6
Phönix, 07.11.13	Protegin [®] XN	12229210	1
Gehe	Protegin [®] XN	12229211	1
Großhandel	Protegin [®] XN	12229211	1
Caelo	Protegin [®] XN	122910	1
Phönix	Protegin [®] XN	13081901	3
Caelo	Protegin [®] XN	13081901	6
Caelo	Protegin [®] XN	13081902	6
Caelo	Protegin [®] XN	13081903	1
Caelo	Protegin [®] XN	13081904	2
Großhandel	Protegin [®] XN	13081905	1
Caelo	Protegin [®] XN	13081907	3
Caelo	Protegin [®] XN	13081908	1
Caelo	Protegin [®] XN	20011402	1
Caelo	Protegin [®] XN	25061301	1
	Protegin [®] XN	2711E-03598	1
VDL	Marjoram ointment	14090002	1
Sanacorp, 21.03.17, 5,73EUR	Marjoram ointment	16068602	1
Sanacorp	Marjoram ointment	12295402	1
Henry Lamotte GmbH/Hold...	Marjoram ointment	5067400	1
Caelo	Marjoram ointment	5322500	1

- 2826 spectra from 283 *Apo-Ident* customers from a total of 1536 batches from a further 118 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Fat ointments* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Fat ointments* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	600	0	18 149
Type B	0	870	0	21 921
Type C	0	352	15	2826

The substance/substance group *Fat ointments* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9531 %)	100.0000 % (> 99.0000 %)
Type B	100.0000 % (> 99.9535 %)	100.0000 % (> 99.3103 %)
Type C	100.0000 % (> 98.3569 %)	95.9128 % (> 95.0954 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30881	30881	0.00	33.17
30926	30926	0.00	44.16
31013	31013	0.00	40.85
31112	31112	0.00	41.60
31443	31443	0.00	33.72
31475	31475	0.00	42.39
31489	31489	0.00	47.48
31858	31858	0.00	51.10
31872	31872	0.00	53.10
32473	32473	0.00	32.51
33130	33130	0.00	31.46
33131	33131	0.00	33.52
33634	33634	0.00	48.16
33921	33921	0.00	41.41
33986	33986	0.00	47.73

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Fenistil® Gel**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31045-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Fenistil® Gel

Special notes

When selecting the *Fenistil® Gel* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Fenistil® Gel*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Novartis	Fenistil® Gel	N00927A	31045	40	not required
Novartis	Fenistil® Gel	R02021A	31767	40	not required
GSK Consumer ...	Fenistil® Gel	U02364A	33638	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Fenistil® Gel*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Fenistil® Gel*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Novartis	Fenistil® Gel	R02021A	31767 [†]	20
Euro OTC	Fenistil® Gel	T02415A	32787	40
Euro OTC	Fenistil® Gel	T02416A	32825	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Fenistil® Gel*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Fenistil® Gel* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Fenistil® Gel* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	100	0	22 731
Type C	0	0	0	3193

The substance/substance group *Fenistil® Gel* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31045	31045	0.00	19.38
31767	31767	0.00	22.02
33638	33638	0.00	21.61

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Fennel oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31140-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Fennel oil; Oleum foeniculum vulgare var. dulce

Special notes

When selecting the *Fennel oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Fennel oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Fennel oil	L16813D-111973BAG90451	31140	40	not required
Taoasis	Fennel oil	L35513DN-119762	31693	40	not required
Taoasis	Fennel oil	1806-125039	32795	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Fennel oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Fennel oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Fennel oil	L35513DN-119762	31693 [†]	20
Taoasis	Fennel oil	1806-127213	32952	40
Taoasis	Fennel oil	1861-129086	33991	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Fennel oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Fennel oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Fennel oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	96	4	22 731
Type C	0	0	0	3193

The substance/substance group *Fennel oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	96.0000 % (> 93.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31140	31140	0.00	96.23
31693	31693	0.00	94.51
32795	32795	0.00	19.96

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Ginger oil
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30451-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Ginger oil; Oleum zingiber officinalis; Zingiber officinalis oleum

Special notes

When selecting the *Ginger oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Ginger oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Ginger oil	130114-110686BAG90451	30958	40	not required
Taoasis	Ginger oil	40592A-112718BAG90451	31142	40	not required
Taoasis	Ginger oil	1431020-124656	32270	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Ginger oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 3 reference samples of the substance/substance group *Ginger oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Ginger oil	6454-92725	30451	40
Bombastus	Ginger oil	300690	32569	40
Bombastus	Ginger oil	300690	33014	40

- 22 711 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 1 spectra from 1 *Apo-Ident* customers from 1 batches from the substance/substance group *Ginger oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Ginger oil	6454-92725	1

- 3192 spectra from 304 *Apo-Ident* customers from a total of 1658 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Ginger oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Ginger oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	80	40	22 711
Type C	0	1	0	3192

The substance/substance group *Ginger oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	66.6667 % (> 64.1667 %)
Type C	100.0000 % (> 98.4407 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30958	30958	0.00	21.68
31142	31142	0.00	18.66
32270	32270	0.00	31.21

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Grapefruit oil organic / Orange oil
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30960-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Grapefruit oil organic / Orange oil; Citrus sinensis oleum; Grapefruit oil, organic; Oleum citrus paradisi; Oleum citrus sinensis; Orange oil

Special notes

When selecting the *Grapefruit oil organic / Orange oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Grapefruit oil organic / Orange oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Orange oil	121109-110685BAG90451	30960	40	not required
Taoasis	Grapefruit oil, ...	31564-119383	31634	80	not required

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Orange oil	33732-12088	31679	40	not required
Taoasis	Grapefruit oil, ...	38755-122181	31781	40	not required
Apotheker Bau...	Grapefruit oil, ...	7.99615.16.02	31853	40	not required
Primavera	Orange oil	00112K25	33805	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 280 spectra of 6 reference samples from the substance/substance group *Grapefruit oil organic / Orange oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 6 different batches.
- 18 469 spectra from a total of 433 batches from further 149 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 380 spectra from 11 reference samples of the substance/substance group *Grapefruit oil organic / Orange oil*.
- These include spectra of independent samples from 7 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Grapefruit oil, organic	zu erfassen	30538	40
Taoasis	Grapefruit oil, organic	31564-119383	31634 [†]	40
Taoasis	Orange oil	33732-12088	31679 [†]	20
Taoasis	Grapefruit oil, organic	38755-122181	31781 [†]	20
Apotheker Bauer & Cie	Grapefruit oil, organic	7.99615.16.02	31853 [†]	20
Taoasis	Orange oil	43829C-124974	32790	40
Taoasis	Orange oil	44583A-126780	32902	40
Taoasis	Grapefruit oil, organic	44040A-124468	32979	40

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra
Apotheker Bauer & Cie	Grapefruit oil, organic	7.68413.17.012018-03	33019	40
Primavera	Orange oil	00149A29	33956	40
Taoasis	Orange oil	011855A-1538	34105	40

- 22 451 spectra from a total of 539 batches from further 201 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 12 spectra from 2 *Apo-Ident* customers from 12 batches from the substance/substance group *Grapefruit oil organic / Orange oil*.
- These include spectra of independent samples from 12 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Grapefruit oil, organic	31564-120540	1
Taoasis	Grapefruit oil, organic	40038-123110	1
Taoasis	Grapefruit oil, organic	44040A-124468	1
Taoasis	Orange oil	122650-123643	1
Taoasis	Orange oil	140129-116983	1
Taoasis	Orange oil	140129-118141	1
Taoasis	Orange oil	140129-118420	1
Taoasis	Orange oil	140129-118790	1
Bombastus	Orange oil	302428	1
Taoasis	Orange oil	42861A-121691	1
Taoasis	Orange oil	42861A-122532	1
Taoasis	Orange oil	43829C-125351	1

- 3181 spectra from 304 *Apo-Ident* customers from a total of 1647 batches from a further 120 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Grapefruit oil organic / Orange oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Grapefruit oil organic*

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

/ *Orange oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	280	0	18 469
Type B	0	340	40	22 371
Type C	0	12	0	3181

The substance/substance group *Grapefruit oil organic / Orange oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.8571 %)
Type B	100.0000 % (> 99.9532 %)	89.4737 % (> 88.6842 %)
Type C	100.0000 % (> 98.3637 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30960	30960	0.00	46.14
31634	31634	0.00	44.68
31679	31679	0.00	48.03
31781	31781	0.00	48.12
31853	31853	0.00	39.65
33805	33805	0.00	46.21

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Hydrophilic ointments
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30634-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Hydrophilic ointments; Abitima[®] clinic body cream; Abitima[®] clinic face cream; Allergika[®] Base cream; Asche Basis[®] cream; Asche Basis[®] lotion; Base cream Taoasis; Dermatop[®] base cream; DMS[®] base cream classic; DMS[®] base cream high classic; DMS[®] base cream high classic plus; Eucerinum O/W basis; Eucerinum W/O basis; Excipial[®] hydro cream; Excipial[®] U Lipolotio; Excipial[®] U10 Lipolotio; Hans Karrer Lipocream MicroSilver; Hans Karrer Lipolotion MicroSilver; Linola[®] body milk; Lipoderm[®] lotion; Neribas[®] cream; Thrombocid[®] ointment

Special notes

When selecting the *Hydrophilic ointments* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Hydrophilic ointments*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Intendis	Neribas [®] cream	21076A	30634	40	not required
Spirig Pharma	Lipoderm [®] lotion	K057	30635	40	not required
Chiesi	Asche Basis [®] cream	03117A	30713	40	not required
Spirig Pharma	Excipial [®] hydro ...	I013	30725	40	not required
Spirig Pharma	Excipial [®] U Lipo...	M042	30732	60	not required
Spirig Pharma	Excipial [®] U10 Li...	M021	30733	60	not required
Bene	Thrombocid [®] oint...	410B113	30772	40	not required
Chiesi	Asche Basis [®] cream	22197A	30775	40	not required
Chiesi	Asche Basis [®] lot...	21042A	30828	40	not required
Actavis	Abitima [®] clinic ...	46820	30845	40	not required
Beiersdorf	Eucerinum W/O ba...	31230108WA	30993	31	not required
Taoasis	Base cream Taoasis	121205-2276	31041	80	not required
Intendis	Neribas [®] cream	21076A	31050	40	not required
Actavis	Abitima [®] clinic ...	110712	31058	40	not required
Actavis	Abitima [®] clinic ...	051112	31061	40	not required
Spirig Pharma	Excipial [®] U10 Li...	N025	31066	40	not required
Spirig Pharma	Excipial [®] U Lipo...	N031	31067	40	not required
Chiesi	Asche Basis [®] lot...	24049A	31068	40	not required
Spirig Pharma	Lipoderm [®] lotion	M065	31072	40	not required
Chiesi	Asche Basis [®] cream	24220A	31085	40	not required
Spirig Pharma	Excipial [®] hydro ...	N013	31089	40	not required
Intendis	Neribas [®] cream	21076A	31095	40	not required
Bene	Thrombocid [®] oint...	410B113	31098	40	not required
Taoasis	Base cream Taoasis	130124-3021	31115	40	not required
Sanofi aventis	Dermatop [®] base c...	3F031A	31234	40	not required
Koko	DMS [®] base cream ...	L010K13	31236	70	not required
Spirig Pharma	Excipial [®] U Lipo...	N085	31280	40	not required
Spirig Pharma	Excipial [®] U10 Li...	N047	31281	40	not required
Koko	DMS [®] base cream ...	L026G13	31307	40	not required
Koko	DMS [®] base cream ...	L032A14	31308	40	not required
Koko	DMS [®] base cream ...	L014L13	31309	40	not required
Dr. Wolff	Linola [®] body milk	433240	31511	40	not required
Spirig Pharma	Lipoderm [®] lotion	R023	31535	40	not required
Beiersdorf	Eucerinum O/W ba...	41830108WA	31539	40	not required
Taoasis	Base cream Taoasis	141222-4350	31578	40	not required
Hans Karrer G...	Hans Karrer Lipo...	15003	31730	40	not required
Koko	DMS [®] base cream ...	L034H15	31869	40	not required
Koko	DMS [®] base cream ...	L041K15	31870	40	not required
Hans Karrer G...	Hans Karrer Lipo...	15008	31873	40	not required
Dr. Wolff	Linola [®] body milk	511750	31874	40	not required
Sanofi aventis	Dermatop [®] base c...	5F056A	32177	40	not required
Koko	DMS [®] base cream ...	L003L15	32178	40	not required
Beiersdorf	Eucerinum W/O ba...	53430208WA	32767	40	not required
Koko	DMS [®] base cream ...	L003H16	32916	40	not required
Actavis	Abitima [®] clinic ...	070915	32997	40	not required
Hans Karrer G...	Hans Karrer Lipo...	15003	33194	40	not required
Hans Karrer G...	Hans Karrer Lipo...	15005	33195	40	not required
Spirig Pharma	Excipial [®] hydro ...	T052	33208	40	not required
Beiersdorf	Eucerinum O/W ba...	64030208WA	33213	40	not required
Galderma Labo...	Excipial [®] U Lipo...	T122	33270	40	not required
Allergika	Allergika [®] Base ...	11/082022	33668	40	not required
Allergika	Allergika [®] Base ...	10/082022	33669	40	not required
Chiesi	Asche Basis [®] lot...	64082A	33737	40	not required
Actavis	Abitima [®] clinic ...	161464	33793	30	not required
Koko	DMS [®] base cream ...	L009K17	33968	40	not required
Actavis	Abitima [®] clinic ...	331017	33969	40	not required
Dr. Wolff	Linola [®] body milk	707930	34043	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 2371 spectra of 57 reference samples from the substance/substance group *Hydrophilic ointments*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 53 different batches.
- 16 378 spectra from a total of 387 batches from further 130 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 2089 spectra from 58 reference samples of the substance/substance group *Hydrophilic ointments*.
- These include spectra of independent samples from 32 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Beiersdorf	Eucerinum O/W basis	zu erfassen	30409	40
Beiersdorf	Eucerinum W/O basis	21430108WA	30494	40
Beiersdorf	Eucerinum O/W basis	0930108WA	30565	40
Actavis	Abitima [®] clinic body cream	072013	30736	39
Spirig Pharma	Excipial [®] hydro cream	I013	30836	40
Taoasis	Base cream Taoasis	121205-2276	31041 [†]	20
Spirig Pharma	Excipial [®] hydro cream	N013	31277	40
Dr. Wolff	Linola [®] body milk	433240	31511 [†]	20
Spirig Pharma	Lipoderm [®] lotion	R023	31535 [†]	20
Beiersdorf	Eucerinum O/W basis	41830108WA	31539 [†]	20
Taoasis	Base cream Taoasis	141222-4350	31578 [†]	20
Hans Karrer GmbH	Hans Karrer Lipolotion Micro...	15003	31730 [†]	20
Koko	DMS [®] base cream classic	L034H15	31869 [†]	20
Koko	DMS [®] base cream high classic	L041K15	31870 [†]	20
Hans Karrer GmbH	Hans Karrer Lipolotion Micro...	15008	31873 [†]	20
Dr. Wolff	Linola [®] body milk	511750	31874 [†]	20
Actavis	Abitima [®] clinic face cream	158584	32179	40
Chiesi	Asche Basis [®] lotion	53075A	32715	40

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra
Sanofi aventis	Dermatop [®] base cream	5F057A	32717	40
Actavis	Abitima [®] clinic face cream	158584	32723	40
Bene	Thrombocid [®] ointment	408C141	32742	40
Hans Karrer GmbH	Hans Karrer Lipolotion Micro...	16005	32753	40
Dr. Wolff	Linola [®] body milk	606430	32765	40
Koko	DMS [®] base cream classic	L003E16	32788	40
Koko	DMS [®] base cream high classic	L003H16	32789	40
Koko	DMS [®] base cream classic	L003E16	32826	40
Koko	DMS [®] base cream high classic...	L018D16	32827	40
Bene	Thrombocid [®] ointment	409A151	32915	40
Chiesi	Asche Basis [®] lotion	53075A	32926	40
Chiesi	Asche Basis [®] cream	63351A	32928	40
Taoasis	Base cream Taoasis	161124-6322	32944	40
Taoasis	Base cream Taoasis	161124-6322	32945	40
Actavis	Abitima [®] clinic face cream	158584	32998	40
Actavis	Abitima [®] clinic body cream	040116	32999	40
Hans Karrer GmbH	Hans Karrer Lipocream MicroS...	17001	33196	40
Hans Karrer GmbH	Hans Karrer Lipocream MicroS...	16002	33197	40
Spirig Pharma	Excipial [®] U Lipolotio	T122	33204	40
Spirig Pharma	Excipial [®] U10 Lipolotio	T072	33205	40
Beiersdorf	Eucerinum O/W basis	64030208WA	33212	40
Spirig Pharma	Lipoderm [®] lotion	V012	33248	40
Dr. Wolff	Linola [®] body milk	609202	33258	40
Galderma Laboratorium G...	Excipial [®] U Lipolotio	T122	33269	40
Jenapharm	Neribas [®] cream	YY01FX3	33289	40
Galderma Laboratorium G...	Excipial [®] U Lipolotio	V011	33298	40
Koko	DMS [®] base cream high classic	L037K16	33433	40
Koko	DMS [®] base cream high classic	L037K16	33434	40
Chiesi	Asche Basis [®] cream	63354A	33435	40
Beiersdorf	Eucerinum W/O basis	72630108WA	33629	40
Allergika	Allergika [®] Base cream	09/072022	33670	40
Actavis	Abitima [®] clinic body cream	040116	33791	40
Actavis	Abitima [®] clinic face cream	161464	33793 [†]	10
Beiersdorf	Eucerinum O/W basis	64030208WA	33850	40
Beiersdorf	Eucerinum W/O basis	72630108WA	33922	40
Jenapharm	Neribas [®] cream	YY01X64	33951	40
Sanofi aventis	Dermatop [®] base cream	7F069A	33966	40
Spirig Pharma	Excipial [®] U10 Lipolotio	50393	33980	40
Spirig Pharma	Excipial [®] hydro cream	50235	33982	40
Spirig Pharma	Lipoderm [®] lotion	V042	33985	40

- 20 742 spectra from a total of 502 batches from further 182 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 738 spectra from 119 *Apo-Ident* customers from 362 batches from the substance/substance group *Hydrophilic ointments*.

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

- These include spectra of independent samples from 349 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Actavis	Abitima [®] clinic body cream	10915	2
Actavis/Noweda	Abitima [®] clinic body cream	21015	1
Actavis Deutschland GmbH	Abitima [®] clinic body cream	31015	1
Actavis	Abitima [®] clinic body cream	31015	2
Actavis	Abitima [®] clinic body cream	31114	1
Actavis/Gehe	Abitima [®] clinic body cream	81013	1
Actavis Deutschland GmbH	Abitima [®] clinic body cream	121115	1
Actavis/Phönix	Abitima [®] clinic face cream	159274	2
Actavis	Abitima [®] clinic face cream	160670	1
Actavis	Abitima [®] clinic body cream	260515	2
Actavis/Noweda	Abitima [®] clinic face cream	161423	1
actavis/Phönix	Abitima [®] clinic face cream	161468	2
Actavis/Noweda	Abitima [®] clinic body cream	260515	2
Persano/Gehe	Abitima [®] clinic face cream	57792	1
actavis	Abitima [®] clinic body cream	60513	2
actavis/Gehe	Abitima [®] clinic body cream	60513	1
actavis	Abitima [®] clinic body cream	70513	1
Caelo	Abitima [®] clinic body cream	110712	2
actavis	Abitima [®] clinic body cream	140113	1
Actavis/Noweda	Abitima [®] clinic body cream	140113	1
Fagron	Abitima [®] clinic face cream	150204	1
Actavis/Noweda	Abitima [®] clinic face cream	150204	1
Fagron	Abitima [®] clinic face cream	150203	2
Actavis / Anzag	Abitima [®] clinic face cream	150237	1
Astellas Pharma	Abitima [®] clinic body cream	190213	1
Actavis Deutschland GmbH	Abitima [®] clinic body cream	350313	2
Actavis / Noweda	Abitima [®] clinic face cream	142371	1
Noweda	Abitima [®] clinic face cream	146735	1
Actavis / Anzag	Abitima [®] clinic face cream	148024	1
Caelo	Abitima [®] clinic face cream	146735	1
Actavis/GEHE	Abitima [®] clinic face cream	148051	1
Actavis/Noweda	Abitima [®] clinic face cream	148417	1
Phoenix/actavis	Abitima [®] clinic face cream	148460	1
actavis	Abitima [®] clinic face cream	148495	1
Actavis/Gehe	Abitima [®] clinic face cream	148495	1
actavis	Abitima [®] clinic face cream	148496	2
Actavis/ Noweda	Abitima [®] clinic face cream	148496	1
Actavis / Anzag	Abitima [®] clinic face cream	148503	1
Caelo	Abitima [®] clinic face cream	149666	2
actavis	Abitima [®] clinic face cream	150202	1
actavis	Abitima [®] clinic body cream	20912	2
Sanacorp	Abitima [®] clinic body cream	12015/110/12	1
Actavis	Abitima [®] clinic body cream	20912	1
Actavis/Gehe	Abitima [®] clinic body cream	51112	1
Sanacorp	Abitima [®] clinic body cream	51112	1
Abitima	Abitima [®] clinic body cream	52015/051112	1
Actavis/ Gehe	Abitima [®] clinic body cream	60513	1
Sanacorp	Abitima [®] clinic body cream	70512	1
Actavis / Gehe	Abitima [®] clinic body cream	110712	1
actavis/Gehe	Abitima [®] clinic body cream	112015	1
actavis/ebert	Abitima [®] clinic body cream	110712	1
Sanacorp	Abitima [®] clinic body cream	140113	1
Roche/AHD	Abitima [®] clinic body cream	142620	1
Sanacorp	Abitima [®] clinic body cream	210612	1
Actavis/Noweda	Abitima [®] clinic body cream	370313	1
Actavis/Sanacorp	Abitima [®] clinic body cream	110712	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Chiesi	Asche Basis [®] cream	30102014A	1
Chiesi/Noweda	Asche Basis [®] cream	32312A	1
Caelo	Asche Basis [®] lotion	32054A	1
Chiesi	Asche Basis [®] lotion	32054A	1
Chiesi	Asche Basis [®] cream	32233A	5
Caelo	Asche Basis [®] lotion	33055A	3
chiesi/gehe	Asche Basis [®] lotion	33056A	2
Chiesi	Asche Basis [®] cream	33239A	2
Chiesi / Noweda	Asche Basis [®] cream	33246A	1
Chiesi GmbH / Anzag	Asche Basis [®] cream	33244A	1
chiesi/gehe	Asche Basis [®] cream	33246A	2
Chiesi	Asche Basis [®] lotion	333056A	2
Chiesi/Noweda	Asche Basis [®] cream	34247A	1
chiesi/gehe	Asche Basis [®] lotion	34058A	1
Chiesi/AHD	Asche Basis [®] cream	34247A	1
Chiesi	Asche Basis [®] cream	34249A	1
Chiesi GmbH / Anzag	Asche Basis [®] cream	34249A	1
Chiesi GmbH / Sanacorp	Asche Basis [®] cream	34247A	1
Chiesi/V.d.L.	Asche Basis [®] cream	34250A	1
Chiesi/Gehe	Asche Basis [®] cream	34248A	1
Chiesi	Asche Basis [®] cream	34250A	1
Chiesi	Asche Basis [®] cream	34248A	1
Chiesi	Asche Basis [®] cream	34251A	3
chiesi / Noweda	Asche Basis [®] cream	34250A	1
Chiesi / Gehe	Asche Basis [®] cream	34252A	1
Chiese	Asche Basis [®] lotion	41060A	1
Chiesi/Noweda	Asche Basis [®] cream	41255A	2
Chiesi	Asche Basis [®] cream	41258A	2
Chiesi	Asche Basis [®] lotion	41059A	1
Chiesi/Noweda	Asche Basis [®] cream	41258A	2
Chiese	Asche Basis [®] cream	41259A	1
Caelo	Asche Basis [®] cream	41261A	1
Chiesi/ Now	Asche Basis [®] cream	41259A	1
Chiesi GmbH / Anzag	Asche Basis [®] cream	41261A	1
Chiesi/Gehe	Asche Basis [®] cream	41260A	1
Chiesi GmbH / Anzag	Asche Basis [®] cream	41260A	1
Chiesi	Asche Basis [®] cream	41260A	2
Chiesi/Gehe	Asche Basis [®] cream	41262A	1
Chiesi	Asche Basis [®] cream	41261A	1
Chiesi	Asche Basis [®] cream	41264A	1
Chiesi GmbH / Anzag	Asche Basis [®] cream	42268A	2
Chiesi / Gehe	Asche Basis [®] cream	41265A	1
Caelo	Asche Basis [®] cream	41265B	1
Chiesi/	Asche Basis [®] cream	41266A	1
Chiesi/Gehe	Asche Basis [®] cream	42271A	1
Anzag/ Chiesi	Asche Basis [®] cream	42269A	1
Chiesi	Asche Basis [®] cream	42270A	1
Chiesi	Asche Basis [®] cream	42271A	2
Chiesi	Asche Basis [®] cream	42275A	1
Chiesi	Asche Basis [®] cream	42272A	1
Anzag/ Chiesi GmbH	Asche Basis [®] cream	42273A	1
Anzag/ Chiesi GmbH	Asche Basis [®] cream	42272A	1
Chiesi/ Phönix	Asche Basis [®] lotion	43066A	1
Chiesi/ Now	Asche Basis [®] cream	42273A	1
Chiesi/Noweda	Asche Basis [®] cream	42275A	2
gehe/Chiesi	Asche Basis [®] lotion	43066A	1
Chiesi	Asche Basis [®] lotion	43066A	1
Chiesi	Asche Basis [®] cream	43278A	1
Chiesi/Phoenix	Asche Basis [®] cream	43278A	1
Chiesi/ Now	Asche Basis [®] cream	43283A	2

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Chiesi	Asche Basis [®] cream	43279A	1
Chiesi	Asche Basis [®] cream	43280A	1
Chiesi/Noweda	Asche Basis [®] cream	43280A	1
Chiesi/ Now	Asche Basis [®] cream	43282A	1
Chiesi / Gehe	Asche Basis [®] cream	43281A	1
Chiesi/Noweda	Asche Basis [®] cream	43284A	1
Chiesi	Asche Basis [®] cream	43287A	1
Chiesi/Noweda	Asche Basis [®] cream	44288A	1
Chiesi/Alliance Healthc...	Asche Basis [®] cream	44290A	4
Chiesi GmbH / Phönix	Asche Basis [®] lotion	51067A	1
Chiesi/ Noweda	Asche Basis [®] cream	51292A	1
Gehe/Chiesi	Asche Basis [®] cream	51296A	1
Chiesi / Phoenix	Asche Basis [®] cream	51300A	1
Chiesi/ Noweda	Asche Basis [®] cream	51294a	1
Chiesi	Asche Basis [®] cream	51294A	1
Chiesi/Alliance Healthc...	Asche Basis [®] cream	51300A	4
Chiesi	Asche Basis [®] cream	51296A	1
Chiesi/Gehe	Asche Basis [®] cream	51301A	1
Chiesi	Asche Basis [®] cream	51305A	2
Phoenix	Asche Basis [®] lotion	52069A	1
Caelo	Asche Basis [®] cream	52306A	3
Gehe20.10.15	Asche Basis [®] lotion	52070A	1
noweda/chiesi	Asche Basis [®] cream	52306A	1
Chiesi/Phoenix	Asche Basis [®] lotion	52070A	1
Gehe-Chiesi	Asche Basis [®] cream	52308A	1
Chiesi	Asche Basis [®] cream	52308A	2
Chiesi	Asche Basis [®] cream	52307A	1
Chiesi/Gehe	Asche Basis [®] cream	52307A	1
Chiesi/ Noweda	Asche Basis [®] cream	52307A	1
Chiesi	Asche Basis [®] cream	52309A	1
Chiesi/Noweda	Asche Basis [®] cream	52316A	1
Chiesi/Noweda	Asche Basis [®] cream	52318a	1
Gehe/Chesie	Asche Basis [®] cream	52318A	2
Chiesi/Noweda	Asche Basis [®] cream	52318A	1
Chiesi GmbH	Asche Basis [®] cream	52319A	1
Caelo	Asche Basis [®] cream	52322A	3
Chiesi/Noweda	Asche Basis [®] cream	52321A	1
Chiesi	Asche Basis [®] cream	52323A	4
Chiesi	Asche Basis [®] cream	52322A	1
Chiesi/Phönix	Asche Basis [®] cream	52323A	2
Caelo	Asche Basis [®] cream	52323a	1
Chiesi GmbH	Asche Basis [®] lotion	53071A	3
Chiesi/Phönix	Asche Basis [®] lotion	53071A	1
Chiesi/Phoenix	Asche Basis [®] lotion	53071A	1
Chiesi GmbH	Asche Basis [®] lotion	53072A	2
Caelo	Asche Basis [®] lotion	53072A	1
Caelo	Asche Basis [®] lotion	53073A	1
Chiesi/Fiebig	Asche Basis [®] lotion	53073A	2
Chiesi	Asche Basis [®] lotion	53073A	1
Chiesi GmbH	Asche Basis [®] lotion	53074A	1
Chiesi	Asche Basis [®] lotion	53075A	1
Chiesi/Anzag	Asche Basis [®] cream	53327A	1
Chiesi/Phönix	Asche Basis [®] cream	53329A	1
Chiesi/Alliance Healthc...	Asche Basis [®] cream	53329A	1
Chiesi	Asche Basis [®] cream	53329A	1
Chiesi	Asche Basis [®] cream	53330A	1
Caelo	Asche Basis [®] cream	53330A	1
Chiesi	Asche Basis [®] cream	53331A	2
Phoenix	Asche Basis [®] cream	53331A	1
Caelo	Asche Basis [®] cream	53331A	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Phoenix	Asche Basis [®] cream	54332A	1
Chiesi	Asche Basis [®] cream	54332A	1
Chiesi GmbH	Asche Basis [®] lotion	61076A	2
Chiesi/Noweda	Asche Basis [®] lotion	61077A	1
Chiesi GmbH	Asche Basis [®] cream	61335A	1
Caelo	Asche Basis [®] cream	61338A	1
Chiesi	Asche Basis [®] lotion	61077A	1
Caelo	Asche Basis [®] cream	61341A	2
Chiesi	Asche Basis [®] cream	62343A	2
PHÖNIX	Asche Basis [®] cream	62343A	1
Chiesi GmbH	Asche Basis [®] lotion	63078A	2
Chiesi	Asche Basis [®] lotion	63079A	2
Chisi	Asche Basis [®] cream	63347A	3
Chiesi GmbH	Asche Basis [®] lotion	63980A	1
Noweda	Asche Basis [®] lotion	63980A	1
Chiesi	Asche Basis [®] cream	64357A	3
Chiesi	Asche Basis [®] lotion	64082A	1
Chiesi/Fiebig	Asche Basis [®] lotion	63980A	1
Chiesi//Phönix	Asche Basis [®] cream	63354A	1
Phoenix/Chiesi	Asche Basis [®] cream	64358A	1
Gehe	Asche Basis [®] cream	64358A	2
Chiesi	Asche Basis [®] cream	64360A	1
Chiesi GmbH / Anzag	Asche Basis [®] cream	22196A	1
Chiesi	Asche Basis [®] cream	14178A	1
Chisi	Asche Basis [®] cream	23203A	1
Anzag	Asche Basis [®] cream	23204A	1
Chiesi	Asche Basis [®] cream	23206A	1
Chiesi	Asche Basis [®] cream	24214A	2
Chiesi GmbH	Asche Basis [®] lotion	73087A	1
Chiesi	Asche Basis [®] cream	23204A	1
chiesi / Noweda	Asche Basis [®] cream	23213A	1
Chiesi/Phoenix	Asche Basis [®] cream	24215A	1
Chiesi GmbH	Asche Basis [®] cream	24214A	2
Chiesi	Asche Basis [®] cream	24217A	1
Chiesi/Gehe	Asche Basis [®] cream	24218A	1
All.Heal./Chiesi	Asche Basis [®] cream	24218A	1
Chiesi	Asche Basis [®] cream	24219A	1
Chiesi	Asche Basis [®] cream	24220A	2
Chiesi/Gehe	Asche Basis [®] cream	24221A	4
Chiesi/Anzag	Asche Basis [®] cream	24221A	1
Ichthyol/Sanacorp	Asche Basis [®] cream	24221A	1
Chiese	Asche Basis [®] cream	24221A	1
Chiesi/AHD	Asche Basis [®] cream	31222A	1
chiesi / Noweda	Asche Basis [®] cream	31222A	1
Chiesi GmbH / Anzag	Asche Basis [®] cream	31222A	1
Chiesi/AHD	Asche Basis [®] cream	31223A	1
Chiesi/Ebert	Asche Basis [®] cream	31224A	1
Chiesi/Noweda	Asche Basis [®] cream	31224A	1
Chiesi/AHD	Asche Basis [®] cream	31224A	1
Chiesi/Gehe	Asche Basis [®] cream	31224A	2
Chiesi/Phoenix	Asche Basis [®] cream	31225A	1
chiesi / Noweda	Asche Basis [®] cream	31225A	1
Caelo	Asche Basis [®] cream	31226A	1
chiesi /Jenne	Asche Basis [®] cream	31227a	1
Chiesi/Phoenix	Asche Basis [®] cream	31227A	2
Caelo	Asche Basis [®] cream	31227A	1
Chiesi/Sanacorp	Asche Basis [®] cream	31227A	1
Chiesi	Asche Basis [®] cream	31227A	2
Chiesi/AHD	Asche Basis [®] cream	31228A	2
Chiesi/Noweda	Asche Basis [®] cream	31229A	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
chiesi/gehe	Asche Basis [®] cream	31228A	1
Chiesi GmbH / Anzag	Asche Basis [®] cream	31229A	1
Chiesi/Phoenix	Asche Basis [®] cream	31229A	1
Chiesi GmbH / Anzag	Asche Basis [®] cream	31230A	1
Chiesi GmbH / Anzag	Asche Basis [®] cream	31231A	2
Chiesi/Noweda	Asche Basis [®] cream	31232A	2
Chiesi	Asche Basis [®] cream	31232A	1
Chiesi/Gehe	Asche Basis [®] cream	32233A	1
Gehe/Chiesi	Asche Basis [®] cream	32233A	1
Chiesi/Phoenix	Asche Basis [®] cream	32233A	1
Phönix	Asche Basis [®] cream	32234A	1
Chiesi	Asche Basis [®] cream	32235A	1
Chiesi/AHD	Asche Basis [®] cream	32234A	1
chiesi/gehe	Asche Basis [®] cream	32236A	1
Chiesi/AHD	Asche Basis [®] cream	32235A	1
Chiesi	Asche Basis [®] cream	32236A	1
Gehe	Asche Basis [®] cream	32236A	1
Euro OTC	Asche Basis [®] cream	3229A	2
Chiesi/ Noweda	Asche Basis [®] cream	33237A	1
Chiesi	Asche Basis [®] cream	33237A	1
Alliance Healthcare	Asche Basis [®] cream	33237A	1
Chiesi/Noweda	Asche Basis [®] cream	33237A	1
Chiesi/ phoenix	Asche Basis [®] cream	33239A	1
Chiesi/AHD	Asche Basis [®] cream	33238A	1
Chiesi / Noweda	Asche Basis [®] cream	33238A	1
Chiesi/ Now	Asche Basis [®] cream	33240A	1
Chiesi/ Noweda	Asche Basis [®] cream	33241A	1
Alliance Healthcare	Asche Basis [®] cream	33240A	1
Chiesi	Asche Basis [®] cream	33240A	1
Chiesi/AHD	Asche Basis [®] cream	33242A	1
Chiesi	Asche Basis [®] cream	33243A	1
Chiesi / Gehe	Asche Basis [®] cream	33244A	1
Phoenix/Chiesi	Asche Basis [®] cream	33246A	1
Gehe	Asche Basis [®] cream	33244A	1
Chiesi/Gehe	Asche Basis [®] cream	34253A	1
Lamotte / Gehe	Asche Basis [®] cream	34247A	1
Chiesi/Noweda	Asche Basis [®] cream	53329A	1
Chiesi/Gehe	Asche Basis [®] lotion	23047A	1
Chiesi/Noweda	Asche Basis [®] lotion	24049A	1
Chiesi/AHD	Asche Basis [®] lotion	24049A	1
Chiesi/Anzag	Asche Basis [®] lotion	32054A	1
Taoasis	Base cream Taoasis	150219-4350	1
Taoasis	Base cream Taoasis	150326-5079	1
Taoasis	Base cream Taoasis	160201-6027	1
Taoasis	Base cream Taoasis	161124-6322	1
Taoasis	Base cream Taoasis	170110-6322	1
Caelo	Base cream Taoasis	171127-7290	1
Sanofi/Gehe	Dermatop [®] base cream	4F050A	1
Sanofi/Noweda	Dermatop [®] base cream	5F056A	1
Sanofi/Anzag	Dermatop [®] base cream	5F056A	1
Sanofi/Phönix	Dermatop [®] base cream	5F051A	1
Sanofi/Noweda	Dermatop [®] base cream	5F055A	2
Sanofi/Noweda	Dermatop [®] base cream	4F045A	1
Sanofi/Anzag	Dermatop [®] base cream	5F051A	1
Sanofi Aventis	Dermatop [®] base cream	4F046A	1
Sanofi	Dermatop [®] base cream	4F046A	1
KOKO GmbH	DMS [®] base cream classic	L003E16	1
Caelo	DMS [®] base cream high classic	L010J15	1
KOKO GmbH	DMS [®] base cream high classic	L030L16	3
KOKO GmbH & Co. KG	DMS [®] base cream high classic...	L018D16	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
KOKO GmbH & Co. KG	DMS [®] base cream high classic	L054B16	1
Koko GmbH & Co.Kg	DMS [®] base cream high classic	L060A15	1
KOKO GmbH	DMS [®] base cream classic	L005A14	1
KOKO GMBH	DMS [®] base cream high classic	L008I14	1
Koko GmbH	DMS [®] base cream high classic	L031A14	1
KOKO GMBH	DMS [®] base cream high classic	L031A14	1
KOKO	DMS [®] base cream high classic	L031A14	1
Beiersdorf	Eucerinum W/O basis	24072015A	1
Beiersdorf AG	Eucerinum W/O basis	32930208WA	3
Beiersdorf	Eucerinum W/O basis	34930308WA	5
Beiersdorf/Phoenix	Eucerinum W/O basis	34930308WA	1
Beiersdorf AG	Eucerinum W/O basis	34930308WA	1
Beiersdorf/Noweda	Eucerinum W/O basis	34930308WA	1
Beiersdorf/ Vedono	Eucerinum W/O basis	34930308WA	1
Beiersdorf/ Sanacorp	Eucerinum W/O basis	418301087WA	1
Beiersdorf/Noweda	Eucerinum W/O basis	41830108WA	2
Beiersdorf	Eucerinum W/O basis	41830108WA	2
Beiersdorf AG	Eucerinum W/O basis	41830108WA	1
Beiersdorf/Noweda	Eucerinum W/O basis	41830108wa	1
Caelo	Eucerinum W/O basis	41830108WA	3
BDF	Eucerinum W/O basis	41830108WA	1
Ph.Eur.8.0	Eucerinum W/O basis	41830108WA	1
Baiersdorf	Eucerinum W/O basis	41839108WA	1
Baiersdorf	Eucerinum W/O basis	43430208WA	1
Beiersdorf/Gehe	Eucerinum W/O basis	43430208WA	1
BDF / Noweda	Eucerinum W/O basis	43430208WA	1
Caelo	Eucerinum W/O basis	51630108	1
Gehe/Beiersdorf	Eucerinum W/O basis	51630108WA	1
Beiersdorf	Eucerinum W/O basis	51630108WA	6
Beiersdorf/Noweda	Eucerinum W/O basis	51630108WA	3
Beiersdorf/Noweda	Eucerinum W/O basis	51630108wa	1
Caelo	Eucerinum W/O basis	51630108wa	1
Noweda/BDF	Eucerinum W/O basis	51630108WA	1
Rondell Apotheke	Eucerinum W/O basis	51630108WA	1
Beiersdorf AG	Eucerinum W/O basis	51630108WA	2
Caelo	Eucerinum W/O basis	51630108WA	1
Beiersdorf/AHD	Eucerinum O/W basis	51630108WA	1
Beiersdorf AG/Noweda	Eucerinum O/W basis	53430208WA	1
Caelo	Eucerinum W/O basis	53430208WA	4
Beiersdorf	Eucerinum W/O basis	53430208WA	1
Beiersdorf/Kehr	Eucerinum O/W basis	53430208WA	3
Beiersdorf/Noweda	Eucerinum W/O basis	53430208WA	4
Beiersdorf/Phönix	Eucerinum W/O basis	53430208WA	1
Beiersdor	Eucerinum W/O basis	541830108WA	1
Beiersdorf/Kehr	Eucerinum O/W basis	61130108WA	1
Beiersdorf/ Noweda	Eucerinum W/O basis	64030108WA	1
Caelo	Eucerinum W/O basis	64030108WA	2
Beiersdorf	Eucerinum W/O basis	64030108WA	1
Allianz Healthcare	Eucerinum W/O basis	64030108wa	1
Beiersdorf/Kehr	Eucerinum W/O basis	64030108WA	1
Beiersdorf/AHD	Eucerinum O/W basis	64030208WA	1
Beiersdorf/VDL	Eucerinum O/W basis	64030208WA	1
Noweda	Eucerinum W/O basis	72630108WA	4
Beiersdorf	Eucerinum W/O basis	72630108WA	1
Caelo	Eucerinum O/W basis	13730308WA	1
Beiersdorf AG	Eucerinum O/W basis	2483030308WA	1
Beiersdorf	Eucerinum O/W basis	24830308WA	1
Hersteller: Beiersdorf/...	Eucerinum O/W basis	24830308WA	1
Beiersdorf AG	Eucerinum O/W basis	24830308WA	1
Beiersdorf	Eucerinum O/W basis	319030108WA	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Caelo	Eucerinum O/W basis	31930108WA	1
Beiersdorf/Gehe	Eucerinum O/W basis	31930108WA	1
Beiersdorf	Eucerinum O/W basis	51630108WA	1
Beiersdorf	Eucerinum O/W basis	53430208WA	1
BDF / Sanacorp	Eucerinum W/O basis	21430108WA	1
Noweda	Eucerinum W/O basis	4730308WA	1
Beiersdorf	Eucerinum W/O basis	21430108WA	2
Beiersdorf	Eucerinum W/O basis	24830208WA	1
Baierdorf	Eucerinum W/O basis	24830208WA	1
Beiersdorf/Sanacorp	Eucerinum W/O basis	31230108WA	1
Beiersdorf/Anzag	Eucerinum W/O basis	31230108WA	1
beiersdorf	Eucerinum W/O basis	24830208WA	1
beiersdorf	Eucerinum W/O basis	31230108WA	1
Beiersdorf/Noweda	Eucerinum W/O basis	31230108WA	1
BDF	Eucerinum W/O basis	31230108WA	1
BeiersdorfPhoenix	Eucerinum W/O basis	31230108WA	1
Beiersdorf	Eucerinum W/O basis	31230108wa	1
Beiersdorf	Eucerinum W/O basis	31230108WA	1
Beiersdorf AG/Noweda	Eucerinum W/O basis	31230108WA	1
Beiersdorf	Eucerinum W/O basis	32930208WA	4
Gehe	Eucerinum W/O basis	32930208WA	1
Beiersdorf	Eucerinum W/O basis	516301328wa	1
eucerinum	Eucerinum W/O basis	41830108WA	1
Spirig Pharma AG	Excipial [®] hydro cream	813993/12	1
Galderma/Spirig Pharma AG	Excipial [®] U Lipolotio	LOTR187	1
Gehe-Spring	Excipial [®] U Lipolotio	LOTR167	1
Galderma, Sanacorp	Excipial [®] U Lipolotio	LOTS042	1
Caelo	Excipial [®] U Lipolotio	LOTS081	1
galderma/G	Excipial [®] U Lipolotio	LOTS202	1
Gehe/Galderma	Excipial [®] U Lipolotio	LotV011	1
Spirig/Fiebig	Excipial [®] hydro cream	N013	1
Spirig/Gehe	Excipial [®] hydro cream	R014	1
Spirig	Excipial [®] hydro cream	R014	2
Caelo	Excipial [®] hydro cream	R045	1
Spirig	Excipial [®] U Lipolotio	R047	1
Galderma (Spirig Pharma...	Excipial [®] U Lipolotio	R061	1
Spirig/Galderma	Excipial [®] U Lipolotio	R146	1
Spirig/Galderma	Excipial [®] U Lipolotio	R156	2
Galderma	Excipial [®] U Lipolotio	R167	2
Galderma	Excipial [®] U Lipolotio	R168	1
Galderma / Kehr	Excipial [®] U Lipolotio	R167	1
Galderma / Kehr	Excipial [®] U Lipolotio	R171	1
Galderma	Excipial [®] U Lipolotio	R171	1
Caelo	Excipial [®] U Lipolotio	R186	1
Galderma/Gehe	Excipial [®] U Lipolotio	R189	1
Noweda	Excipial [®] U Lipolotio	R261	1
Spirig/Gehe	Excipial [®] U Lipolotio	R262	1
Sprig (Galderma)/ G	Excipial [®] U Lipolotio	S123	1
Galderma/Galderma	Excipial [®] U Lipolotio	s017	1
Galderma/Galderma	Excipial [®] U Lipolotio	S017	1
Galderma/Galderma	Excipial [®] hydro cream	S024	2
Galderma/ Sana	Excipial [®] hydro cream	S024	1
Spirig	Excipial [®] hydro cream	S025	6
Galderma/ Sana	Excipial [®] hydro cream	S025	1
spirig	Excipial [®] hydro cream	s025	1
Jenne	Excipial [®] hydro cream	S041	1
Galderma/Phönix	Excipial [®] hydro cream	S045	1
Spirig Pharma/ G	Excipial [®] hydro cream	S053	1
Galderma	Excipial [®] hydro cream	S054	1
Galderma/Galderma	Excipial [®] hydro cream	S054	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Galderma/Noweda	Excipial [®] hydro cream	S054	1
Galderma	Excipial [®] U Lipolotio	S064	1
Galderma/Kehr	Excipial [®] U Lipolotio	S071	1
Galderma	Excipial [®] U Lipolotio	S091	1
Phoenix	Excipial [®] U Lipolotio	S091	1
Spirig Pharma	Excipial [®] U Lipolotio	S122	1
Gehe	Excipial [®] U Lipolotio	S132	1
Phoenix	Excipial [®] U Lipolotio	S138	2
Galderma/Galderma	Excipial [®] U Lipolotio	S142	1
Gehe	Excipial [®] U Lipolotio	S143	1
Galderma	Excipial [®] U Lipolotio	S147	1
Spirig/Fiebig	Excipial [®] U Lipolotio	S143	1
Phoenix	Excipial [®] U Lipolotio	S148	1
Galderma/Kehr	Excipial [®] U Lipolotio	S203	1
Galderma/Anzag	Excipial [®] U Lipolotio	S206	1
Caelo	Excipial [®] U Lipolotio	S206	1
Galderma/Alliance Healt...	Excipial [®] hydro cream	S025	1
Galderma/Galderma	Excipial [®] hydro cream	so25	1
Spirig (Galderma)	Excipial [®] hydro cream	T011	1
Spirig (Galderma)	Excipial [®] hydro cream	T012	2
Galderma / Kehr	Excipial [®] U Lipolotio	T029	2
Kehr/Galderma	Excipial [®] U Lipolotio	T029	1
PHÖNIX	Excipial [®] U Lipolotio	T063	1
Spirig (Galderma)	Excipial [®] hydro cream	V011	2
Galderma/Noweda	Excipial [®] U Lipolotio	V011	1
Galderma/Noweda	Excipial [®] U Lipolotio	V041	1
Spirig Pharma	Excipial [®] U Lipolotio	V053	1
Spirig Pharma	Excipial [®] U Lipolotio	V082	1
Spirig Pharma/Gehe	Excipial [®] hydro cream		1
Spirig/Noweda	Excipial [®] U Lipolotio	LOTN113	1
Spirig Pharma AG/ Anzag	Excipial [®] hydro cream	LOTN034	1
Spirig / Anzag	Excipial [®] hydro cream	LOTN043	1
Spirig / Gehe	Excipial [®] hydro cream	LOTN044	1
Spirig/AHD	Excipial [®] U Lipolotio	lotno86	1
Spirig Pharma AG/ Anzag	Excipial [®] hydro cream	LOTR013	1
Fischar	Excipial [®] U Lipolotio	LOTN085	1
Phoenix/Spirig	Excipial [®] hydro cream	N044	1
Spirig / Gehe	Excipial [®] hydro cream	N011	1
spirig	Excipial [®] hydro cream	N034	2
Spirig / Noweda	Excipial [®] hydro cream	N035	1
Spirig/Noweda	Excipial [®] U10 Lipolotio	N043	1
spirig	Excipial [®] hydro cream	N043	2
Spirig	Excipial [®] hydro cream	N043	1
Spirig/Noweda	Excipial [®] hydro cream	N043	1
spirig	Excipial [®] hydro cream	N045	2
spirig	Excipial [®] U Lipolotio	N058	3
Spirig/Noweda	Excipial [®] U Lipolotio	N061	1
Spirig	Excipial [®] U Lipolotio	N123	1
Fagron	Excipial [®] U Lipolotio	N168	2
spirig	Excipial [®] hydro cream	No14	1
Spirig Pharma	Excipial [®] hydro cream	N044	1
Spirig	Excipial [®] U Lipolotio	N086	1
AHD	Excipial [®] hydro cream	R011	1
spirig	Excipial [®] hydro cream	R013	2
Spirig	Excipial [®] hydro cream	R013	1
spirig	Excipial [®] U Lipolotio	R037	1
Spirig / Sanacorp	Excipial [®] U Lipolotio	R041	1
Galaderma	Excipial [®] U Lipolotio	R072	1
Galderma/Noweda	Excipial [®] U Lipolotio	R124	1
Galderma	Excipial [®] U Lipolotio	R111	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Galderma / Galderma	Excipial [®] U Lipolotio	R086	1
Spirig / Anzag	Excipial [®] hydro cream	813993/09	1
Spirig Pharma	Excipial [®] hydro cream	813993/10	1
Spirig Pharma	Excipial [®] hydro cream	LON014	1
Spirig / Anzag	Excipial [®] hydro cream	813993/10	1
Sanacorp	Excipial [®] hydro cream	LOTN011	1
spirig	Excipial [®] hydro cream	813993/10	1
Spirig	Excipial [®] hydro cream	K013	1
Spirig/Anzag	Excipial [®] hydro cream	LOTM041	1
Spirig/Gehe	Excipial [®] hydro cream	M012	1
spirig	Excipial [®] hydro cream	M014	2
Phoenix/Spirig	Excipial [®] hydro cream	M015	1
spirig	Excipial [®] hydro cream	M034	1
spirig	Excipial [®] hydro cream	M041	1
Spirig Pharma	Excipial [®] hydro cream	N011	1
Sanacorp	Excipial [®] hydro cream	N011	1
Spirig / Anzag	Excipial [®] hydro cream	N034	1
Phönix/Spirig	Excipial [®] hydro cream	N034	1
Spirig Pharma	Excipial [®] hydro cream	N013	1
Spirig/Anzag	Excipial [®] hydro cream	N035	1
Spirig	Excipial [®] hydro cream	N035	1
Alliance/Spirig	Excipial [®] hydro cream	N043	1
Spirig Pharma AG/GEHE	Excipial [®] hydro cream	R011	1
Spirig Pharma	Excipial [®] U Lipolotio	LOTK131	1
Spirig/ Noweda	Excipial [®] U Lipolotio	LOTN027	1
Spirig/ Noweda	Excipial [®] U Lipolotio	LOTN056	1
Spirig Pharma	Excipial [®] U Lipolotio	LOTN013	1
Spirig/AHD	Excipial [®] U Lipolotio	M085	1
Spirig	Excipial [®] U Lipolotio	M091	1
Spirig/Gehe	Excipial [®] U Lipolotio	M102	2
spirig	Excipial [®] U Lipolotio	M142	1
Spirig Pharma	Excipial [®] U Lipolotio	M134	1
Spirig/Gehe	Excipial [®] U Lipolotio	M146	1
Spirig / Anzag	Excipial [®] U Lipolotio	M182	1
Spirig/Gehe	Excipial [®] U Lipolotio	M182	1
Spirig/Noweda	Excipial [®] U Lipolotio	N027	1
Spirig / Anzag	Excipial [®] U Lipolotio	N032	1
Spirig / Anzag	Excipial [®] U Lipolotio	N041	1
Phoenix/Spirig	Excipial [®] U Lipolotio	N041	1
Spirig/AHD	Excipial [®] U Lipolotio	N058	1
spirig	Excipial [®] U Lipolotio	N061	1
Phoenix	Excipial [®] U Lipolotio	N105	1
spirig	Excipial [®] U Lipolotio	N013	1
Spirig/Noweda	Excipial [®] U Lipolotio	N032	1
spirig	Excipial [®] U Lipolotio	N086	1
Spirig/AHD	Excipial [®] U10 Lipolotio	M025	1
Galderma	Lipoderm [®] lotion	R044	2
Spirig Pharma / Allianc...	Lipoderm [®] lotion	R044	3
Galderma/Kehr	Lipoderm [®] lotion	S021	2
Galderma/AllianceHC	Lipoderm [®] lotion	S021	3
Spirig	Lipoderm [®] lotion	S041	2
Galderma	Lipoderm [®] lotion	S021	1
Galderma/Phoenix	Lipoderm [®] lotion	S041	1
Galderma/AllianceHC	Lipoderm [®] lotion	T011	2
Jenne	Lipoderm [®] lotion	T011	1
Galderma	Lipoderm [®] lotion	T011	1
Spirig Pharma/Phoenix	Lipoderm [®] lotion	T043	2
Spirig/AHD	Lipoderm [®] lotion	LOTN041	1
Spirig	Lipoderm [®] lotion	LOTR011	1
Spirig Pharma AG/Allian...	Lipoderm [®] lotion	N041	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Spirig Pharma AG/Allian...	Lipoderm [®] lotion	R011	4
Spirig	Lipoderm [®] lotion	R011	2
Spirig Pharma AG/Allian...	Lipoderm [®] lotion	R023	8
Spirig	Lipoderm [®] lotion	R023	2
Spirig/Phoenix	Lipoderm [®] lotion	M065	4
Spirig/AHD	Lipoderm [®] lotion	N041	1
Spirig Pharma AG	Lipoderm [®] lotion	N041	1
Spirig Pharma AG/Allian...	Lipoderm [®] lotion	N041	1
Hans Karrer/Gehe	Hans Karrer Lipolotion Micro...	17002	20
Jenapharm/Gehe	Neribas [®] cream	YY005HT	1
Jenapharm/Gehe	Neribas [®] cream	YY006KY	1
GPGrenzach Produktion/P...	Neribas [®] cream	YY01673	1
Phönix / Jenapharm	Neribas [®] cream	YY01FX3	1
Caelo	Neribas [®] cream	YY01X64	1
Intendis / Sanacorp	Neribas [®] cream	24105A	1
Jenapharm/Alliance	Neribas [®] cream	YY000C3	1
Jenapharm/Alliance	Neribas [®] cream	YY001HS	1
Bayer / Sanacorp	Neribas [®] cream	YY002L6	1
Intendis / Gehe	Neribas [®] cream	21076A	1
Intendis GmbH	Neribas [®] cream	24089A	1
Chiesi / Gehe	Neribas [®] cream	32097A	1
Jenapharm / Gehe	Neribas [®] cream	34104C	1

- 2455 spectra from 296 *Apo-Ident* customers from a total of 1306 batches from a further 105 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Hydrophilic ointments* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Hydrophilic ointments* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	2371	0	16 378
Type B	0	1972	117	20 395
Type C	0	687	51	2451

The substance/substance group *Hydrophilic ointments* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9530 %)	100.0000 % (> 99.7469 %)
Type B	100.0000 % (> 99.9564 %)	94.3992 % (> 94.2556 %)
Type C	100.0000 % (> 98.3661 %)	93.0894 % (> 92.6829 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30634	30634	0.00	25.82
30635	30635	0.00	27.73
30713	30713	0.00	27.58
30725	30725	0.00	34.47
30732	30732	0.00	33.49
30733	30733	0.00	31.19
30772	30772	0.00	22.04
30775	30775	0.00	32.49
30828	30828	0.00	23.87
30845	30845	0.00	17.61
30993	30993	0.00	35.01
31041	31041	0.00	18.52
31050	31050	0.00	32.17
31058	31058	0.00	28.78
31061	31061	0.00	33.47
31066	31066	0.00	31.07
31067	31067	0.00	28.54
31068	31068	0.00	26.73
31072	31072	0.00	31.60
31085	31085	0.00	32.37
31089	31089	0.00	21.85
31095	31095	0.00	31.25
31098	31098	0.00	22.95
31115	31115	0.00	17.42
31234	31234	0.00	27.34
31236	31236	0.00	29.90
31280	31280	0.00	31.89
31281	31281	0.00	29.73
31307	31307	0.00	15.00
31308	31308	0.00	28.94
31309	31309	0.00	28.30
31511	31511	0.00	23.56
31535	31535	0.00	27.93
31539	31539	0.00	27.58
31578	31578	0.00	21.25
31730	31730	0.00	25.80
31869	31869	0.00	15.11
31870	31870	0.00	30.25
31873	31873	0.00	24.92
31874	31874	0.00	25.97
32177	32177	0.00	24.14
32178	32178	0.00	27.94

continued on the next page

continued from previous page

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
32767	32767	0.00	31.92
32916	32916	0.00	32.26
32997	32997	0.00	31.14
33194	33194	0.00	25.55
33195	33195	0.00	27.20
33208	33208	0.00	24.34
33213	33213	0.00	25.79
33270	33270	0.00	31.12
33668	33668	0.00	32.91
33669	33669	0.00	31.19
33737	33737	0.00	21.06
33793	33793	0.00	15.56
33968	33968	0.00	32.29
33969	33969	0.00	30.27
34043	34043	0.00	25.71

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Immortelle oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30450-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Immortelle oil; Oleum helichrysum italicum

Special notes

When selecting the *Immortelle oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Immortelle oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Immortelle oil	L24113D-113148BAG90451	31138	40	not required
Taoasis	Immortelle oil	L24415DN-122092BAG90451	32069	40	not required
Taoasis	Immortelle oil	114717DNJ-137	33727	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Immortelle oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 4 reference samples of the substance/substance group *Immortelle oil*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Immortelle oil	L03111DN-102431	30450	40
Taoasis	Immortelle oil	L24415DN-122092BAG90451	32069 [†]	20
Taoasis	Immortelle oil	L17716DN-126770	32975	40
Taoasis	Immortelle oil	L17716DN-124955	33021	40

- 22 691 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 6 spectra from 3 *Apo-Ident* customers from 5 batches from the substance/substance group *Immortelle oil*.
- These include spectra of independent samples from 5 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Immortelle oil	L03111DN	1
Taoasis	Immortelle oil	L38914DN-118878	2
Taoasis	Immortelle oil	33111D	1
Fagron	Immortelle oil	L10712D-111673	1
Taoasis	Immortelle oil	L13613DN	1

- 3187 spectra from 304 *Apo-Ident* customers from a total of 1654 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Immortelle oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Immortelle oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	140	0	22 691
Type C	0	5	1	3187

The substance/substance group *Immortelle oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 95.7143 %)
Type C	100.0000 % (> 98.3707 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31138	31138	0.00	35.93
32069	32069	0.00	36.16
33727	33727	0.00	32.17

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Juniper oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31559-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Juniper oil; Juniperi aetheroleum; Oleum juniperi e baccarae

Special notes

When selecting the *Juniper oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Juniper oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Juniper oil	4155-116757	31559	40	not required
Taoasis	Juniper oil	4155-120900	31687	40	not required
Taoasis	Juniper oil	02JB0B-126047	32809	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Juniper oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *Juniper oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Juniper oil	4155-116757	31559 [†]	20
Taoasis	Juniper oil	4155-120900	31687 [†]	20
Taoasis	Juniper oil	02JB0B-127283	32932	40
Taoasis	Juniper oil	1713-128196	33651	40

- 22 711 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 2 spectra from 2 *Apo-Ident* customers from 2 batches from the substance/substance group *Juniper oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Juniper oil	155-118896	1
Caelo	Juniper oil	16160706	1

- 3191 spectra from 304 *Apo-Ident* customers from a total of 1657 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Juniper oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Juniper oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	80	40	22 711
Type C	0	2	0	3191

The substance/substance group *Juniper oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	66.6667 % (> 64.1667 %)
Type C	100.0000 % (> 98.3987 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31559	31559	0.00	23.52
31687	31687	0.00	25.33
32809	32809	0.00	21.82

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	La Roche-Posay Cold Cream Naturel
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30829-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

La Roche-Posay Cold Cream Naturel

Special notes

When selecting the *La Roche-Posay Cold Cream Naturel* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *La Roche-Posay Cold Cream Naturel*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
La Roche-Posay	La Roche-Posay C...	54J01P	30829	40	not required
La Roche-Posay	La Roche-Posay C...	54K31P	31074	40	not required
La Roche-Posay	La Roche-Posay C...	54LN1P	31765	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *La Roche-Posay Cold Cream Naturel*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *La Roche-Posay Cold Cream Naturel*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
La Roche-Posay	La Roche-Posay Cold Cream Na...	54LN1P	31765 [†]	20
La Roche-Posay	La Roche-Posay Cold Cream Na...	54M51P	33262	40
La Roche-Posay	La Roche-Posay Cold Cream Na...	54P61P	33984	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 3 spectra from 1 *Apo-Ident* customers from 3 batches from the substance/substance group *La Roche-Posay Cold Cream Naturel*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
La Roche-Posay	La Roche-Posay Cold Cream Na...	54M01P	1
La Roche-Posay	La Roche-Posay Cold Cream Na...	54M11P	1
La Roche-Posay	La Roche-Posay Cold Cream Na...	54M91P	1

- 3190 spectra from 304 *Apo-Ident* customers from a total of 1656 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *La Roche-Posay Cold Cream Naturel* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *La Roche-Posay Cold Cream Naturel* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	100	0	22 731
Type C	0	3	0	3190

The substance/substance group *La Roche-Posay Cold Cream Naturel* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3847 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30829	30829	0.00	87.60
31074	31074	0.00	85.10
31765	31765	0.00	82.14

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **La Roche-Posay Toleriane**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30827-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

La Roche-Posay Toleriane

Special notes

When selecting the *La Roche-Posay Toleriane* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *La Roche-Posay Toleriane*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
La Roche-Posay	La Roche-Posay T...	54J901	30827	40	not required
La Roche-Posay	La Roche-Posay T...	54K500	31094	40	not required
La Roche-Posay	La Roche-Posay T...	54N201	32563	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *La Roche-Posay Toleriane*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 80 spectra from 2 reference samples of the substance/substance group *La Roche-Posay Toleriane*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
La Roche-Posay	La Roche-Posay Toleriane	54N600	32759	40
La Roche-Posay	La Roche-Posay Toleriane	54P401	33983	40

- 22 751 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 2 spectra from 1 *Apo-Ident* customers from 2 batches from the substance/substance group *La Roche-Posay Toleriane*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	La Roche-Posay Toleriane	54J500	1
Roche/AHD	La Roche-Posay Toleriane	54J800	1

- 3191 spectra from 304 *Apo-Ident* customers from a total of 1657 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *La Roche-Posay Toleriane* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *La Roche-Posay Toleriane* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	80	0	22 751
Type C	0	2	0	3191

The substance/substance group *La Roche-Posay Toleriane* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9531 %)	100.0000 % (> 92.5000 %)
Type C	100.0000 % (> 98.3987 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30827	30827	0.00	28.79
31094	31094	0.00	24.15
32563	32563	0.00	23.94

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Lemon grass oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30661-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Lemon grass oil; Cymbopogonis aetheroleum

Special notes

When selecting the *Lemon grass oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Lemon grass oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Bombastus	Lemon grass oil	263571	30661	40	not required
Bombastus	Lemon grass oil	278923	31185	40	not required
Bombastus	Lemon grass oil	278526	31187	40	not required
Taoasis	Lemon grass oil	130930-116754	31549	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Lemon grass oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 589 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Lemon grass oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Lemon grass oil	130930-116754	31549 [†]	20
Taoasis	Lemon grass oil	1448-127048	32977	40
Bombastus	Lemon grass oil	294338	33012	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 8 spectra from 6 *Apo-Ident* customers from 8 batches from the substance/substance group *Lemon grass oil*.
- These include spectra of independent samples from 8 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Bombastus	Lemon grass oil	270423	1
Taoasis	Lemon grass oil	130930-115336	1
Taoasis	Lemon grass oil	8911011-121890	1
Taoasis	Lemon grass oil	FFL457-123568	1
Taoasis	Lemon grass oil	1663-125423	1
Primavera	Lemon grass oil	11401144	1
Taoasis	Lemon grass oil	13131-112982	1
Taoasis	Lemon grass oil	532897-111140	1

- 3185 spectra from 304 *Apo-Ident* customers from a total of 1651 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Lemon grass oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Lemon grass oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	96	4	22 731
Type C	0	7	1	3185

The substance/substance group *Lemon grass oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9530 %)	96.0000 % (> 93.0000 %)
Type C	100.0000 % (> 98.3672 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to

exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30661	30661	0.00	43.75
31185	31185	0.00	41.34
31187	31187	0.00	38.93
31549	31549	0.00	45.26

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Lime oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31544-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Lime oil; Oleum citrus aurantifolia

Special notes

When selecting the *Lime oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Lime oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Lime oil	059-117571	31544	40	not required
Taoasis	Lime oil	15506-1458	33874	40	not required
Taoasis	Lime oil	15506-1762	33942	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Lime oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 160 spectra from 4 reference samples of the substance/substance group *Lime oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Lime oil	059-117571	31544 [†]	20
Taoasis	Lime oil	059-117571	31654	60
Taoasis	Lime oil	867-124956	32800	40
Taoasis	Lime oil	867-124956	32976	40

- 22 671 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 1 spectra from 1 *Apo-Ident* customers from 1 batches from the substance/substance group *Lime oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Lime oil	867-124956	1

- 3192 spectra from 304 *Apo-Ident* customers from a total of 1658 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Lime oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Lime oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	135	25	22 671
Type C	0	1	0	3192

The substance/substance group *Lime oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9529 %)	84.3750 % (> 82.5000 %)
Type C	100.0000 % (> 98.4407 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested

reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31544	31544	0.00	70.57
33874	33874	0.00	64.72
33942	33942	0.00	65.34

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Liniment ointments**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30731-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Liniment ointments; Amciderm[®] Base cream; Excipial[®] U Hydrolotio; Linola[®] O/W cream; Linola[®] Sept; Nourivan[™] Antiox

Special notes

When selecting the *Liniment ointments* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Liniment ointments*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Spirig Pharma	Excipial [®] U Hydr...	M016	30731	40	not required
Dr. Wolff	Linola [®] O/W cream	238482	30834	40	not required
Spirig Pharma	Excipial [®] U Hydr...	M063	31073	40	not required
Dr. Wolff	Linola [®] Sept	334080	31079	40	not required

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Dr. Wolff	Linola [®] O/W cream	334010	31133	40	not required
Spirig Pharma	Excipial [®] U Hydr...	N013	31279	40	not required
Dr. Wolff	Linola [®] O/W cream	432480	31312	20	not required
Fagron	Nourivan [™] Antiox	14C11-T09-010959	31495	40	not required
Almirall	Amciderm [®] Base c...	313311	31509	40	not required
Spirig Pharma	Excipial [®] U Hydr...	R014	31536	40	not required
Almirall	Amciderm [®] Base c...	313311	31628	55	not required
Dr. Wolff	Linola [®] Sept	509361	31856	40	not required
Fagron	Nourivan [™] Antiox	1503258	31958	30	not required
Almirall	Amciderm [®] Base c...	650341	33502	40	not required
Fagron	Nourivan [™] Antiox	1703842	33789	30	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 575 spectra of 15 reference samples from the substance/substance group *Liniment ointments*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 14 different batches.
- 18 174 spectra from a total of 426 batches from further 146 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 480 spectra from 16 reference samples of the substance/substance group *Liniment ointments*.
- These include spectra of independent samples from 5 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Dr. Wolff	Linola [®] O/W cream	432480	31312 [†]	20
Fagron	Nourivan [™] Antiox	14C11-T09-010959	31495 [†]	20
Almirall	Amciderm [®] Base cream	313311	31509 [†]	20

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra
Spirig Pharma	Excipial [®] U Hydrolotio	R014	31536 [†]	20
Almirall	Amciderm [®] Base cream	313311	31628 [†]	20
Dr. Wolff	Linola [®] Sept	509361	31856 [†]	20
Fagron	Nourivan [™] Antiox	1503258	31958 [†]	30
Almirall	Amciderm [®] Base cream	538331	32758	40
Dr. Wolff	Linola [®] Sept	603190	32760	40
Fagron	Nourivan [™] Antiox	1503258	32778	40
Fagron	Nourivan [™] Antiox	1503258	32822	40
Spirig Pharma	Excipial [®] U Hydrolotio	S061	33206	40
Dr. Wolff	Linola [®] Sept	603190	33257	40
Dr. Wolff	Linola [®] O/W cream	610960	33259	40
Fagron	Nourivan [™] Antiox	1703842	33789 [†]	10
Spirig Pharma	Excipial [®] U Hydrolotio	V052	33981	40

- 22351 spectra from a total of 537 batches from further 198 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 106 spectra from 31 *Apo-Ident* customers from 57 batches from the substance/substance group *Liniment ointments*.
- These include spectra of independent samples from 53 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Excipial [®] U Hydrolotio	L0TR092	1
Galderma/Noweda	Excipial [®] U Hydrolotio	R054	1
Spirig Pharma AG/ NOWEDA	Excipial [®] U Hydrolotio	L0TN022	1
Galderma	Excipial [®] U Hydrolotio	L0TR054	1
Spirig	Excipial [®] U Hydrolotio	N013	1
Spirig	Excipial [®] U Hydrolotio	N013	1
Phoenix/Spirig	Excipial [®] U Hydrolotio	M016	1
Dr. Wolff	Linola [®] O/W cream	610960	1
Dr. Wolff	Linola [®] O/W cream	505200	4
Dr. August Wolff	Linola [®] O/W cream	505560	1
Dr. Wolff	Linola [®] O/W cream	510792	1
Dr. Wolff	Linola [®] O/W cream	602190	3
Dr. Wolff	Linola [®] O/W cream	602300	1
Sanofi/Noweda	Linola [®] O/W cream	602300	1
Caelo	Linola [®] O/W cream	602310	1
Dr. August Wolff	Linola [®] O/W cream	605560	1
Dr. Wolff/Gehe	Linola [®] O/W cream	606100	1
Dr. Wolff GmbH / Phoenix	Linola [®] O/W cream	606100	1

continued on the next page

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

continued from previous page

Supplier	Substance	Batch	Spectra
Dr. August Wolff	Linola [®] O/W cream	607140	1
Dr. August Wolff	Linola [®] O/W cream	608900	1
Dr. Wolff/Gehe	Linola [®] O/W cream	138350	1
Dr. Wolff	Linola [®] O/W cream	434430	3
Dr. Wolff	Linola [®] O/W cream	438200	2
Dr. Wolff	Linola [®] O/W cream	438200	1
Dr. Wolff/Gehe	Linola [®] O/W cream	438360	1
Dr. Wolff	Linola [®] O/W cream	504100	1
Dr. Wolff/Noweda	Linola [®] O/W cream	505280	1
Dr. Wolff/Malteser	Linola [®] O/W cream	338990	1
Wolff / Noweda	Linola [®] O/W cream	431120	4
Wolff / Gehe	Linola [®] O/W cream	431130	3
Dr. Wolff/Noweda	Linola [®] O/W cream	431130	2
Dr. Wolff	Linola [®] O/W cream	431200	3
Dr. August Wolff/Ebert	Linola [®] O/W cream	431250	1
Dr. Wolff	Linola [®] O/W cream	432480	1
Dr. Wolff/Gehe	Linola [®] O/W cream	432490	2
Dr. Wolff	Linola [®] O/W cream	432490	1
Dr. August Wolff/Ebert	Linola [®] O/W cream	432970	1
wolff	Linola [®] O/W cream	434320	1
Dr. Wolff/Gehe	Linola [®] O/W cream	434320	2
Dr. Wolff	Linola [®] O/W cream	435530	1
AHD	Linola [®] O/W cream	437110	1
Dr. Wolff	Linola [®] O/W cream	437120	1
Dr. Wolff/Noweda	Linola [®] O/W cream	438350	2
Caelo	Linola [®] O/W cream	438350	1
Dr. August Wolff	Linola [®] O/W cream	233140	1
Dr. Wolff/Noweda	Linola [®] O/W cream	233200	1
Dr. Wolff	Linola [®] O/W cream	233200	1
DR. WOLFF	Linola [®] O/W cream	234192	1
Dr. Wolff/ Anzag	Linola [®] O/W cream	234193	1
Caelo	Linola [®] O/W cream	236281	1
Dr. Wolff	Linola [®] O/W cream	236290	1
Caelo	Linola [®] O/W cream	238484	1
Dr. Wolff	Linola [®] O/W cream	238482	1
Dr. Wolff	Linola [®] O/W cream	238484	1
Caelo	Linola [®] O/W cream	238520	2
Dr. Wolff	Linola [®] O/W cream	238520	2
Dr. Wolff/Gehe	Linola [®] O/W cream	238520	1
Dr. Wolff/Anzag	Linola [®] O/W cream	238520	2
Caelo	Linola [®] O/W cream	238530	2
Dr. Wolff	Linola [®] O/W cream	238530	1
Dr. Wolff	Linola [®] O/W cream	238530	4
Caelo	Linola [®] O/W cream	239181	1
Dr. Wolff/Malteser	Linola [®] O/W cream	330751	1
Caelo	Linola [®] O/W cream	330760	2
chiesi / Noweda	Linola [®] O/W cream	331460	1
Wolff/Sanacorp	Linola [®] O/W cream	333630	2
Phoenix/Wolff	Linola [®] O/W cream	333640	1
Dr. Wolff	Linola [®] O/W cream	333640	1
Dr. August Wolff/Ebert	Linola [®] O/W cream	333640	2
Euro OTC	Linola [®] O/W cream	333730	1
Dr. Wolff/Malteser	Linola [®] O/W cream	333880	1
Dr. Wolff	Linola [®] O/W cream	334003	1
Dr. Wolff / Sanacorp	Linola [®] O/W cream	338990	3
Dr. Wolff	Linola [®] O/W cream	435080	1
Dr. Wolf/GEHE	Linola [®] O/W cream	437110	1

- 3087 spectra from 304 *Apo-Ident* customers from a total of 1606 batches from a further 120 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Liniment ointments* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Liniment ointments* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	575	0	18 174
Type B	0	417	63	22 244
Type C	0	103	3	3087

The substance/substance group *Liniment ointments* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9531 %)	100.0000 % (> 98.9565 %)
Type B	100.0000 % (> 99.9542 %)	86.8750 % (> 86.2500 %)
Type C	100.0000 % (> 98.3575 %)	97.1698 % (> 94.3396 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30731	30731	0.00	12.37
30834	30834	0.00	35.78
31073	31073	0.00	11.96
31079	31079	0.00	38.05
31133	31133	0.00	35.59
31279	31279	0.00	11.87
31312	31312	0.00	32.05
31495	31495	0.00	21.62
31509	31509	0.00	32.90
31536	31536	0.00	12.16

continued on the next page

continued from previous page

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31628	31628	0.00	33.24
31856	31856	0.00	36.45
31958	31958	0.00	22.28
33502	33502	0.00	35.03
33789	33789	0.00	20.45

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Linola[®] fat creme / Linola[®] H fat N / Alfason[®] R...**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30724-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Linola[®] fat creme / Linola[®] H fat N / Alfason[®] Repair; Alfason[®] Repair; Linola[®] fat cream;
Linola[®] H fat N

Special notes

When selecting the *Linola[®] fat creme / Linola[®] H fat N / Alfason[®] Repair* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Linola[®] fat creme / Linola[®] H fat N / Alfason[®] Repair*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Astellas Pharma	Alfason [®] Repair	12F18/77	30724	50	not required
Dr. Wolff	Linola [®] fat cream	235260	30771	40	not required
Astellas Pharma	Alfason [®] Repair	13E16/76	31069	40	not required
Dr. Wolff	Linola [®] H fat N	333080	31092	40	not required

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Dr. Wolff	Linola [®] fat cream	337280	31096	40	not required
Astellas Pharma	Alfason [®] Repair	16G60/75	32993	40	not required
Dr. Wolff	Linola [®] H fat N	611060	33211	40	not required
Dr. Wolff	Linola [®] fat cream	611011	33243	40	not required
Dr. Wolff	Linola [®] fat cream	612211	33260	40	not required
LEO Pharma GmbH	Alfason [®] Repair	17E20/75	33790	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 410 spectra of 10 reference samples from the substance/substance group *Linola[®] fat creme / Linola[®] H fat N / Alfason[®] Repair*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 10 different batches.
- 18 339 spectra from a total of 429 batches from further 148 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 205 spectra from 6 reference samples of the substance/substance group *Linola[®] fat creme / Linola[®] H fat N / Alfason[®] Repair*.
- These include spectra of independent samples from 4 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Dr. Wolff	Linola [®] H fat N	231610	30576	40
Astellas Pharma	Alfason [®] Repair	12F18/77	30724 [†]	5
Dr. Wolff	Linola [®] H fat N	603230	32560	40
Dr. Wolff	Linola [®] fat cream	603820	32561	40
Astellas Pharma	Alfason [®] Repair	15K07/77	32562	40
Dr. Wolff	Linola [®] H fat N	611060	33038	40

- 22 626 spectra from a total of 543 batches from further 200 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 20 spectra from 14 *Apo-Ident* customers from 18 batches from the substance/substance group *Linola® fat creme / Linola® H fat N / Alfason® Repair*.
- These include spectra of independent samples from 18 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Phoenix	Alfason® Repair	15/03/75	1
Phönix	Alfason® Repair	15J09/75	1
Astellas	Alfason® Repair	13H02/76	1
Sanofi/Noweda	Linola® fat cream	501000	1
Wolff/Noweda	Linola® fat cream	501230	1
dr.wolff	Linola® fat cream	337770	1
Dr. Wolff / Gehe	Linola® fat cream	337860	1
Wolff / Wolff	Linola® fat cream	431190	1
Dr. Wolff/NOW	Linola® fat cream	435160	1
Dr. Wolff	Linola® fat cream	437270	1
Caelo	Linola® fat cream	435161	1
Löwen City-Apotheke Bar...	Linola® fat cream	231060	1
Caelo	Linola® fat cream	438070	1
Dr. Wolff /Anzag	Linola® fat cream	237710	1
Caelo	Linola® fat cream	238540	1
Dr. Wolff	Linola® fat cream	237920	1
Phönix	Linola® fat cream	238540	1
dr.wolff	Linola® fat cream	337760	1
Wolff/Phönix	Linola® fat cream	337770	1
Dr. Wolff	Linola® fat cream	432591	1

- 3173 spectra from 304 *Apo-Ident* customers from a total of 1641 batches from a further 120 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Linola® fat creme / Linola® H fat N / Alfason® Repair* can clearly be distinguished from all other substances using NIR spectroscopy

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Linola® fat creme* / *Linola® H fat N* / *Alfason® Repair* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	410	0	18 339
Type B	0	203	2	22 601
Type C	0	19	1	3173

The substance/substance group *Linola® fat creme* / *Linola® H fat N* / *Alfason® Repair* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9532 %)	100.0000 % (> 98.5366 %)
Type B	100.0000 % (> 99.9536 %)	99.0244 % (> 97.5610 %)
Type C	100.0000 % (> 98.3609 %)	95.0000 % (> 80.0000 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30724	30724	0.00	20.08
30771	30771	0.00	19.73
31069	31069	0.00	8.79
31092	31092	0.00	21.30
31096	31096	0.00	23.05
32993	32993	0.00	14.92
33211	33211	0.00	26.67
33243	33243	0.00	25.37
33260	33260	0.00	24.44
33790	33790	0.00	20.03

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Lipolotio urea 5% F body lotion**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30973-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Lipolotio urea 5% F body lotion

Special notes

When selecting the *Lipolotio urea 5% F body lotion* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Lipolotio urea 5% F body lotion*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Allergika	Lipolotio urea 5...	1205050	30973	80	not required
Allergika	Lipolotio urea 5...	13112021	31766	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 2 reference samples from the substance/substance group *Lipolotio urea 5% F body lotion*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 629 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Lipolotio urea 5% F body lotion*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Allergika	Lipolotio urea 5% F body lot...	13112021	31766 [†]	20
Allergika	Lipolotio urea 5% F body lot...	1502011	32766	40
Allergika	Lipolotio urea 5% F body lot...	1502011	33482	40

- 22 731 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Lipolotio urea 5% F body lotion*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Lipolotio urea 5% F body lotion* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Lipolotio urea 5% F body lotion* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	100	0	22 731
Type C	0	0	0	3193

The substance/substance group *Lipolotio urea 5% F body lotion* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30973	30973	0.00	30.66
31766	31766	0.00	42.30

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Lygal® head ointment N 3%**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31049-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Lygal® head ointment N 3%

Special notes

When selecting the *Lygal® head ointment N 3%* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Lygal® head ointment N 3%*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taurus	Lygal® head oint...	124311	31049	40	not required
Taurus	Lygal® head oint...	143710	31763	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Lygal[®] head ointment N 3%*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Lygal[®] head ointment N 3%*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taurus	Lygal [®] head ointment N 3%	143710	31763 [†]	20
Taurus	Lygal [®] head ointment N 3%	160732	32763	40
Taurus	Lygal [®] head ointment N 3%	160732	33261	40

- 22 731 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 2 spectra from 2 *Apo-Ident* customers from 2 batches from the substance/substance group *Lygal[®] head ointment N 3%*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
TaurusPharma/Noweda	Lygal [®] head ointment N 3%	141703	1
Taurus Pharma	Lygal [®] head ointment N 3%	134507	1

- 3191 spectra from 304 *Apo-Ident* customers from a total of 1657 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Lygal[®] head ointment N 3%* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Lygal[®] head ointment N 3%* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	100	0	22 731
Type C	0	2	0	3191

The substance/substance group *Lygal[®] head ointment N 3%* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3987 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31049	31049	0.00	38.34
31763	31763	0.00	38.09

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Lygal[®] ointment base**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31014-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Lygal[®] ointment base

Special notes

When selecting the *Lygal[®] ointment base* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Lygal[®] ointment base*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taurus	Lygal [®] ointment ...	121302P	31014	40	not required
Taurus	Lygal [®] ointment ...	154841	32749	40	not required
Taurus	Lygal [®] ointment ...	164201	32772	40	not required
Taurus	Lygal [®] ointment ...	164201	33244	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Lygal[®] ointment base*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 589 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 3 reference samples of the substance/substance group *Lygal[®] ointment base*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taurus	Lygal [®] ointment base	124904P	31704	60
Taurus	Lygal [®] ointment base	164201	33297	40
Almirall	Lygal [®] ointment base	174104	33988	40

- 22 691 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 18 spectra from 6 *Apo-Ident* customers from 10 batches from the substance/substance group *Lygal[®] ointment base*.
- These include spectra of independent samples from 9 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taurus	Lygal [®] ointment base	133402	1
Taurus/ Phönix	Lygal [®] ointment base	133401T	1
TAUROS PHARMA	Lygal [®] ointment base	141601P	1
Caelo	Lygal [®] ointment base	141601P	1
Taurus	Lygal [®] ointment base	141601T	1
TAURUS PHARMA	Lygal [®] ointment base	141602T	1
Taurus	Lygal [®] ointment base	141603	2
TAURUS PHARMA	Lygal [®] ointment base	154841	1
TAUROS PHARMA	Lygal [®] ointment base		1
TAUROS PHARMA	Lygal [®] ointment base	124803	1
Taurus	Lygal [®] ointment base	124803	2
Taurus Pharma	Lygal [®] ointment base	133401P	1
Taurus Pharma	Lygal [®] ointment base	133401T	2
Taurus	Lygal [®] ointment base	133401T	2

- 3175 spectra from 304 *Apo-Ident* customers from a total of 1649 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Lygal[®] ointment base* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Lygal[®] ointment base* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	116	24	22 691
Type C	0	6	12	3175

The substance/substance group *Lygal[®] ointment base* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9530 %)	82.8571 % (> 80.7143 %)
Type C	100.0000 % (> 98.3613 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several

new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31014	31014	0.00	21.73
32749	32749	0.00	26.69
32772	32772	0.00	33.51
33244	33244	0.00	29.08

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Mandarin oil, green**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 33798-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Mandarin oil, green

Special notes

When selecting the *Mandarin oil, green* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Mandarin oil, green*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Primavera	Mandarin oil, gr...	00431E25	33798	40	not required
Primavera	Mandarin oil, gr...	00381H25	33806	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Mandarin oil, green*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 40 spectra from 1 reference samples of the substance/substance group *Mandarin oil, green*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Primavera	Mandarin oil, green	00425M25	33960	40

- 22 791 spectra from a total of 548 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Mandarin oil, green*.

- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Mandarin oil, green* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Mandarin oil, green* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	40	0	22 791
Type C	0	0	0	3193

The substance/substance group *Mandarin oil, green* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9535 %)	100.0000 % (> 85.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
33798	33798	0.00	29.95
33806	33806	0.00	30.58

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Manuka oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31545-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Manuka oil; Oleum leptospermum scoparium

Special notes

When selecting the *Manuka oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Manuka oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Manuka oil	1235-116650	31545	40	not required
Taoasis	Manuka oil	14286-119821	31655	40	not required
Taoasis	Manuka oil	83579-2408	33994	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Manuka oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *Manuka oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Manuka oil	1235-116650	31545 [†]	20
Taoasis	Manuka oil	14286-119821	31655 [†]	20
Taoasis	Manuka oil	42798-125299	32799	40
Taoasis	Manuka oil	42798-125299	33198	40

- 22 711 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 5 spectra from 1 *Apo-Ident* customers from 4 batches from the substance/substance group *Manuka oil*.
- These include spectra of independent samples from 4 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Manuka oil	23694-122415	1
Taoasis	Manuka oil	23694-123569	1
Taoasis	Manuka oil	36968-121509	1
Taoasis	Manuka oil	42798-125299	2

- 3188 spectra from 304 *Apo-Ident* customers from a total of 1655 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Manuka oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Manuka oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	120	0	22 711
Type C	0	5	0	3188

The substance/substance group *Manuka oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 95.0000 %)
Type C	100.0000 % (> 98.3735 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31545	31545	0.00	37.73
31655	31655	0.00	37.49
33994	33994	0.00	28.94

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Marjoram oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30454-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Marjoram oil; Majoranae aetheroleum; Oleum Majoranae

Special notes

When selecting the *Marjoram oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Marjoram oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Marjoram oil	130312-111991BAG90451	31137	40	not required
Taoasis	Marjoram oil	3701004-121123	32067	40	not required
Caelo	Marjoram oil	16170005	32819	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Marjoram oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 4 reference samples of the substance/substance group *Marjoram oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Marjoram oil	29293-98717	30454	40
Taoasis	Marjoram oil	3701004-121123	32067 [†]	20
Taoasis	Marjoram oil	1838-125624	32478	40
Caelo	Marjoram oil	16170005	32919	40

- 22 691 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 25 spectra from 8 *Apo-Ident* customers from 16 batches from the substance/substance group *Marjoram oil*.
- These include spectra of independent samples from 16 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Marjoram oil	12041602	7
Taoasis	Marjoram oil	130312-111347	1
Taoasis	Marjoram oil	130312-111870	1
Caelo	Marjoram oil	13041601	1
Taoasis	Marjoram oil	130910-116960	1
Taoasis	Marjoram oil	130910-117430	1
Caelo	Marjoram oil	13121301	1
Caelo	Marjoram oil	15051505	1
Taoasis	Marjoram oil	1631-122347	1
Taoasis	Marjoram oil	1631-124621	1
Taoasis/Gehe	Marjoram oil	29293-98717	1
Taoasis	Marjoram oil	3701004-119994	2
Caelo	Marjoram oil	13041603	2
Caelo	Marjoram oil	13041606	1
Caelo	Marjoram oil	14198304	1
Taoasis	Marjoram oil	29293-98717	1
Bombastus	Marjoram oil	278279	1

- 3168 spectra from 304 *Apo-Ident* customers from a total of 1643 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Marjoram oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Marjoram oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	138	2	22 691
Type C	0	17	8	3168

The substance/substance group *Marjoram oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	98.5714 % (> 96.4286 %)
Type C	100.0000 % (> 98.3600 %)	68.0000 % (> 56.0000 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31137	31137	0.00	74.84
32067	32067	0.00	46.71
32819	32819	0.00	53.18

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Matricaria chamomilla oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 33811-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Matricaria chamomilla oil; Chamomile oil; German chamomile oil

Special notes

When selecting the *Matricaria chamomilla oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Matricaria chamomilla oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Primavera	Matricaria chamo...	00888G25	33811	40	not required
Taoasis	Matricaria chamo...	934	33825	40	not required
Taoasis	Matricaria chamo...	2132-1981	33952	40	not required
Primavera	Matricaria chamo...	00715L25	33963	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Matricaria chamomilla oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 589 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 40 spectra from 1 reference samples of the substance/substance group *Matricaria chamomilla oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Matricaria chamomilla oil	100477-850	33824	40

- 22 791 spectra from a total of 548 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Matricaria chamomilla oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Matricaria chamomilla oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Matricaria chamomilla oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	40	0	22 791
Type C	0	0	0	3193

The substance/substance group *Matricaria chamomilla oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9535 %)	100.0000 % (> 85.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
33811	33811	0.00	51.19
33825	33825	0.00	18.90
33952	33952	0.00	21.80
33963	33963	0.00	51.66

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Matricaria oil, morrocan**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30844-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Matricaria oil, morrocan; Oleum ormenis multicaulis; Ormenis multicaulis oleum

Special notes

When selecting the *Matricaria oil, morrocan* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Matricaria oil, morrocan*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Matricaria oil, ...	140124-116959	31579	40	not required
Taoasis	Matricaria oil, ...	6121017-122184	31841	40	not required
Taoasis	Matricaria oil, ...	1629-124596	32269	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Matricaria oil, morrocan*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 160 spectra from 5 reference samples of the substance/substance group *Matricaria oil, morrocan*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Matricaria oil, morrocan	6982-1074109BAC90451	30844	40
Taoasis	Matricaria oil, morrocan	140124-116959	31579 [†]	20
Taoasis	Matricaria oil, morrocan	6121017-122184	31841 [†]	20
Taoasis	Matricaria oil, morrocan	1629-125295	32480	40
Taoasis	Matricaria oil, morrocan	1814-126707	33020	40

- 22 671 spectra from a total of 544 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B*

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 2 spectra from 1 *Apo-Ident* customers from 2 batches from the substance/substance group *Matricaria oil, morrocan*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Matricaria oil, morrocan	6159-93239	1
Taoasis	Matricaria oil, morrocan	6631-104465	1

- 3191 spectra from 304 *Apo-Ident* customers from a total of 1657 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Matricaria oil, morrocan* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Matricaria oil, morrocan* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	159	1	22 671
Type C	0	1	1	3191

The substance/substance group *Matricaria oil, morrocan* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9529 %)	99.3750 % (> 97.5000 %)
Type C	100.0000 % (> 98.3987 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31579	31579	0.00	38.77
31841	31841	0.00	41.47
32269	32269	0.00	54.46

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Matricaria oil, roman
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	31254-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Matricaria oil, roman; Oleum anthemis nobilis

Special notes

When selecting the *Matricaria oil, roman* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Matricaria oil, roman*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Matricaria oil, ...	34561-112874	31254	40	not required
Taoasis	Matricaria oil, ...	1850119-119823	31686	40	not required
Taoasis	Matricaria oil, ...	2033-843	33989	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Matricaria oil, roman*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Matricaria oil, roman*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Matricaria oil, roman	1850119-119823	31686 [†]	20
Taoasis	Matricaria oil, roman	8941-125875	32796	40
Taoasis	Matricaria oil, roman	8941-125875	32978	40

- 22 731 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 7 spectra from 1 *Apo-Ident* customers from 7 batches from the substance/substance group *Matricaria oil, roman*.
- These include spectra of independent samples from 7 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Matricaria oil, roman	1511-122183	1
Taoasis	Matricaria oil, roman	1850119-116753	1
Taoasis	Matricaria oil, roman	1850119-118138	1
Taoasis	Matricaria oil, roman	1850119-119451	1
Taoasis	Matricaria oil, roman	1850119-121467	1
Taoasis	Matricaria oil, roman	1638-123706	1
Taoasis	Matricaria oil, roman	8941-125875	1

- 3186 spectra from 304 *Apo-Ident* customers from a total of 1652 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Matricaria oil, roman* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Matricaria oil, roman* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	73	27	22 731
Type C	0	6	1	3186

The substance/substance group *Matricaria oil, roman* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	73.0000 % (> 70.0000 %)
Type C	100.0000 % (> 98.3687 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to

exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31254	31254	0.00	32.67
31686	31686	0.00	31.95
33989	33989	0.00	31.27

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Myrrh oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30748-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Myrrh oil; Oleum myrrha

Special notes

When selecting the *Myrrh oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Myrrh oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Myrrh oil	30007-120958	31692	40	not required
Taoasis	Myrrh oil	41197-124418BAG90451	32271	40	not required
Taoasis	Myrrh oil	2206-2428	33992	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Myrrh oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 4 reference samples of the substance/substance group *Myrrh oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Myrrh oil	22920-105398BAG90451	30748	40
Taoasis	Myrrh oil	30007-120958	31692 [†]	20
Taoasis	Myrrh oil	41197-126694	33050	40
Taoasis	Myrrh oil	41197-126694	33051	40

- 22 691 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 1 spectra from 1 *Apo-Ident* customers from 1 batches from the substance/substance group *Myrrh oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Myrrh oil	2067-107953	1

- 3192 spectra from 304 *Apo-Ident* customers from a total of 1658 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Myrrh oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Myrrh oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	100	40	22 691
Type C	0	0	1	3192

The substance/substance group *Myrrh oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	71.4286 % (> 69.2857 %)
Type C	100.0000 % (> 98.4407 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested

reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31692	31692	0.00	152.31
32271	32271	0.00	207.75
33992	33992	0.00	61.29

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Myrtle oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 33797-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Myrtle oil

Special notes

When selecting the *Myrtle oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Myrtle oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Primavera	Myrtle oil	00401D25	33797	40	not required
Taoasis	Myrtle oil	013	33830	40	not required
Primavera	Myrtle oil	00877L25	33957	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Myrtle oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 40 spectra from 1 reference samples of the substance/substance group *Myrtle oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Primavera	Myrtle oil	00971J25	33807	40

- 22 791 spectra from a total of 548 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Myrtle oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Myrtle oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Myrtle oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	40	0	22 791
Type C	0	0	0	3193

The substance/substance group *Myrtle oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9535 %)	100.0000 % (> 85.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the [Mahalanobis distance](#) to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
33797	33797	0.00	33.07
33830	33830	0.00	27.91
33957	33957	0.00	33.73

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Neroli oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31260-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Neroli oil; Oleum citrus aurantium var. amara; Oleum neroli

Special notes

When selecting the *Neroli oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Neroli oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Neroli oil	91029-115714	31260	40	not required
Taoasis	Neroli oil	6142-120057BAG90451	31695	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Neroli oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 4 reference samples of the substance/substance group *Neroli oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Neroli oil	6142-120057BAG90451	31695 [†]	20
Taoasis	Neroli oil	125052-124958	32805	40
Taoasis	Neroli oil	125052-124958	33042	40
Taoasis	Neroli oil	071704-1966	33993	40

- 22 691 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Neroli oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Neroli oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Neroli oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	60	80	22 691
Type C	0	0	0	3193

The substance/substance group *Neroli oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9530 %)	42.8571 % (> 40.7143 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31260	31260	0.00	46.06
31695	31695	0.00	47.67

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Neuroderm [®] care lotion / moisturising cream
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30722-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Neuroderm[®] care lotion / moisturising cream; Neuroderm[®] care lotion; Neuroderm[®] moisturising cream

Special notes

When selecting the *Neuroderm[®] care lotion / moisturising cream* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Neuroderm[®] care lotion / moisturising cream*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Infectopharm	Neuroderm [®] moist...	S041214.1	30722	40	not required
Infectopharm	Neuroderm [®] care ...	S041204.1	31070	40	not required
Infectopharm	Neuroderm [®] moist...	S021307.1	31102	40	not required
Infectopharm	Neuroderm [®] moist...	S101428.1	31761	40	not required
Infectopharm	Neuroderm [®] care ...	S031505.1	31772	40	not required
Infectopharm	Neuroderm [®] care ...	S071706.1	34044	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 240 spectra of 6 reference samples from the substance/substance group *Neuroderm[®] care lotion / moisturising cream*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 6 different batches.
- 18 509 spectra from a total of 433 batches from further 149 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 200 spectra from 6 reference samples of the substance/substance group *Neuroderm[®] care lotion / moisturising cream*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Infectopharm	Neuroderm [®] moisturising cream	S101428.1	31761 [†]	20
Infectopharm	Neuroderm [®] care lotion	S031505.1	31772 [†]	20
Infectopharm	Neuroderm [®] care lotion	S111514.1	32740	40
Infectopharm	Neuroderm [®] moisturising cream	S051612.1	32905	40
Infectopharm	Neuroderm [®] moisturising cream	S051612.1	32906	40
Infectopharm	Neuroderm [®] care lotion	S081610.1	33286	40

- 22 631 spectra from a total of 544 batches from further 201 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 16 spectra from 7 *Apo-Ident* customers from 15 batches from the substance/substance group *Neuroderm® care lotion / moisturising cream*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Infectopharm	Neuroderm® moisturising cream		6
Infectopharm/Kehr	Neuroderm® moisturising cream		1
Infectopharm/Gehe	Neuroderm® moisturising cream	S0315081	1
Infectopharm/Fiebig	Neuroderm® moisturising cream		1
Bombastus	Neuroderm® moisturising cream		1
Noweda	Neuroderm® moisturising cream		1
Infectopharm/Noweda	Neuroderm® care lotion		2
Sanacorp	Neuroderm® moisturising cream		1
Infectpharm/ Noweda	Neuroderm® moisturising cream		1
infectopharm	Neuroderm® moisturising cream		1

- 3177 spectra from 304 *Apo-Ident* customers from a total of 1644 batches from a further 120 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Neuroderm® care lotion / moisturising cream* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Neuroderm® care lotion / moisturising cream* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	240	0	18 509
Type B	0	200	0	22 510
Type C	0	16	0	3175

The substance/substance group *Neuroderm® care lotion / moisturising cream* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.5000 %)
Type B	100.0000 % (> 99.9541 %)	100.0000 % (> 97.0000 %)
Type C	100.0000 % (> 98.3713 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30722	30722	0.00	30.97
31070	31070	0.00	18.91
31102	31102	0.00	25.38
31761	31761	0.00	27.76
31772	31772	0.00	19.30
34044	34044	0.00	21.26

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Neuroderm[®] moisturising cream lipo**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30721-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Neuroderm[®] moisturising cream lipo

Special notes

When selecting the *Neuroderm[®] moisturising cream lipo* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Neuroderm[®] moisturising cream lipo*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Infectopharm	Neuroderm [®] moist...	S031202.1	30721	40	not required
Infectopharm	Neuroderm [®] moist...	S031506.1	31672	40	not required
Infectopharm	Neuroderm [®] moist...	S051723.1	34045	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Neuroderm[®] moisturising cream lipo*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Neuroderm[®] moisturising cream lipo*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Infectopharm	Neuroderm [®] moisturising crea...	S031506.1	31672 [†]	20
Infectopharm	Neuroderm [®] moisturising crea...	S071617.2	32739	40
Infectopharm	Neuroderm [®] moisturising crea...	S021706.1	33287	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 2 spectra from 2 *Apo-Ident* customers from 2 batches from the substance/substance group *Neuroderm[®] moisturising cream lipo*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Infectopharm	Neuroderm [®] moisturising crea...		1
Infectopharm/AHD	Neuroderm [®] moisturising crea...		1

- 3191 spectra from 304 *Apo-Ident* customers from a total of 1658 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Neuroderm[®] moisturising cream lipo* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Neuroderm[®] moisturising cream lipo* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	100	0	22 731
Type C	0	2	0	3191

The substance/substance group *Neuroderm[®] moisturising cream lipo* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3987 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30721	30721	0.00	30.62
31672	31672	0.00	24.25
34045	34045	0.00	30.45

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Nicotine**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31019-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Nicotine

Special notes

When selecting the *Nicotine* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Nicotine*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Roth	Nicotine	152183017	31019	20	1402687
Roth	Nicotine	194210173	31508	40	1410523
Roth	Nicotine	194210173	31518	40	not required
Roth	Nicotine	116210173	31948	30	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 130 spectra of 4 reference samples from the substance/substance group *Nicotine*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 619 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 105 spectra from 5 reference samples of the substance/substance group *Nicotine*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Roth	Nicotine	451178717	30376	10
Roth	Nicotine	194210173	31508 [†]	20
Roth	Nicotine	194210173	31518 [†]	20
Roth	Nicotine	116210173	31948 [†]	15
Roth	Nicotine	137210173	33291	40

- 22 726 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Nicotine*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Nicotine* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Nicotine* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	130	0	18 619
Type B	0	105	0	22 726
Type C	0	0	0	3193

The substance/substance group *Nicotine* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9535 %)	100.0000 % (> 95.3846 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.2857 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31019	31019	0.00	151.38
31508	31508	0.00	149.88
31518	31518	0.00	150.58
31948	31948	0.00	151.00

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Ointment ash basis / Neribas [®]
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30774-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Ointment ash basis / Neribas[®]; Asche Basis[®] ointment; Neribas[®] ointment

Special notes

When selecting the *Ointment ash basis / Neribas[®]* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Ointment ash basis / Neribas[®]*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Chiesi	Asche Basis [®] oin...	21038A	30774	40	not required
Chiesi	Asche Basis [®] oin...	21038A	30833	40	not required
Bayer	Neribas [®] ointment	21053B	30969	40	not required
Bayer	Neribas [®] ointment	21053B	30994	40	not required
Chiesi	Asche Basis [®] oin...	21038A	31080	40	not required
Jenapharm	Neribas [®] ointment	31064C	31093	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 240 spectra of 6 reference samples from the substance/substance group *Ointment ash basis / Neribas®*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 509 spectra from a total of 436 batches from further 149 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 240 spectra from 6 reference samples of the substance/substance group *Ointment ash basis / Neribas®*.
- These include spectra of independent samples from 4 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Jenapharm	Neribas [®] ointment	YY00TJ0	32564	40
Chiesi	Asche Basis [®] ointment	61067A	32590	40
Chiesi	Asche Basis [®] ointment	61067A	32924	40
Chiesi	Asche Basis [®] ointment	61067A	32927	40
Jenapharm	Neribas [®] ointment	YY016EC	33288	40
Chiesi	Asche Basis [®] ointment	17072A	33840	40

- 22 591 spectra from a total of 545 batches from further 201 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 72 spectra from 21 *Apo-Ident* customers from 37 batches from the substance/substance group *Ointment ash basis / Neribas®*.
- These include spectra of independent samples from 37 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Chiese	Asche Basis [®] ointment	31045A	4
Chiesi/AHD	Asche Basis [®] ointment	32048A	2
Chiesi / Gehe	Asche Basis [®] ointment	33050A	1
Chiesi	Asche Basis [®] ointment	41052A	1
Chiesi/Noweda	Asche Basis [®] ointment	41053A	1
Chiesi/Sanacorp	Asche Basis [®] ointment	41053A	1
Chiesi	Asche Basis [®] ointment	41054A	3
Chiesi/Gehe	Asche Basis [®] ointment	42056A	1
chiesi	Asche Basis [®] ointment	42055A	1
chiesi	Asche Basis [®] ointment	44057A	1
Chiesi/Phönix	Asche Basis [®] ointment	51059A	1
Chiesi	Asche Basis [®] ointment	51060A	1
CHIESI/ NOWEDA	Asche Basis [®] ointment	51060A	2
Chiesi	Asche Basis [®] ointment	52062A	2
Chiesi	Asche Basis [®] ointment	52063A	3
Chiesi GmbH	Asche Basis [®] ointment	52062A	1
Chiesi GmbH	Asche Basis [®] ointment	53065A	4
Chiesi GmbH	Asche Basis [®] ointment	61066A	2
Caelo	Asche Basis [®] ointment	61067A	1
Caelo	Asche Basis [®] ointment	63068A	2
Chiesi	Asche Basis [®] ointment	63068A	1
Chiesi	Asche Basis [®] ointment	63069A	3
Chiesi GmbH	Asche Basis [®] ointment	63070A	2
Chiesi GmbH	Asche Basis [®] ointment	71071A	1
Chiesi GmbH	Asche Basis [®] ointment	71073A	1
Chiesi GmbH	Asche Basis [®] ointment	72074A	1
Chiesi GmbH	Asche Basis [®] ointment	21039A	1
Chiesi/Phoenix	Asche Basis [®] ointment	23042A	1
Anzag/Ichtyol-Gesellsch. ...	Asche Basis [®] ointment	23043A	1
Phönix	Asche Basis [®] ointment	31047A	1
Chiesi/AHD	Asche Basis [®] ointment	31047A	1
Caelo	Asche Basis [®] ointment	31047A	1
Chiesi	Asche Basis [®] ointment	32048A	1
Chiesi/AHD	Asche Basis [®] ointment	33049A	1
Chiesi / Gehe	Asche Basis [®] ointment	33049A	1
Chiesi	Asche Basis [®] ointment	33049A	1
Chiesi	Asche Basis [®] ointment	34051A	1
Jenapharm/ Sanacorp	Neribas [®] ointment	YY006JT	3
Jenapharm	Neribas [®] ointment	YY006JT	1
Caelo	Neribas [®] ointment	YY00CC8	1
Jenapharm/ Anzag	Neribas [®] ointment	YY00K7E	2
Jenapharm / Anzag	Neribas [®] ointment	YY00THL	1
Jenapharm/ Sanacorp	Neribas [®] ointment	YY00THL	1
Jenapharm/Noweda	Neribas [®] ointment	YY00THL	1
Caelo	Neribas [®] ointment	YY00TJ0	1
Jenapharm/Anzag	Neribas [®] ointment	YY00XEH	2
Caelo	Neribas [®] ointment	YY00XEH	1
Jenapharm	Neribas [®] ointment	YY00XEH	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Caelo	Neribas [®] ointment	YY016LN	1
Jenapharm	Neribas [®] ointment	YY017LF	1

- 3121 spectra from 304 *Apo-Ident* customers from a total of 1622 batches from a further 120 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Ointment ash basis / Neribas[®]* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Ointment ash basis / Neribas[®]* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	240	0	18 509
Type B	0	239	1	22 551
Type C	0	67	5	3121

The substance/substance group *Ointment ash basis / Neribas[®]* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.5000 %)
Type B	100.0000 % (> 99.9536 %)	99.5833 % (> 98.3333 %)
Type C	100.0000 % (> 98.3578 %)	93.0556 % (> 88.8889 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30774	30774	0.00	36.35
30833	30833	0.00	37.75

continued on the next page

continued from previous page

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30969	30969	0.00	36.81
30994	30994	0.00	34.20
31080	31080	0.00	36.24
31093	31093	0.00	27.84

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Orange aroma**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31190-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Orange aroma; Orange flavour

Special notes

When selecting the *Orange aroma* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Orange aroma*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Orange aroma	13123307	31190	40	not required
Caelo	Orange aroma	13406108	31402	40	not required
Fagron	Orange aroma	13H23-B41-295530	31510	40	not required
Caelo	Orange aroma	163470	33100	40	20161221*

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Orange aroma*. These samples are listed above in the section *calibration samples*. The reference samples originate from 4 different batches.
- 18 589 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this *chemometric model*, the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in *Appendix A*. The samples in *Appendix A* were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 220 spectra from 6 reference samples of the substance/substance group *Orange aroma*.
- These include spectra of independent samples from 5 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	Orange aroma	11069914	30335	40
Symrise	Orange aroma	113	30414	40
Fagron	Orange aroma	13H23-B41-295530	31510 [†]	20
Caelo	Orange aroma	162063	32331	40
Caelo	Orange aroma	173005	33881	40
Caelo	Orange aroma	180831	34076	40

*The spectra were taken from a part of the same sample which the supplier used for its analysis (identity, purity, content). In this case the manufacturer certificates proof the identity and quality of the sample from those the spectra descend.

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

- 22 611 spectra from a total of 543 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 59 spectra from 28 *Apo-Ident* customers from 34 batches from the substance/substance group *Orange aroma*.
- These include spectra of independent samples from 33 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Orange aroma	13330207	1
Caelo	Orange aroma	13406101	2
Caelo	Orange aroma	13406104	1
Caelo	Orange aroma	13406108	3
Caelo	Orange aroma	14150703	1
Caelo	Orange aroma	14150706	6
Caelo	Orange aroma	14150707	3
Caelo	Orange aroma	14150714	1
Caelo	Orange aroma	14352904	2
Caelo	Orange aroma	14352903	1
Fagron	Orange aroma	14352904	1
Caelo	Orange aroma	14352906	1
Caelo	Orange aroma	15215401	1
Caelo	Orange aroma		1
Caelo	Orange aroma	15215402	3
Caelo	Orange aroma	15215403	2
Caelo	Orange aroma	15215408	2
Audor Pharma	Orange aroma	15215409	1
Caelo	Orange aroma	15215410	1
Caelo	Orange aroma	16056105	3
Allergika/Kehr	Orange aroma	16056107	1
Caelo	Orange aroma	16056107	1
Caelo	Orange aroma	16056108	1
Caelo	Orange aroma	16347005	1
Caelo	Orange aroma	17055104	1
Caelo	Orange aroma	17300501	1
Fagron	Orange aroma	17A30-B02-335197	1
Caelo	Orange aroma	12149707	1
Caelo	Orange aroma	12216712	2
Caelo	Orange aroma	12356502	2
Sanacorp	Orange aroma	12356503	1
Caelo	Orange aroma	12356503	2
Caelo	Orange aroma	13123305	2
Caelo	Orange aroma	13123309	2
Caelo	Orange aroma	13330206	1
Caelo	Orange aroma	23051201	1
Caelo	Orange aroma	25031304	1

- 3134 spectra from 302 *Apo-Ident* customers from a total of 1625 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Orange aroma* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Orange aroma* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	211	9	22 611
Type C	0	58	1	3134

The substance/substance group *Orange aroma* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9529 %)	95.9091 % (> 94.5455 %)
Type C	100.0000 % (> 98.3581 %)	98.3051 % (> 93.2203 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31190	31190	0.00	93.24
31402	31402	0.00	90.09
31510	31510	0.00	136.83
33100	33100	0.00	93.52

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Oregano oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31554-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Oregano oil; Oleum origani cretici

Special notes

When selecting the *Oregano oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Oregano oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Oregano oil	0131022-117999	31554	40	not required
Taoasis	Oregano oil	7921018-119483	31779	40	not required
Taoasis	Oregano oil	43878A-125814	32769	40	not required
Taoasis	Oregano oil	44314A-129096	33649	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Oregano oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 589 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 80 spectra from 3 reference samples of the substance/substance group *Oregano oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Oregano oil	0131022-117999	31554 [†]	20
Taoasis	Oregano oil	7921018-119483	31779 [†]	20
Taoasis	Oregano oil	43878A-123201	32939	40

- 22 751 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Oregano oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Oregano oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Oregano oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	80	0	22 751
Type C	0	0	0	3193

The substance/substance group *Oregano oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9531 %)	100.0000 % (> 92.5000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31554	31554	0.00	100.68
31779	31779	0.00	143.83
32769	32769	0.00	48.51
33649	33649	0.00	51.32

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Palmarosa oil, organic**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30597-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Palmarosa oil, organic; Oleum cymbopogon martinii; Oleum palmarosae

Special notes

When selecting the *Palmarosa oil, organic* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Palmarosa oil, organic*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Palmarosa oil, o...	2461021-121891	31828	40	not required
Taoasis	Palmarosa oil, o...	4941003-126708	33043	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Palmarosa oil, organic*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 180 spectra from 5 reference samples of the substance/substance group *Palmarosa oil, organic*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Palmarosa oil, organic	17524-105015BAG90451	30597	40
Taoasis	Palmarosa oil, organic	17524-105015BAG90451	30671	40
Taoasis	Palmarosa oil, organic	2461021-121891	31828 [†]	20
Taoasis	Palmarosa oil, organic	4941003-126708	33044	40
Taoasis	Palmarosa oil, organic	104704-129227	33856	40

- 22 651 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 4 spectra from 3 *Apo-Ident* customers from 2 batches from the substance/substance group *Palmarosa oil, organic*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis/Taoasis	Palmarosa oil, organic	121018-111872	1
Taoasis	Palmarosa oil, organic	121018-111094	1
Taoasis	Palmarosa oil, organic	121018-111872	2

- 3189 spectra from 304 *Apo-Ident* customers from a total of 1657 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Palmarosa oil, organic* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Palmarosa oil, organic* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	169	11	22 651
Type C	0	4	0	3189

The substance/substance group *Palmarosa oil, organic* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9529 %)	93.8889 % (> 92.2222 %)
Type C	100.0000 % (> 98.3777 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31828	31828	0.00	168.29
33043	33043	0.00	141.06

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Panthenol ointment Lichtenstein**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30727-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Panthenol ointment Lichtenstein

Special notes

When selecting the *Panthenol ointment Lichtenstein* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Panthenol ointment Lichtenstein*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Winthrop	Panthenol ointme...	2290680H	30727	40	not required
Winthrop	Panthenol ointme...	2290718H	31091	40	not required
Zentiva Pharm...	Panthenol ointme...	EH4594	31764	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Panthenol ointment Lichtenstein*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Panthenol ointment Lichtenstein*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Zentiva Pharma GmbH	Panthenol ointment Lichtenst. ...	EH4594	31764 [†]	20
Zentiva Pharma GmbH	Panthenol ointment Lichtenst. ...	GC5567	33284	40
Zentiva Pharma GmbH	Panthenol ointment Lichtenst. ...	GD5661	33425	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Panthenol ointment Lichtenstein*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Panthenol ointment Lichtenstein* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Panthenol ointment Lichtenstein* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	100	0	22 731
Type C	0	0	0	3193

The substance/substance group *Panthenol ointment Lichtenstein* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30727	30727	0.00	205.50
31091	31091	0.00	201.57
31764	31764	0.00	168.34

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Patchouli oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31547-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Patchouli oil; Oleum pogostemon cablin

Special notes

When selecting the *Patchouli oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Patchouli oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Patchouli oil	130930-112789	31547	40	not required
Caelo	Patchouli oil	14167204	31619	40	not required
Taoasis	Patchouli oil	FFL735-123003	32172	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Patchouli oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *Patchouli oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Patchouli oil	130930-112789	31547 [†]	20
Caelo	Patchouli oil	14167204	31619 [†]	20
Taoasis	Patchouli oil	1633-126207	33031	40
Taoasis	Patchouli oil	1633-126207	33032	40

- 22 711 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 2 spectra from 1 *Apo-Ident* customers from 2 batches from the substance/substance group *Patchouli oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Patchouli oil	130930-116684	1
Taoasis	Patchouli oil	33474-119161	1

- 3191 spectra from 304 *Apo-Ident* customers from a total of 1657 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Patchouli oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Patchouli oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	40	80	22 711
Type C	0	1	1	3191

The substance/substance group *Patchouli oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	33.3333 % (> 30.8333 %)
Type C	100.0000 % (> 98.3987 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31547	31547	0.00	26.60
31619	31619	0.00	33.41
32172	32172	0.00	41.16

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Peppermint oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30930-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Peppermint oil; Menthae piperitae aetheroleum; Oleum menthae piperitae dopp. rectific.

Special notes

When selecting the *Peppermint oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Peppermint oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Peppermint oil	443428-83579	30930	40	not required
Taoasis	Peppermint oil	525728-109217B	30965	40	not required
Taoasis	Peppermint oil	2016104277-125426	32481	40	not required
Primavera	Peppermint oil	00098J25	33809	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Peppermint oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 589 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 3 reference samples of the substance/substance group *Peppermint oil*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Peppermint oil	2016104277-126375	32804	40
Taoasis	Peppermint oil	104277-126684	33030	40
Primavera	Peppermint oil	00374B26	33958	40

- 22 711 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 12 spectra from 6 *Apo-Ident* customers from 12 batches from the substance/substance group *Peppermint oil*.
- These include spectra of independent samples from 12 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Peppermint oil	12114225	1
Taoasis/Kehr	Peppermint oil	127711	1
Fagron	Peppermint oil	15B11-B08-304117	1
Taoasis	Peppermint oil	24699-117643	1
Bombastus	Peppermint oil	286204	1
Taoasis	Peppermint oil	40076-120885	1
Taoasis/ Kehr	Peppermint oil	41798-120746H08	1
Taoasis	Peppermint oil	41798-121607	1
Taoasis	Peppermint oil	52216-123564	1
Taoasis	Peppermint oil	63561-125104	1
Taoasis	Peppermint oil	63561-125729	1
Taoasis	Peppermint oil	63561-126008	1

- 3181 spectra from 303 *Apo-Ident* customers from a total of 1647 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Peppermint oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Peppermint oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	116	4	22 711
Type C	0	12	0	3181

The substance/substance group *Peppermint oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9530 %)	96.6667 % (> 94.1667 %)
Type C	100.0000 % (> 98.3637 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30930	30930	0.00	91.87
30965	30965	0.00	99.88
32481	32481	0.00	79.23
33809	33809	0.00	75.52

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Perfume oil vanilla**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30665-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Perfume oil vanilla; Oleum vanillae planifoliae

Special notes

When selecting the *Perfume oil vanilla* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Perfume oil vanilla*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Perfume oil vani...	10293212	30665	40	not required
Caelo	Perfume oil vani...	13393501	31191	40	1403100
Caelo	Perfume oil vani...	13393501	31400	40	not required
Caelo	Perfume oil vani...	14284107	31786	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Perfume oil vanilla*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 589 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 112 spectra from 4 reference samples of the substance/substance group *Perfume oil vanilla*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	Perfume oil vanilla	14284107	31786 [†]	20
Caelo	Perfume oil vanilla	15316308	33439	40
Caelo	Perfume oil vanilla	15316308	33440	40
Caelo	Perfume oil vanilla	15316308	not required	12

- 22 719 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Perfume oil vanilla*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Perfume oil vanilla* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Perfume oil vanilla* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	112	0	22 719
Type C	0	0	0	3193

The substance/substance group *Perfume oil vanilla* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.6429 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30665	30665	0.00	143.73
31191	31191	0.00	137.53
31400	31400	0.00	138.02
31786	31786	0.00	150.39

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Raspberry aroma**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30562-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Raspberry aroma; Raspberry flavour

Special notes

When selecting the *Raspberry aroma* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Raspberry aroma*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Raspberry aroma	12246612	31171	40	1402588
Caelo	Raspberry aroma	14115902	31396	40	not required
Caelo	Raspberry aroma	16126605	33222	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Raspberry aroma*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 160 spectra from 4 reference samples of the substance/substance group *Raspberry aroma*.
- These include spectra of independent samples from 4 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	Raspberry aroma	11303813	30562	40
Caelo	Raspberry aroma	12054307	30611	40
Caelo	Raspberry aroma	15438503	33054	40
Caelo	Raspberry aroma	172581	33697	40

- 22 671 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 64 spectra from 40 *Apo-Ident* customers from 32 batches from the substance/substance group *Raspberry aroma*.
- These include spectra of independent samples from 29 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caesar/Noweda	Raspberry aroma	212m-07020	1
Caelo	Raspberry aroma	13255507	2
Caelo	Raspberry aroma	13255514	3
Caelo	Raspberry aroma	1355514	1
Caelo	Raspberry aroma	14115902	2
Noweda	Raspberry aroma	14115909	1
Caelo	Raspberry aroma	14115912	1
Caelo	Raspberry aroma	14115913	2
Caelo	Raspberry aroma	14376103	2
Caelo	Raspberry aroma	14376109	3
Caelo	Raspberry aroma	14376115	4
Caelo	Raspberry aroma	15219102	3
Caelo	Raspberry aroma	14378103	1
Caelo	Raspberry aroma	15438501	5
Rondell Apotheke	Raspberry aroma	15438501	1
Caelo	Raspberry aroma	15438503	3
Phönix	Raspberry aroma	15438506	1
Caelo	Raspberry aroma	16126605	2
Caelo	Raspberry aroma	16126609	2
Caelo	Raspberry aroma	17187212	4
Caelo	Raspberry aroma	17258103	1
Caelo	Raspberry aroma	11303813	1
Gehe 16.04.2013	Raspberry aroma	12054307	1
Caelo	Raspberry aroma	12054302	1
Caelo	Raspberry aroma	12054310	2
Caelo	Raspberry aroma	12246604	1
Caelo	Raspberry aroma	12246608	2
Phönix Gotha	Raspberry aroma	12246612	1
Caelo	Raspberry aroma	12246612	2
Caelo	Raspberry aroma	12246616	2
Caelo	Raspberry aroma	12246617	1
Caelo	Raspberry aroma	1254307	1
Caelo	Raspberry aroma	13255502	2
Caelo	Raspberry aroma	14115909	1
Caelo	Raspberry aroma	12054311	1

- 3129 spectra from 298 *Apo-Ident* customers from a total of 1627 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Raspberry aroma* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Raspberry aroma* and it was evaluated

how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	153	7	22 671
Type C	0	62	2	3129

The substance/substance group *Raspberry aroma* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9529 %)	95.6250 % (> 93.7500 %)
Type C	100.0000 % (> 98.3580 %)	96.8750 % (> 92.1875 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31171	31171	0.00	80.43
31396	31396	0.00	76.22
33222	33222	0.00	81.45

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Ravensara**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30386-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Ravensara

Special notes

When selecting the *Ravensara* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Ravensara*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Ravensara	231012-118739	31635	40	not required
Taoasis	Ravensara	1687-125316	32803	40	not required
Taoasis	Ravensara	1687-966	33854	40	not required
Taoasis	Ravensara	1687-966	33872	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Ravensara*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 589 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Ravensara*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Ravensara	T170I11	30386	40
Taoasis	Ravensara	231012-118739	31635 [†]	20
Taoasis	Ravensara	1687-125316	33029	40

- 22 731 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 13 spectra from 1 *Apo-Ident* customers from 8 batches from the substance/substance group *Ravensara*.
- These include spectra of independent samples from 6 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Ravensara	231012-119831	1
Taoasis	Ravensara	1687-125316	1
Taoasis	Ravensara	2101005-116830	3
Taoasis	Ravensara	231012-118738	1
Taoasis	Ravensara	231012-118739	2
Taoasis	Ravensara	231012-118892	1
Taoasis	Ravensara	231012-120291	1
Taoasis	Ravensara	9331006-121650	3

- 3180 spectra from 304 *Apo-Ident* customers from a total of 1651 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Ravensara* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Ravensara* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	100	0	22 731
Type C	0	13	0	3180

The substance/substance group *Ravensara* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.0000 %)
Type C	100.0000 % (> 98.3631 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to

exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31635	31635	0.00	23.48
32803	32803	0.00	26.75
33854	33854	0.00	26.51
33872	33872	0.00	25.35

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Rose-geranium oil, organic**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30536-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Rose-geranium oil, organic; Oleum pelargonium graveolens

Special notes

When selecting the *Rose-geranium oil, organic* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Rose-geranium oil, organic*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Rose-geranium oi...	21027-119997	31698	40	not required
Taoasis	Rose-geranium oi...	21027-121893	31776	40	not required
Taoasis	Rose-geranium oi...	21027-121893	31951	40	not required
Taoasis	Rose-geranium oi...	1616008-125056	33036	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Rose-geranium oil, organic*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 589 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 180 spectra from 6 reference samples of the substance/substance group *Rose-geranium oil, organic*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Rose-geranium oil, organic	zu erfassen	30536	40
Taoasis	Rose-geranium oil, organic	21027-119997	31698 [†]	20
Taoasis	Rose-geranium oil, organic	21027-121893	31776 [†]	20
Taoasis	Rose-geranium oil, organic	21027-121893	31951 [†]	20
Taoasis	Rose-geranium oil, organic	1616008-125056	33037	40
Taoasis	Rose-geranium oil, organic	2260-2430	34042	40

- 22 651 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 13 spectra from 7 *Apo-Ident* customers from 12 batches from the substance/substance group *Rose-geranium oil, organic*.
- These include spectra of independent samples from 12 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Bombastus	Rose-geranium oil, organic	22625-106677	1
Taoasis	Rose-geranium oil, organic	21027-119121	1
Taoasis	Rose-geranium oil, organic	21027-120292	1
Taoasis	Rose-geranium oil, organic	401011-117806	2
Taoasis	Rose-geranium oil, organic	1500969-123554	1
Taoasis	Rose-geranium oil, organic	21027-121337	1
Taoasis/Taoasis	Rose-geranium oil, organic	22625-104463	1
Taoasis	Rose-geranium oil, organic	401011-117641	1
Primavera	Rose-geranium oil, organic	8304131	1
Taoasis	Rose-geranium oil, organic	130111-110467	1
Taoasis	Rose-geranium oil, organic	130111-112615	1
Taoasis	Rose-geranium oil, organic	130729-113060	1

- 3180 spectra from 304 *Apo-Ident* customers from a total of 1647 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Rose-geranium oil, organic* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Rose-geranium oil, organic* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	180	0	22 651
Type C	0	13	0	3180

The substance/substance group *Rose-geranium oil, organic* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9529 %)	100.0000 % (> 96.6667 %)
Type C	100.0000 % (> 98.3631 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31698	31698	0.00	30.25
31776	31776	0.00	34.74
31951	31951	0.00	35.25
33036	33036	0.00	37.38

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Rosewood oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31556-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Rosewood oil; Oleum aniba roseodora

Special notes

When selecting the *Rosewood oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Rosewood oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Rosewood oil	11572-116545	31556	40	not required
Taoasis	Rosewood oil	4381004-120355	31688	40	not required
Taoasis	Rosewood oil	42543-2657	34106	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Rosewood oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *Rosewood oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Rosewood oil	11572-116545	31556 [†]	20
Taoasis	Rosewood oil	4381004-120355	31688 [†]	20
Taoasis	Rosewood oil	5101015-124967	32801	40
Taoasis	Rosewood oil	5101015-124967	33048	40

- 22 711 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 11 spectra from 1 *Apo-Ident* customers from 7 batches from the substance/substance group *Rosewood oil*.
- These include spectra of independent samples from 7 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Rosewood oil	4381004-1200355	1
Taoasis	Rosewood oil	4381004-121143	2
Taoasis	Rosewood oil	4381004-122041	2
Taoasis	Rosewood oil	5101017-122566	2
Taoasis	Rosewood oil	5101015-123299	1
Taoasis	Rosewood oil	5101015-124243	1
Taoasis	Rosewood oil	5101015-124967	2

- 3182 spectra from 304 *Apo-Ident* customers from a total of 1652 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Rosewood oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Rosewood oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	120	0	22 711
Type C	0	11	0	3182

The substance/substance group *Rosewood oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 95.0000 %)
Type C	100.0000 % (> 98.3643 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to

exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31556	31556	0.00	52.79
31688	31688	0.00	58.97
34106	34106	0.00	47.20

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Sage oil
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	31139-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Sage oil; Oleum salviae; Salvia officinalis oleum

Special notes

When selecting the *Sage oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Sage oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Sage oil	526975-112642BAG90451	31139	40	not required
Taoasis	Sage oil	34052-114939	31257	40	not required
Taoasis	Sage oil	8461016-122272	31783	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Sage oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 4 reference samples of the substance/substance group *Sage oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Sage oil	8461016-122272	31783 [†]	20
Taoasis	Sage oil	8461016-126496	33034	40
Taoasis	Sage oil	8461016-126496	33035	40
Taoasis	Sage oil	24858-79297	36005	40

- 22 691 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 5 spectra from 1 *Apo-Ident* customers from 5 batches from the substance/substance group *Sage oil*.
- These include spectra of independent samples from 5 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Sage oil	13988-117449	1
Taoasis	Sage oil	21190-118009	1
Taoasis	Sage oil	21190-119239	1
Taoasis	Sage oil	29944-120889	1
Taoasis	Sage oil	29944-120965	1

- 3188 spectra from 304 *Apo-Ident* customers from a total of 1654 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Sage oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Sage oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	105	35	22 691
Type C	0	5	0	3188

The substance/substance group *Sage oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	75.0000 % (> 72.8571 %)
Type C	100.0000 % (> 98.3735 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31139	31139	0.00	35.75
31257	31257	0.00	41.92
31783	31783	0.00	36.32

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Salicylic white paraffin 10%**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31521-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Salicylic white paraffin 10%; Acidum salicylicum cum vaselino albo 10 %

Special notes

When selecting the *Salicylic white paraffin 10%* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Salicylic white paraffin 10%*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Bombastus	Salicylic white ...	287820	31521	40	not required
Bombastus	Salicylic white ...	291886	31629	55	not required
Bombastus	Salicylic white ...	310227	33473	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 135 spectra of 3 reference samples from the substance/substance group *Salicylic white paraffin 10%*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 614 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *Salicylic white paraffin 10%*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Bombastus	Salicylic white paraffin 10%	287820	31521 [†]	20
Bombastus	Salicylic white paraffin 10%	291886	31629 [†]	20
Bombastus	Salicylic white paraffin 10%	302737	32741	40
Bombastus	Salicylic white paraffin 10%	309004	33416	40

- 22 711 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 7 spectra from 5 *Apo-Ident* customers from 6 batches from the substance/substance group *Salicylic white paraffin 10%*.
- These include spectra of independent samples from 6 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Eigenrezeptur	Salicylic white paraffin 10%	150702	1
Caelo	Salicylic white paraffin 10%	16120701	2
Caelo	Salicylic white paraffin 10%	15394702	1
Caelo	Salicylic white paraffin 10%	16214202	1
Caelo	Salicylic white paraffin 10%	16214203	1
Bombastus	Salicylic white paraffin 10%	293212	1

- 3186 spectra from 303 *Apo-Ident* customers from a total of 1653 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Salicylic white paraffin 10%* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Salicylic white paraffin 10%* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	135	0	18 614
Type B	0	120	0	22 711
Type C	0	6	1	3186

The substance/substance group *Salicylic white paraffin 10%* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9535 %)	100.0000 % (> 95.5556 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 95.0000 %)
Type C	100.0000 % (> 98.3687 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31521	31521	0.00	80.54
31629	31629	0.00	80.22
33473	33473	0.00	82.51

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Sandalwood oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 33799-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Sandalwood oil

Special notes

When selecting the *Sandalwood oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Sandalwood oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Primavera	Sandalwood oil	00891G25	33799	40	not required
Primavera	Sandalwood oil	00141L25	33801	40	not required
Taoasis	Sandalwood oil	145-505	33832	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Sandalwood oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 40 spectra from 1 reference samples of the substance/substance group *Sandalwood oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Sandalwood oil	128319	33833	40

- 22 791 spectra from a total of 548 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Sandalwood oil*.

- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Sandalwood oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Sandalwood oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	40	0	22 791
Type C	0	0	0	3193

The substance/substance group *Sandalwood oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9535 %)	100.0000 % (> 85.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
33799	33799	0.00	147.49
33801	33801	0.00	151.30
33832	33832	0.00	133.03

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Sebexol® formula basis / Sebexol® cream lotion / ...**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30839-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Sebexol® formula basis / Sebexol® cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver; Hans Karrer Hydrocream MicroSilver; Sebexol® basic pH 5 formula basis; Sebexol® cream lotion pH 5; Wofacutan wash lotion

Special notes

When selecting the *Sebexol® formula basis / Sebexol® cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Sebexol® formula basis / Sebexol® cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Devesa	Sebexol® cream l...	1220	30839	40	1403097

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Kesla Pharma	Wofacutan wash l...	0974.12	30972	80	not required
Kesla Pharma	Wofacutan wash l...	0389.13	31065	40	not required
Devesa	Sebexol [®] cream l...	1317	31103	40	not required
Devesa	Sebexol [®] basic p...	1307	31104	40	1403098
Devesa	Sebexol [®] cream l...	1412	31592	30	not required
Devesa	Sebexol [®] cream l...	1531	31839	40	not required
Devesa	Sebexol [®] basic p...	1522	31961	40	not required
Devesa	Sebexol [®] cream l...	1552.2	32743	40	not required
Devesa	Sebexol [®] basic p...	1638124	32744	40	not required
Hans Karrer G...	Hans Karrer Hydr...	16001	33190	40	not required
Hans Karrer G...	Hans Karrer Hydr...	16002	33191	40	not required
Kesla Pharma	Wofacutan wash l...	0868.16	33483	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 550 spectra of 13 reference samples from the substance/substance group *Sebexol[®] formula basis / Sebexol[®] cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 13 different batches.
- 18 199 spectra from a total of 426 batches from further 147 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 415 spectra from 12 reference samples of the substance/substance group *Sebexol[®] formula basis / Sebexol[®] cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver*.
- These include spectra of independent samples from 7 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Devesa	Sebexol [®] basic pH 5 formula ...	1131	30521	40
Devesa	Sebexol [®] cream lotion pH 5	1412	31592 [†]	15
Devesa	Sebexol [®] cream lotion pH 5	1531	31839 [†]	20
Devesa	Sebexol [®] basic pH 5 formula ...	1522	31961 [†]	20
Devesa	Sebexol [®] cream lotion pH 5	1605101	32907	40
Devesa	Sebexol [®] cream lotion pH 5	1605101	32908	40
Kesla Pharma	Wofacutan wash lotion	0729.16	32913	40
Kesla Pharma	Wofacutan wash lotion	0729.16	32914	40
Hans Karrer GmbH	Hans Karrer Hydrocream Micro...	16003	33192	40
Hans Karrer GmbH	Hans Karrer Hydrocream Micro...	16004	33193	40
Devesa	Sebexol [®] basic pH 5 formula ...	1712122	33427	40
Devesa	Sebexol [®] basic pH 5 formula ...	1644102	33432	40

- 22 416 spectra from a total of 539 batches from further 199 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 196 spectra from 43 *Apo-Ident* customers from 72 batches from the substance/substance group *Sebexol[®] formula basis* / *Sebexol[®] cream lotion* / *Wofacutan washing lotion* / *Hans Karrer Hydrocream MicroSilver*.
- These include spectra of independent samples from 62 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Devesa	Sebexol [®] basic pH 5 formula ...	1307	5
Devesa	Sebexol [®] basic pH 5 formula ...	1440	4
DEVESA	Sebexol [®] basic pH 5 formula ...	1440	1
Devesa	Sebexol [®] basic pH 5 formula ...	1511	2
DEVESA	Sebexol [®] basic pH 5 formula ...	1511	1
Fiebig	Sebexol [®] basic pH 5 formula ...	1511	3
Devesa/Noweda	Sebexol [®] basic pH 5 formula ...	1511	1
Devesa	Sebexol [®] basic pH 5 formula ...	1522	3
Devesa/Sanacorp	Sebexol [®] basic pH 5 formula ...	1522	1
Devesa/Sanacorp	Sebexol [®] basic pH 5 formula ...	1543	4
Devesa	Sebexol [®] basic pH 5 formula ...	1543	5
Devesa/Phönix	Sebexol [®] basic pH 5 formula ...	1543	1
Devesa	Sebexol [®] basic pH 5 formula ...	1604127	4
Caelo	Sebexol [®] basic pH 5 formula ...	1604127	2
Devesa/Noweda	Sebexol [®] basic pH 5 formula ...	1604127	1
Caelo	Sebexol [®] basic pH 5 formula ...	1620117	2
Devesa/Sanacorp	Sebexol [®] basic pH 5 formula ...	1620117	1
Devesa/Phönix	Sebexol [®] basic pH 5 formula ...	1620117	1
Devesa	Sebexol [®] basic pH 5 formula ...	1620117	1

continued on the next page

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

continued from previous page

Supplier	Substance	Batch	Spectra
Devesa/Sanacorp	Sebexol [®] basic pH 5 formula ...	1638124	1
Devesa	Sebexol [®] basic pH 5 formula ...	1638124	2
Devesa/Sanacorp	Sebexol [®] basic pH 5 formula ...	1644102	2
Devesa / Noweda	Sebexol [®] basic pH 5 formula ...	1620117	1
Devesa	Sebexol [®] basic pH 5 formula ...	1644102	3
Devesa/Sanacorp	Sebexol [®] basic pH 5 formula ...	1712122	1
Devesa	Sebexol [®] basic pH 5 formula ...	1712122	2
Phönix / Devesa	Sebexol [®] basic pH 5 formula ...	1644102	1
Devesa	Sebexol [®] basic pH 5 formula ...	1683124	1
Devesa/Sanacorp	Sebexol [®] basic pH 5 formula ...	1719111	1
Devesa	Sebexol [®] basic pH 5 formula ...	1719111	2
Caelo	Sebexol [®] cream lotion pH 5	1417	1
Devesa/Fiebig	Sebexol [®] cream lotion pH 5	1426	1
Devesa/Noweda	Sebexol [®] cream lotion pH 5	1447	3
Astellas	Sebexol [®] cream lotion pH 5	1504	2
Caelo	Sebexol [®] cream lotion pH 5	1504	1
Devesa/NOW	Sebexol [®] cream lotion pH 5	1504	2
Devesa/Gehe	Sebexol [®] cream lotion pH 5	1504	1
Devesa/Noweda	Sebexol [®] cream lotion pH 5	1504	1
Devesa/NOW	Sebexol [®] cream lotion pH 5	1504/01	1
Hedinger/Phoenix	Sebexol [®] cream lotion pH 5	1510	2
DEVESA	Sebexol [®] cream lotion pH 5	1505	1
Devesa	Sebexol [®] cream lotion pH 5	1510	2
DEVESA	Sebexol [®] cream lotion pH 5	1510	1
Devesa/Phoenix	Sebexol [®] cream lotion pH 5	1511	1
Caelo	Sebexol [®] cream lotion pH 5	1511	2
Fagron	Sebexol [®] cream lotion pH 5	1517	1
DEVESA	Sebexol [®] cream lotion pH 5	1523	1
Devesa/NOW	Sebexol [®] cream lotion pH 5	1523	1
Devesa	Sebexol [®] cream lotion pH 5	1523	3
Rondell Apotheke	Sebexol [®] cream lotion pH 5	1523	1
Devesa/ Phönix	Sebexol [®] cream lotion pH 5	1524	1
Devesa/Sanacorp	Sebexol [®] cream lotion pH 5	1530	1
Caelo	Sebexol [®] cream lotion pH 5	1531	1
Devesa / Noweda	Sebexol [®] cream lotion pH 5	1531	1
DEVESA	Sebexol [®] cream lotion pH 5	1531	1
Dr. Wolff	Sebexol [®] cream lotion pH 5		1
Caelo	Sebexol [®] cream lotion pH 5		2
Devesa/Noweda	Sebexol [®] cream lotion pH 5		1
Devesa/fiebig	Sebexol [®] cream lotion pH 5		1
Devesa/NOW	Sebexol [®] cream lotion pH 5		1
Devesa/NOW	Sebexol [®] cream lotion pH 5	1552	1
Devesea	Sebexol [®] cream lotion pH 5	1604128	1
Phönix	Sebexol [®] cream lotion pH 5	1604129	1
Devesa/NOW	Sebexol [®] cream lotion pH 5	1604129	1
Devesa/NOW	Sebexol [®] cream lotion pH 5	1604228	1
sanacorp, 27.9.16,4,312EUR	Sebexol [®] cream lotion pH 5	1604328	1
Devesa/Noweda	Sebexol [®] cream lotion pH 5	1605201	3
Devesa/NOW	Sebexol [®] cream lotion pH 5	1644102	1
Devesa/Noweda	Sebexol [®] cream lotion pH 5	1651120	1
Devesa	Sebexol [®] cream lotion pH 5	1651121	1
Noweda	Sebexol [®] cream lotion pH 5	1651120	1
Devesa	Sebexol [®] cream lotion pH 5	1708120	2
Devesa/Noweda	Sebexol [®] cream lotion pH 5	1712123	1
Caelo	Sebexol [®] cream lotion pH 5	1712223	1
Devesa/Noweda	Sebexol [®] cream lotion pH 5	1736105	1
Devesa	Sebexol [®] cream lotion pH 5	1727107	1
Devesa/Noweda	Sebexol [®] cream lotion pH 5	1736106	1
Devesa/Sanacorp	Sebexol [®] cream lotion pH 5	1743125	1
Devesa/Noweda	Sebexol [®] cream lotion pH 5	1743225	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Caelo	Sebexol [®] basic pH 5 formula ...	1328	4
Devesa	Sebexol [®] basic pH 5 formula ...	1328	8
DEVESA	Sebexol [®] basic pH 5 formula ...	1328	2
DEVESA Dr.Reingraber GmbH	Sebexol [®] basic pH 5 formula ...	1328	1
Devesa	Sebexol [®] basic pH 5 formula ...	1335	2
Devesa/ Fiebig	Sebexol [®] basic pH 5 formula ...	1338	1
devesa/gehe	Sebexol [®] basic pH 5 formula ...	1346	1
Bombastus	Sebexol [®] basic pH 5 formula ...	1346	2
devesa/gehe	Sebexol [®] basic pH 5 formula ...	1356	1
Sebexol/ Noweda	Sebexol [®] basic pH 5 formula ...	1403	1
Devesa	Sebexol [®] basic pH 5 formula ...	1404	1
Phoenix/Devesa	Sebexol [®] basic pH 5 formula ...	1407	1
Devesa	Sebexol [®] basic pH 5 formula ...	1407	2
Devesa/Kehr	Sebexol [®] basic pH 5 formula ...	1407	2
Phönix	Sebexol [®] basic pH 5 formula ...	1416	1
Devesa Dr.Reingraber GmbH	Sebexol [®] basic pH 5 formula ...	1425	1
Phönix	Sebexol [®] basic pH 5 formula ...	1425	2
Fagron	Sebexol [®] basic pH 5 formula ...	1425	1
Devesa/Noweda	Sebexol [®] basic pH 5 formula ...	1430	1
Sebexol/ Noweda	Sebexol [®] basic pH 5 formula ...	1430	1
Phoenix	Sebexol [®] basic pH 5 formula ...	1440	1
Devesa	Sebexol [®] basic pH 5 formula ...	1430	1
Devesa/Sanacorp	Sebexol [®] basic pH 5 formula ...	1440	1
Fagron	Sebexol [®] basic pH 5 formula ...	1510	2
Noweda/Devesa	Sebexol [®] basic pH 5 formula ...	2x1407	1
DEVESA	Sebexol [®] basic pH 5 formula ...	1318	2
Eigenherstellung	Sebexol [®] basic pH 5 formula ...	1328	1
Devesa	Sebexol [®] basic pH 5 formula ...		3
Caelo	Sebexol [®] basic pH 5 formula ...	1131	2
Phoenix/Devesa	Sebexol [®] basic pH 5 formula ...	1215	1
Dr.Reingraber GmbH	Sebexol [®] basic pH 5 formula ...	1232	1
Devesa	Sebexol [®] basic pH 5 formula ...	1232	1
Ichthyol	Sebexol [®] basic pH 5 formula ...	1232	2
Caelo	Sebexol [®] basic pH 5 formula ...	1239	1
Devesa	Sebexol [®] basic pH 5 formula ...	1239	1
Devesa Dr.Reingraber Gm...	Sebexol [®] basic pH 5 formula ...	1244	1
Devesa/AHD	Sebexol [®] basic pH 5 formula ...	1245	4
Sebexol/AHD	Sebexol [®] basic pH 5 formula ...	1303	2
Devesa Dr.Reingraber Gm...	Sebexol [®] basic pH 5 formula ...	1303	1
Phönix	Sebexol [®] basic pH 5 formula ...	1313	2
Sebexol	Sebexol [®] basic pH 5 formula ...	1313	1
Sanacorp	Sebexol [®] basic pH 5 formula ...	1313	1
Devesa/Gehe	Sebexol [®] basic pH 5 formula ...		1
Sebexol	Sebexol [®] basic pH 5 formula ...	1333	1
Devesa	Sebexol [®] basic pH 5 formula ...	I232	1
Devesa	Sebexol [®] basic pH 5 formula ...	I307	1
Phönix Gotha	Wofacutan wash lotion	97913	1

- 2997 spectra from 303 *Apo-Ident* customers from a total of 1587 batches from a further 119 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Sebexol[®] formula basis / Sebexol[®] cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Sebexol[®] formula basis / Sebexol[®] cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	550	0	18 199
Type B	0	415	0	22 416
Type C	0	196	0	2997

The substance/substance group *Sebexol[®] formula basis / Sebexol[®] cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9531 %)	100.0000 % (> 98.9091 %)
Type B	100.0000 % (> 99.9528 %)	100.0000 % (> 98.5542 %)
Type C	100.0000 % (> 98.3571 %)	100.0000 % (> 96.9388 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30839	30839	0.00	20.87
30972	30972	0.00	25.04
31065	31065	0.00	26.47
31103	31103	0.00	21.01
31104	31104	0.00	19.09
31592	31592	0.00	19.10
31839	31839	0.00	17.51
31961	31961	0.00	16.40
32743	32743	0.00	19.71
32744	32744	0.00	16.20
33190	33190	0.00	20.65
33191	33191	0.00	19.49
33483	33483	0.00	26.04

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Siam benzoin**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31490-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Siam benzoin

Special notes

When selecting the *Siam benzoin* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Siam benzoin*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Primavera	Siam benzoin	3602151	31490	40	not required
Primavera	Siam benzoin	00051M23	31844	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Siam benzoin*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 80 spectra from 3 reference samples of the substance/substance group *Siam benzoin*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Primavera	Siam benzoin	3602151	31490 [†]	20
Primavera	Siam benzoin	00051M23	31844 [†]	20
Primavera	Siam benzoin	00218C24	32184	40

- 22 751 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Siam benzoin*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Siam benzoin* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Siam benzoin* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	67	13	22 751
Type C	0	0	0	3193

The substance/substance group *Siam benzoin* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9531 %)	83.7500 % (> 80.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31490	31490	0.00	388.08
31844	31844	0.00	288.18

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Solutio Cordes®**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30996-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Solutio Cordes®

Special notes

When selecting the *Solutio Cordes®* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Solutio Cordes®*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Ichthyol	Solutio Cordes®	12B006	30996	40	not required
Ichthyol	Solutio Cordes®	13B001	31076	40	not required
Ichthyol	Solutio Cordes®	15B005	32065	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Solutio Cordes*[®]. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 100 spectra from 3 reference samples of the substance/substance group *Solutio Cordes*[®].
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Ichthyol	<i>Solutio Cordes</i> [®]	09B008	30224	40
Ichthyol	<i>Solutio Cordes</i> [®]	15B005	32065 [†]	20
Ichthyol	<i>Solutio Cordes</i> [®]	15B005	32483	40

- 22 731 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 47 spectra from 15 *Apo-Ident* customers from 15 batches from the substance/substance group *Solutio Cordes*[®].
- These include spectra of independent samples from 12 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Ichthyol /Noweda	Solutio Cordes [®]	138002	1
ICHTYOL-GESELLSCHAFT	Solutio Cordes [®]	13B001	1
Ichthyol/Phönix	Solutio Cordes [®]	13B002	3
Phönix	Solutio Cordes [®]	13B002	1
Caelo	Solutio Cordes [®]	148012	1
Ichthyol / Gehe	Solutio Cordes [®]	14B006	1
Gehe/Ichthyol	Solutio Cordes [®]	14b006	1
Ichthyol Gesellschaft	Solutio Cordes [®]	14B006	1
Ichthyol/Phönix	Solutio Cordes [®]	14B011	1
Ichthyol Gesellschaft	Solutio Cordes [®]	14B012	1
Ichthyol / G	Solutio Cordes [®]	14B012	1
Ichthyol/Noweda	Solutio Cordes [®]	14B012	1
Ichthyol / K	Solutio Cordes [®]	15B005	3
Ichthyol/Phönix	Solutio Cordes [®]	15B001	1
Ichthyol/Noweda	Solutio Cordes [®]	15B005	1
Ichthyol-Gesellschaft	Solutio Cordes [®]	15B005	1
Ichthyol / G	Solutio Cordes [®]	16B005	2
Fiebig	Solutio Cordes [®]	22314	1
Ichthyol/Phönix	Solutio Cordes [®]	12B0005	1
Ichthyol/Phönix	Solutio Cordes [®]	12B005	15
Ichthyol	Solutio Cordes [®]	12B005	1
Ichthyol /Noweda	Solutio Cordes [®]	12B005	1
Ichtyol	Solutio Cordes [®]	12B006	2
Ichthyol	Solutio Cordes [®]	12B006	1
Ichthyol	Solutio Cordes [®]	13B001	1
Ichthyol- Gesellschaft ...	Solutio Cordes [®]	13B001	1
Ichtyol	Solutio Cordes [®]	13B002	1

- 3146 spectra from 304 *Apo-Ident* customers from a total of 1644 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Solutio Cordes*[®] can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Solutio Cordes*[®] and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	70	30	22 731
Type C	0	42	5	3146

The substance/substance group *Solutio Cordes*[®] can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	70.0000 % (> 67.0000 %)
Type C	100.0000 % (> 98.3585 %)	89.3617 % (> 82.9787 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30996	30996	0.00	58.65
31076	31076	0.00	66.43
32065	32065	0.00	65.91

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Spearmint oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30453-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Spearmint oil; Oleum mentha spicata

Special notes

When selecting the *Spearmint oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Spearmint oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Spearmint oil	33894-111575BAG90451	31135	30	not required
Taoasis	Spearmint oil	946-120296	31685	40	not required
Taoasis	Spearmint oil	421012-129111	33481	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 110 spectra of 3 reference samples from the substance/substance group *Spearmint oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 639 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 140 spectra from 4 reference samples of the substance/substance group *Spearmint oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Spearmint oil	27537-96244	30453	40
Taoasis	Spearmint oil	946-120296	31685 [†]	20
Taoasis	Spearmint oil	421012-126408	33200	40
Taoasis	Spearmint oil	421012-126408	33201	40

- 22 691 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 3 spectra from 2 *Apo-Ident* customers from 3 batches from the substance/substance group *Spearment oil*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Spearment oil	27537-96244	1
Taoasis	Spearment oil	421012-123260	1
Taoasis	Spearment oil	421012-125373	1

- 3190 spectra from 304 *Apo-Ident* customers from a total of 1656 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Spearment oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Spearment oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	110	0	18 639
Type B	0	130	10	22 691
Type C	0	3	0	3190

The substance/substance group *Spearment oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 94.5455 %)
Type B	100.0000 % (> 99.9530 %)	92.8571 % (> 90.7143 %)
Type C	100.0000 % (> 98.3847 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31135	31135	0.00	66.97
31685	31685	0.00	63.15
33481	33481	0.00	53.66

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Spruce needle oil / Frankincense oil / Silvire fir oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31032-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Spruce needle oil / Frankincense oil / Silvire fir oil; Frankincense oil; Oleum abies alba; Oleum boswellia serrata; Oleum olibanum; Oleum pini sibiricum; Piceae aetheroleum; Silvire fir oil; Spruce needle oil

Special notes

When selecting the *Spruce needle oil / Frankincense oil / Silvire fir oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Spruce needle oil / Frankincense oil / Silvire fir oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Silvir fir oil	120830-111781	31032	40	not required
Taoasis	Silvir fir oil	211-114751	31255	40	not required

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Frankincense oil	15181-119714	31633	40	not required
Taoasis	Silvir fir oil	01A0B15-121623	31754	40	not required
Taoasis	Spruce needle oil	1507-122520	31819	40	not required
Taoasis	Frankincense oil	1509-122197	31820	40	not required
Taoasis	Spruce needle oil	1719-124011	32066	40	not required
Taoasis	Frankincense oil	1509-125625	33052	40	not required
Taoasis	Frankincense oil	4314-129238	33646	40	not required
Taoasis	Spruce needle oil	10213-149	33726	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 400 spectra of 10 reference samples from the substance/substance group *Spruce needle oil* / *Frankincense oil* / *Silvir fir oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 10 different batches.
- 18 349 spectra from a total of 429 batches from further 148 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 500 spectra from 15 reference samples of the substance/substance group *Spruce needle oil* / *Frankincense oil* / *Silvir fir oil*.
- These include spectra of independent samples from 7 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Frankincense oil	zu erfassen	30298	40
Caelo	Spruce needle oil	11220801	30613	40
Taoasis	Spruce needle oil	29828-103169BAG90451	30652	40
Taoasis	Spruce needle oil	29828-103169BAG90451	30672	40
Taoasis	Frankincense oil	3.26550	30843	40

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Frankincense oil	15181-119714	31633 [†]	20
Taoasis	Silvir fir oil	01A0B15-121623	31754 [†]	20
Taoasis	Spruce needle oil	1507-122520	31819 [†]	20
Taoasis	Frankincense oil	1509-122197	31820 [†]	20
Taoasis	Spruce needle oil	1719-124011	32066 [†]	20
Taoasis	Spruce needle oil	1769-124969	32479	40
Taoasis	Silvir fir oil	01A0B16-125623	33026	40
Taoasis	Silvir fir oil	01A0B16-125623	33027	40
Taoasis	Frankincense oil	1500910-128786	33650	40
Taoasis	Frankincense oil	1500910-128786	33652	40

- 22 331 spectra from a total of 538 batches from further 200 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 22 spectra from 12 *Apo-Ident* customers from 17 batches from the substance/substance group *Spruce needle oil* / *Frankincense oil* / *Silvir fir oil*.
- These include spectra of independent samples from 17 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Spruce needle oil	1769-124949	1
Taoasis	Spruce needle oil	1769-124969	2
Caelo	Spruce needle oil	10185604	1
Caelo	Spruce needle oil	11324401	4
Caelo	Spruce needle oil	13152001	2
Primavera	Spruce needle oil	1670814	1
Taoasis	Spruce needle oil	29828-107964	1
Taoasis	Spruce needle oil	442-113922	1
Taoasis	Spruce needle oil	531766-109893	1
Taoasis	Spruce needle oil	534374-22869	1
Taoasis	Spruce needle oil	534474-111585	1
Taoasis/Sanacorp	Spruce needle oil	534474-111869	1
Taoasis/Sanacorp	Spruce needle oil	534474-113171	1
Apoth.Bauer&Co/Noweda	Frankincense oil		1
Taoasis	Frankincense oil	549702-112299	1
Apoth.Bauer&Co/Noweda	Frankincense oil	3175391505	1
Taoasis	Silvir fir oil	212-120342	1

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

- 3171 spectra from 304 *Apo-Ident* customers from a total of 1642 batches from a further 119 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Spruce needle oil / Frankincense oil / Silvir fir oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Spruce needle oil / Frankincense oil / Silvir fir oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	400	0	18 349
Type B	0	205	295	22 293
Type C	0	6	16	3171

The substance/substance group *Spruce needle oil / Frankincense oil / Silvir fir oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9532 %)	100.0000 % (> 98.5000 %)
Type B	100.0000 % (> 99.9532 %)	41.0000 % (> 40.4000 %)
Type C	100.0000 % (> 98.3605 %)	27.2727 % (> 13.6364 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31032	31032	0.00	32.74
31255	31255	0.00	38.16
31633	31633	0.00	73.12
31754	31754	0.00	20.70
31819	31819	0.00	16.53
31820	31820	0.00	38.24
32066	32066	0.00	20.98
33052	33052	0.00	39.05
33646	33646	0.00	17.96
33726	33726	0.00	14.43

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Stomahesive® adhesive paste**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30974-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Stomahesive® adhesive paste

Special notes

When selecting the *Stomahesive® adhesive paste* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Stomahesive® adhesive paste*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
ConvaTec Inc.	Stomahesive® adh...	2D117	30974	70	not required
ConvaTec Inc.	Stomahesive® adh...	2M084	31087	40	not required
ConvaTec Inc.	Stomahesive® adh...	6L039	33474	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 150 spectra of 3 reference samples from the substance/substance group *Stomahesive® adhesive paste*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 599 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 114 spectra from 4 reference samples of the substance/substance group *Stomahesive® adhesive paste*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
ConvaTec Inc.	Stomahesive® adhesive paste	2D117	30974 [†]	10
ConvaTec Inc.	Stomahesive® adhesive paste	6D079	32761	40
ConvaTec Inc.	Stomahesive® adhesive paste	5G062	32954	40
Convatec	Stomahesive® adhesive paste	5G062	not required	24

- 22 717 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Stomahesive® adhesive paste*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Stomahesive® adhesive paste* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Stomahesive® adhesive paste* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	150	0	18 599
Type B	0	114	0	22 717
Type C	0	0	0	3193

The substance/substance group *Stomahesive® adhesive paste* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9535 %)	100.0000 % (> 96.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.7368 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30974	30974	0.00	51.54
31087	31087	0.00	56.91
33474	33474	0.00	56.25

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Strawberry aroma**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30643-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Strawberry aroma; Strawberry flavour

Special notes

When selecting the *Strawberry aroma* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Strawberry aroma*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Strawberry aroma	11178804	30643	40	1402439
Caelo	Strawberry aroma	13252407	31395	40	not required
Caelo	Strawberry aroma	15411807	32814	40	not required
Caelo	Strawberry aroma	15411807	32838	40	1704074

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Strawberry aroma*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 589 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 160 spectra from 4 reference samples of the substance/substance group *Strawberry aroma*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	Strawberry aroma	14311215	32567	40
Caelo	Strawberry aroma	14311215	32837	40
Caelo	Strawberry aroma	170246	33056	40
Caelo	Strawberry aroma	180832	34077	40

- 22 671 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details

provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 27 spectra from 17 *Apo-Ident* customers from 15 batches from the substance/substance group *Strawberry aroma*.
- These include spectra of independent samples from 13 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Strawberry aroma	9101405	1
Caelo	Strawberry aroma	12183513	4
Caelo	Strawberry aroma	13252405	1
Caelo	Strawberry aroma	13252407	2
Caelo	Strawberry aroma	14311204	1
Caelo	Strawberry aroma	13252417	1
Caelo	Strawberry aroma	14311209	4
Caelo	Strawberry aroma	14311215	2
Hedinger	Strawberry aroma	14311215	1
Caelo	Strawberry aroma	15411805	3
Caelo	Strawberry aroma	15411807	1
Caelo	Strawberry aroma	15411813	1
Caelo	Strawberry aroma	10184917	1
Caelo	Strawberry aroma	11178813	1
Caelo	Strawberry aroma	12183504	2
Caelo	Strawberry aroma	12183511	1

- 3166 spectra from 303 *Apo-Ident* customers from a total of 1644 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Strawberry aroma* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Strawberry aroma* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	160	0	22 671
Type C	0	27	0	3166

The substance/substance group *Strawberry aroma* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9529 %)	100.0000 % (> 96.2500 %)
Type C	100.0000 % (> 98.3598 %)	100.0000 % (> 77.7778 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30643	30643	0.00	112.15
31395	31395	0.00	88.82
32814	32814	0.00	103.87
32838	32838	0.00	70.74

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Swiss pine oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31553-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Swiss pine oil; Oleum pinus cembra

Special notes

When selecting the *Swiss pine oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Swiss pine oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Swiss pine oil	4892-117271	31553	40	not required
Taoasis	Swiss pine oil	3023-120087	31680	40	not required
Taoasis	Swiss pine oil	12069-127103	32934	40	not required
Taoasis	Swiss pine oil	10304-128929	33645	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 160 spectra of 4 reference samples from the substance/substance group *Swiss pine oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 4 different batches.
- 18 589 spectra from a total of 435 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 80 spectra from 3 reference samples of the substance/substance group *Swiss pine oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Swiss pine oil	4892-117271	31553 [†]	20
Taoasis	Swiss pine oil	3023-120087	31680 [†]	20
Taoasis	Swiss pine oil	201610159-126370	32793	40

- 22 751 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Swiss pine oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Swiss pine oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Swiss pine oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	160	0	18 589
Type B	0	40	40	22 748
Type C	0	0	0	3193

The substance/substance group *Swiss pine oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9534 %)	100.0000 % (> 96.2500 %)
Type B	100.0000 % (> 99.9535 %)	50.0000 % (> 46.2500 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31553	31553	0.00	20.90
31680	31680	0.00	18.29
32934	32934	0.00	27.88
33645	33645	0.00	22.04

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Tea tree oil
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	30959-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Tea tree oil; Melaleuca alternifolia oleum; Oleum melaleuca alternifolia

Special notes

When selecting the *Tea tree oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7	European Pharmacopoeia 8th Edition, Basic Version 2014 [3]
Komm2.2.40	Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident [4]
AA004	Erstellung und Validierung eines IdentModul-Updates
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Tea tree oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Tea tree oil	533902-109942BAG91303	30959	40	not required
Taoasis	Tea tree oil	34828-113642BAG91303	31134	40	not required
Taoasis	Tea tree oil	44737-121997	31762	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Tea tree oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 112 spectra from 4 reference samples of the substance/substance group *Tea tree oil*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Tea tree oil	44737-121997	31762 [†]	20
Taoasis	Tea tree oil	52642-125539	33017	40
Taoasis	Tea tree oil	52642-124527	33018	40
Taoasis	Tea tree oil	52642-125539BAG91303	not required	12

- 22 719 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 16 spectra from 5 *Apo-Ident* customers from 12 batches from the substance/substance group *Tea tree oil*.
- These include spectra of independent samples from 12 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Tea tree oil	1469-117418	2
Caelo	Tea tree oil	15051806	2
Caelo	Tea tree oil	15051807	1
Caelo	Tea tree oil	15337105	1
Caelo	Tea tree oil	15337106	1
Taoasis	Tea tree oil	20269-116833	2
Taoasis	Tea tree oil	28348-119841	1
Taoasis	Tea tree oil	28348-121515	1
Taoasis	Tea tree oil	28348-121597	2
Taoasis	Tea tree oil	52642-12078	1
Taoasis	Tea tree oil	52642-124559	1
Taoasis	Tea tree oil	52642-124919	1

- 3177 spectra from 303 *Apo-Ident* customers from a total of 1647 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Tea tree oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Tea tree oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	112	0	22 719
Type C	0	16	0	3177

The substance/substance group *Tea tree oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 94.6429 %)
Type C	100.0000 % (> 98.3619 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several

new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30959	30959	0.00	89.20
31134	31134	0.00	92.32
31762	31762	0.00	101.25

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Thyme oil, red**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30929-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Thyme oil, red; Oleum thymus vulgaris

Special notes

When selecting the *Thyme oil, red* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Thyme oil, red*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Thyme oil, red	31911-109378	30929	40	not required
Taoasis	Thyme oil, red	9434-115427	31259	40	not required
Taoasis	Thyme oil, red	56081-124964	32482	40	not required
Taoasis	Thyme oil, red	1829-125364	32981	40	not required
Taoasis	Thyme oil, red	1976-2943	34103	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 200 spectra of 5 reference samples from the substance/substance group *Thyme oil, red*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 5 different batches.
- 18 549 spectra from a total of 434 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 40 spectra from 1 reference samples of the substance/substance group *Thyme oil, red*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Thyme oil, red	40517-121881	32472	40

- 22 791 spectra from a total of 548 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 15 spectra from 2 *Apo-Ident* customers from 11 batches from the substance/substance group *Thyme oil, red*.
- These include spectra of independent samples from 11 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Thyme oil, red	43637-122852	1
Taoasis	Thyme oil, red	43637-124964	1
Taoasis	Thyme oil, red	14381-117385	1
Taoasis	Thyme oil, red	25394-118296	1
Taoasis	Thyme oil, red	32444-118785	4
Taoasis	Thyme oil, red	33635-120088	1
Taoasis	Thyme oil, red	40517-121881	2
Taoasis	Thyme oil, red	43637-122713	1
Taoasis	Thyme oil, red	43637-123574	1
Taoasis	Thyme oil, red	9434-115525	1
Taoasis	Thyme oil, red	33092-111620	1

- 3178 spectra from 304 *Apo-Ident* customers from a total of 1648 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Thyme oil, red* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Thyme oil, red* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	200	0	18 549
Type B	0	40	0	22 791
Type C	0	13	2	3178

The substance/substance group *Thyme oil, red* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.0000 %)
Type B	100.0000 % (> 99.9535 %)	100.0000 % (> 85.0000 %)
Type C	100.0000 % (> 98.3623 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30929	30929	0.00	27.59
31259	31259	0.00	20.86
32482	32482	0.00	42.52
32981	32981	0.00	34.12
34103	34103	0.00	83.41

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Thyme oil, white / Coriander oil
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	31552-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Thyme oil, white / Coriander oil; Coriander oil; Oleum coriandri; Thyme oil, white

Special notes

When selecting the *Thyme oil, white / Coriander oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Thyme oil, white / Coriander oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Coriander oil	9324-114961	31552	40	not required
Taoasis	Coriander oil	25134-117444	31690	40	not required
Taoasis	Coriander oil	104709-126301	32792	40	not required
Primavera	Thyme oil, white	01067J25	33796	40	not required
Taoasis	Thyme oil, white	111400-970	33818	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 200 spectra of 5 reference samples from the substance/substance group *Thyme oil, white / Coriander oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 5 different batches.
- 18 549 spectra from a total of 434 batches from further 149 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 160 spectra from 5 reference samples of the substance/substance group *Thyme oil, white / Coriander oil*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Coriander oil	9324-114961	31552 [†]	20
Taoasis	Coriander oil	25134-117444	31690 [†]	20
Taoasis	Coriander oil	104709-127188	33053	40
Taoasis	Thyme oil, white	925	33819	40
Taoasis	Coriander oil	9291-1307	33990	40

- 22 671 spectra from a total of 544 batches from further 201 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B*

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 2 spectra from 2 *Apo-Ident* customers from 2 batches from the substance/substance group *Thyme oil, white / Coriander oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Coriander oil	15229607	1
Caelo	Coriander oil	16140401	1

- 3191 spectra from 304 *Apo-Ident* customers from a total of 1657 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Thyme oil, white / Coriander oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Thyme oil, white / Coriander oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	200	0	18 549
Type B	0	147	13	22 671
Type C	0	2	0	3191

The substance/substance group *Thyme oil, white / Coriander oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.0000 %)
Type B	100.0000 % (> 99.9529 %)	91.8750 % (> 90.0000 %)
Type C	100.0000 % (> 98.3987 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31552	31552	0.00	22.51
31690	31690	0.00	21.55
32792	32792	0.00	30.51
33796	33796	0.00	41.68
33818	33818	0.00	23.14

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Vetiver bourbon oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31546-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Vetiver bourbon oil; Oleum vetiveria zizanoides

Special notes

When selecting the *Vetiver bourbon oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Vetiver bourbon oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Vetiver bourbon ...	7141-116418	31546	40	not required
Taoasis	Vetiver bourbon ...	7141-119962	31684	40	not required
Taoasis	Vetiver bourbon ...	7141-128198	33647	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Vetiver bourbon oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *Vetiver bourbon oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Vetiver bourbon oil	7141-116418	31546 [†]	20
Taoasis	Vetiver bourbon oil	7141-119962	31684 [†]	20
Taoasis	Vetiver bourbon oil	7141-124922	32810	40
Taoasis	Vetiver bourbon oil	7141-124922	33046	40

- 22 711 spectra from a total of 546 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 5 spectra from 1 *Apo-Ident* customers from 3 batches from the substance/substance group *Vetiver bourbon oil*.
- These include spectra of independent samples from 3 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Taoasis	Vetiver bourbon oil	7141-122195	2
Taoasis	Vetiver bourbon oil	7141-119174	1
Taoasis	Vetiver bourbon oil	7141-124600	2

- 3188 spectra from 304 *Apo-Ident* customers from a total of 1656 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Vetiver bourbon oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Vetiver bourbon oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	120	0	22 711
Type C	0	5	0	3188

The substance/substance group *Vetiver bourbon oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 95.0000 %)
Type C	100.0000 % (> 98.3735 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31546	31546	0.00	31.03
31684	31684	0.00	29.75
33647	33647	0.00	28.37

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Wart ointment InfectoPharm® (NRF 11.31)**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30730-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Wart ointment InfectoPharm® (NRF 11.31)

Special notes

When selecting the *Wart ointment InfectoPharm® (NRF 11.31)* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Wart ointment InfectoPharm® (NRF 11.31)*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Infectopharm	Wart ointment In...	S031201.1	30730	40	not required
Infectopharm	Wart ointment In...	S031201.1	30905	40	not required
Infectopharm	Wart ointment In...	S121103.1	30906	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Wart ointment InfectoPharm®* (NRF 11.31). These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 629 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 3 reference samples of the substance/substance group *Wart ointment InfectoPharm®* (NRF 11.31).
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Infectopharm	Wart ointment InfectoPharm® ...	S021601.1	32593	40
Infectopharm	Wart ointment InfectoPharm® ...	S071602.1	32953	40
Infectopharm	Wart ointment InfectoPharm® ...	S071602.1	33515	40

- 22 711 spectra from a total of 547 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 78 spectra from 42 *Apo-Ident* customers from 24 batches from the substance/substance group *Wart ointment InfectoPharm® (NRF 11.31)*.
- These include spectra of independent samples from 6 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
InfectoPharm/Anzag	Wart ointment InfectoPharm® ...	716A-03898	1
Infectopharm	Wart ointment InfectoPharm® ...		25
Euro OTC	Wart ointment InfectoPharm® ...		1
gehe/infecto	Wart ointment InfectoPharm® ...		1
Infectopharm / Gehe	Wart ointment InfectoPharm® ...		1
InfectopharmNoweda	Wart ointment InfectoPharm® ...		1
Infectopharm/Phoenix Le...	Wart ointment InfectoPharm® ...		1
Infectopharm/Gehe	Wart ointment InfectoPharm® ...		1
Infectopharm Arzneimittel	Wart ointment InfectoPharm® ...		1
Infectopharm/Kehr	Wart ointment InfectoPharm® ...		1
Infectopharm/K	Wart ointment InfectoPharm® ...		1
InfectoPharm	Wart ointment InfectoPharm® ...	S0215011	1
PX/Infectop	Wart ointment InfectoPharm® ...	S0215011	3
Infectopharm/Noweda	Wart ointment InfectoPharm® ...		4
Infectopharm/Phönix	Wart ointment InfectoPharm® ...		2
Infectopharm/Alliance H...	Wart ointment InfectoPharm® ...		1
InfectoPharm/Noweda	Wart ointment InfectoPharm® ...		2
Eigenherstellung	Wart ointment InfectoPharm® ...		1
Infectopharm / Alliance...	Wart ointment InfectoPharm® ...		1
Infecto/Spangro	Wart ointment InfectoPharm® ...		1
InfectoPharm	Wart ointment InfectoPharm® ...		2
Sanacorp	Wart ointment InfectoPharm® ...		2
Infectopharm/G	Wart ointment InfectoPharm® ...		1
Noweda/Infectopharm	Wart ointment InfectoPharm® ...		1
Infectopharm/Phoenix	Wart ointment InfectoPharm® ...		1
InfectoPharm/Anzag	Wart ointment InfectoPharm® ...		2
Fagron	Wart ointment InfectoPharm® ...		1
Ichthyol-Gesellschaft C...	Wart ointment InfectoPharm® ...		1
InfectoPharm/AHD	Wart ointment InfectoPharm® ...		1
Infecto/Gehe	Wart ointment InfectoPharm® ...		1
Phönix Gotha	Wart ointment InfectoPharm® ...		1
Infect/PX	Wart ointment InfectoPharm® ...		2
infectopharm	Wart ointment InfectoPharm® ...		2
Löwen City-Apotheke Bar...	Wart ointment InfectoPharm® ...	8032013	1
Infectopharm	Wart ointment InfectoPharm® ...	3013I-07003	1
nfectoPharm	Wart ointment InfectoPharm® ...		1
InfectoPharm/ E	Wart ointment InfectoPharm® ...		1
infectipharm/Phönix	Wart ointment InfectoPharm® ...		1
Caelo	Wart ointment InfectoPharm® ...		1
Gehe	Wart ointment InfectoPharm® ...		1
Infectopharm (Phoenix)	Wart ointment InfectoPharm® ...		1
Infectopharm	Wart ointment InfectoPharm® ...	S111202	1

- 3115 spectra from 296 *Apo-Ident* customers from a total of 1636 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Wart ointment InfectoPharm®* (NRF 11.31) can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Wart ointment InfectoPharm®* (NRF 11.31) and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	120	0	22 711
Type C	0	78	0	3115

The substance/substance group *Wart ointment InfectoPharm®* (NRF 11.31) can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	100.0000 % (> 95.0000 %)
Type C	100.0000 % (> 98.3577 %)	100.0000 % (> 92.3077 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
30730	30730	0.00	234.49
30905	30905	0.00	228.33
30906	30906	0.00	223.94

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group	Water and aqueous solutions
Substance class	APIs & excipients, liquid/semi-solid (other)
Report date	18/07/2018
Report number	31047-2018-07-18
Executing company	HiperScan GmbH Weißeritzstraße 3 01067 Dresden Germany

Relevant substance names

Water and aqueous solutions; Aloe Vera gel, 1:1; Aloe vera gel, 10x concentrated; Aqua; Bepanthen[®] solution; Retterspitz external; SyrSpend[®] SF pH4 cherry aroma; SyrSpend[®] SF pH4 aroma free; Water

Special notes

When selecting the *Water and aqueous solutions* substance/substance group, the following information is displayed to the user:

- Advice for the identification of water: Preserved, purified and distilled water can not be conclusively distinguished by spectral analysis. An additional analysis of identity may be required.

Applicable documents

978-3-7692-5416-7	<i>European Pharmacopoeia 8th Edition, Basic Version 2014</i> [3]
Komm2.2.40	<i>Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident</i> [4]
AA004	<i>Erstellung und Validierung eines IdentModul-Updates</i>
Anhang F	Zusatz zu den Modellen der zweiten Stufe

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Water and aqueous solutions*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Bayer	Bepanthen [®] solut...	KP08DR1	31047	40	not required
Caelo	Aloe Vera gel, 1:1	13025802	31145	40	not required

continued on the next page

continued from previous page

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Aloe vera gel, 1...	13045901	31146	40	not required
Hedinger	Water	219066	31623	40	not required
Fagron	SyrSpend [®] SF pH4...	1503D009	31670	40	not required
Fagron	SyrSpend [®] SF pH4...	1503D017	31671	40	not required
Caelo	Water	15068701	31675	40	not required
Fagron	SyrSpend [®] SF pH4...	1412D002	31722	40	not required
Fagron	SyrSpend [®] SF pH4...	1503D017	31723	40	not required
Bayer	Bepanthen [®] solut...	KPOASHC	31857	40	not required
Caelo	Aloe vera gel, 1...	15185503	31862	40	not required
Caelo	Aloe Vera gel, 1:1	15329203	31866	40	not required
Caelo	Aloe Vera gel, 1:1	153292	31880	40	20151111*
Caelo	Aloe Vera gel, 1:1	153292	31881	40	20151111*
Caelo	Water	153943	32079	40	20151215*
Caelo	Aloe vera gel, 1...	161922	32313	40	20160707*
Retterspitz G...	Retterspitz exte...	016100	32785	40	not required
Retterspitz G...	Retterspitz exte...	017058	33430	40	not required
Retterspitz G...	Retterspitz exte...	016240	33431	40	not required
Bayer	Bepanthen [®] solut...	KPOC3FU	33612	40	not required
Fagron	SyrSpend [®] SF pH4...	1704492/1	33787	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 840 spectra of 21 reference samples from the substance/substance group *Water and aqueous solutions*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 19 different batches.
- 17 909 spectra from a total of 420 batches from further 144 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

*The spectra were taken from a part of the same sample which the supplier used for its analysis (identity, purity, content). In this case the manufacturer certificates proof the identity and quality of the sample from those the spectra descend.

- 1490 spectra from 43 reference samples of the substance/substance group *Water and aqueous solutions*.
- These include spectra of independent samples from 30 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Hedinger	Water	219066	31623 [†]	20
Fagron	SyrSpend [®] SF pH4 aroma free	1503D009	31670 [†]	20
Fagron	SyrSpend [®] SF pH4 cherry aroma	1503D017	31671 [†]	20
Caelo	Water	15068701	31675 [†]	20
Fagron	SyrSpend [®] SF pH4 aroma free	1412D002	31722 [†]	20
Fagron	SyrSpend [®] SF pH4 cherry aroma	1503D017	31723 [†]	20
Bayer	Bepanthen [®] solution	KP0ASHC	31857 [†]	20
Caelo	Aloe vera gel, 10x concentra...	15185503	31862 [†]	20
Caelo	Aloe Vera gel, 1:1	15329203	31866 [†]	20
Caelo	Aloe Vera gel, 1:1	153292	31880 [†]	20
Caelo	Aloe Vera gel, 1:1	153292	31881 [†]	20
Caelo	Water	153944	32081	40
Caelo	Water	153712	32083	40
Caelo	Aloe Vera gel, 1:1	162489	32580	40
Bayer	Bepanthen [®] solution	KP0B60L	32719	40
Fagron	SyrSpend [®] SF pH4 aroma free	1507402	32737	40
Fagron	SyrSpend [®] SF pH4 cherry aroma	1503255	32738	40
Retterspitz GmbH	Retterspitz external	016179	32784	30
Fagron	SyrSpend [®] SF pH4 cherry aroma	1603252/2	32842	40
Fagron	SyrSpend [®] SF pH4 aroma free	1507402	32843	40
Caelo	Water	162553	32895	40
Caelo	Water	162867	32896	40
Caelo	Aloe Vera gel, 1:1	162912	32899	40
Bayer	Bepanthen [®] solution	KP0BJ61	32949	40
Caelo	Water	163533	33176	40
Caelo	Water	170018	33177	40
Caelo	Aloe Vera gel, 1:1	171341	33411	40
Fagron	SyrSpend [®] SF pH4 aroma free	1704493	33501	40
Caelo	Aloe vera gel, 10x concentra...	17082002	33511	40
Caelo	Water	172191	33578	40
Caelo	Water	172192	33579	40
Caelo	Water	172220	33580	40
Fagron	SyrSpend [®] SF pH4 aroma free	1704493	33631	40
Caelo	Aloe Vera gel, 1:1	172891	33736	40
Caelo	Water	172533	33741	40
Caelo	Water	172827	33742	40
Caelo	Water	172861	33743	40
Caelo	Water	172862	33744	40
Caelo	Water	172771	33745	40
Caelo	Aloe Vera gel, 1:1	172890	33970	40
Caelo	Water	171147	33977	40
Caelo	Water	172957	33978	40
Retterspitz GmbH	Retterspitz external	017192	34049	40

- 21 341 spectra from a total of 510 batches from further 196 substances. These spectra were

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 50 spectra from 30 *Apo-Ident* customers from 29 batches from the substance/substance group *Water and aqueous solutions*.
- These include spectra of independent samples from 27 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Aloe vera gel, 10x concentra...	15185512	1
Fagron	Water	15H25-T21-025683	1
Fagron	SyrSpend® SF pH4 aroma free	1410D020	1
Fagron	SyrSpend® SF pH4 cherry aroma	1503255	7
Fagron	SyrSpend® SF pH4 cherry aroma	1507318	2
Fagron	SyrSpend® SF pH4 aroma free	1503D009	1
Fagron	SyrSpend® SF pH4 cherry aroma	1503D017	1
Caelo	SyrSpend® SF pH4 cherry aroma	1507318	1
Fagron	SyrSpend® SF pH4 aroma free	1507402	6
Caelo	SyrSpend® SF pH4 aroma free	1507402	1
Apomix/ Sanacorp	SyrSpend® SF pH4 cherry aroma	15C19/C	1
Fagron	SyrSpend® SF pH4 aroma free	1605291	5
Fagron	SyrSpend® SF pH4 cherry aroma	15C19/C	1
Fagron	SyrSpend® SF pH4 aroma free	1609686	1
Fagron	SyrSpend® SF pH4 cherry aroma	1612631	2
Fagron	SyrSpend® SF pH4 aroma free	1615584/2	1
Fagron	SyrSpend® SF pH4 aroma free	161558412	1
Fagron	SyrSpend® SF pH4 aroma free	1704493	2
Fagron	SyrSpend® SF pH4 aroma free	1706769/1	1
Caelo	Water	15394201	1
Fagron	Water	15H25-T14-025670	1
Fagron	Water	15H25-T18-025678	1
Phoenix	Water	15L14-T14-028664	1
Fagron	Water	15L15-T01-028672	1
Fagron	Water	16C23-T06-031423	1
Fagron	Water	16F13-T16-034297	1
Euro OTC	Water	16F14-T07-034326	1
Caelo	Water		1
WEPA	Water	LOTAP0015	1
wepa	Water	LOTAP0019	1
Wepa	Water	LOTAP0031	1
Wepa/Wepa	Water	LOTAP0037	1

- 3143 spectra from 300 *Apo-Ident* customers from a total of 1630 batches from a further 118 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Water and aqueous solutions* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Water and aqueous solutions* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	840	0	17 909
Type B	0	1489	1	21 001
Type C	0	50	0	3117

The substance/substance group *Water and aqueous solutions* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9531 %)	100.0000 % (> 99.2857 %)
Type B	100.0000 % (> 99.9536 %)	99.9329 % (> 99.7315 %)
Type C	100.0000 % (> 98.3727 %)	100.0000 % (> 88.0000 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31047	31047	0.00	25.88
31145	31145	0.00	30.93
31146	31146	0.00	27.87
31623	31623	0.00	29.64
31670	31670	0.00	29.17
31671	31671	0.00	29.99
31675	31675	0.00	32.37
31722	31722	0.00	27.90
31723	31723	0.00	28.16
31857	31857	0.00	25.25
31862	31862	0.00	27.66
31866	31866	0.00	30.89
31880	31880	0.00	32.85
31881	31881	0.00	30.38
32079	32079	0.00	30.96
32313	32313	0.00	26.78

continued on the next page

continued from previous page

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
32785	32785	0.00	19.79
33430	33430	0.00	25.54
33431	33431	0.00	25.08
33612	33612	0.00	25.65
33787	33787	0.00	27.27

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Wax ointment (stabilised)**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31111-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Wax ointment (stabilised); Unguentum cereum

Special notes

When selecting the *Wax ointment (stabilised)* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Wax ointment (stabilised)*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	Wax ointment (st...	13072102	31111	40	not required
Caelo	Wax ointment (st...	13307701	31167	40	not required
Caelo	Wax ointment (st...	14073701	31444	40	not required
Caelo	Wax ointment (st...	154243	31911	40	20151217*
Caelo	Wax ointment (st...	154244	31912	40	20151217*

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 200 spectra of 5 reference samples from the substance/substance group *Wax ointment (stabilised)*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 5 different batches.
- 18 549 spectra from a total of 434 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 280 spectra from 8 reference samples of the substance/substance group *Wax ointment (stabilised)*.
- These include spectra of independent samples from 6 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	Wax ointment (stabilised)	09361101	30267	40
Caelo	Wax ointment (stabilised)	12352501	30967	40
Caelo	Wax ointment (stabilised)	154243	31911 [†]	20
Caelo	Wax ointment (stabilised)	154244	31912 [†]	20
Caelo	Wax ointment (stabilised)	161215	32117	40
Caelo	Wax ointment (stabilised)	161334	32118	40
Caelo	Wax ointment (stabilised)	171714	33550	40
Caelo	Wax ointment (stabilised)	172942	34024	40

*The spectra were taken from a part of the same sample which the supplier used for its analysis (identity, purity, content). In this case the manufacturer certificates proof the identity and quality of the sample from those the spectra descend.

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

- 22 551 spectra from a total of 541 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 201 spectra from 83 *Apo-Ident* customers from 45 batches from the substance/substance group *Wax ointment (stabilised)*.
- These include spectra of independent samples from 42 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Wax ointment (stabilised)	13307802	11
VDL;6,45EUR;21.10.14	Wax ointment (stabilised)	14235901	1
Caelo	Wax ointment (stabilised)	14235901	11
Caelo	Wax ointment (stabilised)	15025801	15
Caelo	Wax ointment (stabilised)	15205402	8
Caelo	Wax ointment (stabilised)	15424301	9
Caelo	Wax ointment (stabilised)	16133401	8
Caelo	Wax ointment (stabilised)	12033701	6
Caelo	Wax ointment (stabilised)	1218702	2
Caelo	Wax ointment (stabilised)	12352501	4
Caelo	Wax ointment (stabilised)	1021303	1
Caelo	Wax ointment (stabilised)	7071204	1
Caelo	Wax ointment (stabilised)	11321302	2
Caelo	Wax ointment (stabilised)	11355202	1
Caelo	Wax ointment (stabilised)	12033602	2
Anzag 07.12.12	Wax ointment (stabilised)	12128702	1
Caelo	Wax ointment (stabilised)	12209402	1
Caelo	Wax ointment (stabilised)	12352051	1
Caelo	Wax ointment (stabilised)	12378701	8
Sanacorp	Wax ointment (stabilised)	12352501	1
Caelo	Wax ointment (stabilised)	13072102	12
Caelo	Wax ointment (stabilised)	13159302	8
Caelo	Wax ointment (stabilised)	13307701	16
Caelo	Wax ointment (stabilised)	14073701	13
Caelo	Wax ointment (stabilised)	20051012-1	1
Caelo	Wax ointment (stabilised)	20121305	1
Caelo	Wax ointment (stabilised)	14072801	1
Caelo	Wax ointment (stabilised)	1407301	1
Sanacorp	Wax ointment (stabilised)	14073701	1
AHD	Wax ointment (stabilised)	14073701	1
Caelo	Wax ointment (stabilised)	140737012	1
Caelo	Wax ointment (stabilised)	140737701	1
Caelo	Wax ointment (stabilised)	14073801	6
Caelo	Wax ointment (stabilised)	14236001	2
Caelo	Wax ointment (stabilised)	14236002	2
Caelo	Wax ointment (stabilised)	15025702	8
Sanacorp	Wax ointment (stabilised)	15205402	2
Caelo	Wax ointment (stabilised)	1524301	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Henry Lamotte Olis GmbH...	Wax ointment (stabilised)	15424301	1
Caeol/Phoenix	Wax ointment (stabilised)	15424301	1
Caelo	Wax ointment (stabilised)	15424401	1
Caelo	Wax ointment (stabilised)	15424402	2
Caelo	Wax ointment (stabilised)	16121501	4
Gehe	Wax ointment (stabilised)	16133401	1
Caelo	Wax ointment (stabilised)	17014901	2
Caelo	Wax ointment (stabilised)	17015001	2
Caelo	Wax ointment (stabilised)	2113Q-03212	1
Caelo	Wax ointment (stabilised)	28051511	1
Caelo	Wax ointment (stabilised)	1114E3212	1
Caelo	Wax ointment (stabilised)	11355201	2
Anzag	Wax ointment (stabilised)	12033701	1
Caelo	Wax ointment (stabilised)	12209401	3
Caesar & Loretz GmbH	Wax ointment (stabilised)	13072102	1
Gehe	Wax ointment (stabilised)	13072102	1
Caelo	Wax ointment (stabilised)	13159303	2
Caelo	Wax ointment (stabilised)	U-150513-2	1
Celo	Wax ointment (stabilised)	15424402	1

- 2992 spectra from 296 *Apo-Ident* customers from a total of 1614 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Wax ointment (stabilised)* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Wax ointment (stabilised)* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	200	0	18 549
Type B	0	280	0	22 551
Type C	0	193	8	2992

The substance/substance group *Wax ointment (stabilised)* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9533 %)	100.0000 % (> 97.0000 %)
Type B	100.0000 % (> 99.9529 %)	100.0000 % (> 97.8571 %)
Type C	100.0000 % (> 98.3571 %)	96.0199 % (> 94.5274 %)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to

exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31111	31111	0.00	35.53
31167	31167	0.00	39.11
31444	31444	0.00	37.49
31911	31911	0.00	37.86
31912	31912	0.00	39.90

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **White paraffin oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 32342-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

White paraffin oil; Oleum Vaselini album

Special notes

When selecting the *White paraffin oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *White paraffin oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Caelo	White paraffin oil	162192	32342	40	20160729*
Caelo	White paraffin oil	15380806	32565	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *White paraffin oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 80 spectra from 2 reference samples of the substance/substance group *White paraffin oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Caelo	White paraffin oil	14338604	32454	40
Caelo	White paraffin oil	14338604	32846	40

- 22 751 spectra from a total of 548 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

*The spectra were taken from a part of the same sample which the supplier used for its analysis (identity, purity, content). In this case the manufacturer certificates proof the identity and quality of the sample from those the spectra descend.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *White paraffin oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *White paraffin oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *White paraffin oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	80	0	22 751
Type C	0	0	0	3193

The substance/substance group *White paraffin oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9531 %)	100.0000 % (> 92.5000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra ([Rule of Three](#) [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the [Mahalanobis distance](#) to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
32342	32342	0.00	23.67
32565	32565	0.00	25.15

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Wintergreen oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 30455-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Wintergreen oil; Oleum gaultheria procumbens

Special notes

When selecting the *Wintergreen oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Wintergreen oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Wintergreen oil	5411006-117696	31543	40	not required
Taoasis	Wintergreen oil	1828-125286	32807	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Wintergreen oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 172 spectra from 5 reference samples of the substance/substance group *Wintergreen oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Wintergreen oil	423129090971	30455	40
Taoasis	Wintergreen oil	5411006-117696	31543 [†]	16
Taoasis	Wintergreen oil	1828-125286	32936	40
Taoasis	Wintergreen oil	1953-128607	33624	40
Taoasis	Wintergreen oil	1953-128607	33625	36

- 22 659 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 9 spectra from 8 *Apo-Ident* customers from 9 batches from the substance/substance group *Wintergreen oil*.
- These include spectra of independent samples from 9 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Spectra
Caelo	Wintergreen oil	15087406	1
Taoasis	Wintergreen oil	423121-90971	1
Taoasis	Wintergreen oil		1
Taoasis	Wintergreen oil	1303-121125	1
Caelo	Wintergreen oil	10318703	1
Caelo	Wintergreen oil	13099202	1
Caelo	Wintergreen oil	10318714	1
Caelo	Wintergreen oil	13099205	1
Taoasis	Wintergreen oil	97540	1

- 3184 spectra from 304 *Apo-Ident* customers from a total of 1650 batches from a further 121 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Wintergreen oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Wintergreen oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	150	22	22 659
Type C	0	7	2	3184

The substance/substance group *Wintergreen oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9529 %)	87.2093 % (> 85.4651 %)
Type C	100.0000 % (> 98.3660 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31543	31543	0.00	307.92
32807	32807	0.00	310.54

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Yarrow oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 33795-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Yarrow oil

Special notes

When selecting the *Yarrow oil* substance/substance group, the following information is displayed to the user:

no information

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Yarrow oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Primavera	Yarrow oil	00516E25	33795	40	not required
Taoasis	Yarrow oil	2041-316	33822	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 80 spectra of 2 reference samples from the substance/substance group *Yarrow oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 2 different batches.
- 18 669 spectra from a total of 437 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a [†]. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 40 spectra from 1 reference samples of the substance/substance group *Yarrow oil*.
- These include spectra of independent samples from 1 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Yarrow oil	128700	33823	40

- 22 791 spectra from a total of 548 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

Type C Spectra from the field, which were not recorded under of the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Yarrow oil*.

- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Yarrow oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Yarrow oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	80	0	18 669
Type B	0	40	0	22 791
Type C	0	0	0	3193

The substance/substance group *Yarrow oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate* (*specificity*) and the weighted *true positive rate* (*recognition rate*) are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9538 %)	100.0000 % (> 92.5000 %)
Type B	100.0000 % (> 99.9535 %)	100.0000 % (> 85.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
33795	33795	0.00	52.18
33822	33822	0.00	24.75

VALIDATION REPORT

IdentModul 1.3-2018-04

Validated substance/substance group **Ylang-ylang oil**
Substance class APIs & excipients, liquid/semi-solid (other)
Report date 18/07/2018
Report number 31506-2018-07-18
Executing company HiperScan GmbH
Weißeritzstraße 3
01067 Dresden
Germany

Relevant substance names

Ylang-ylang oil; Oleum cananga odorata; Oleum ylang-ylang

Special notes

When selecting the *Ylang-ylang oil* substance/substance group, the following information is displayed to the user:

- Advice for the identification of essential oils : Some essential oils are listed in the class APIs & excipients, liquid/semi-solid (with analysis certificate) as well as in the class APIs & excipients, liquid/semi-solid (other).

Applicable documents

978-3-7692-5416-7 *European Pharmacopoeia 8th Edition, Basic Version 2014* [3]
Komm2.2.40 *Erfüllung von Ph. Eur. 2.2.40 durch Apo-Ident* [4]
AA004 *Erstellung und Validierung eines IdentModul-Updates*
Anhang F [Zusatz zu den Modellen der zweiten Stufe](#)

Validation method

Validation is performed after every change to the chemometric model (also “database”) in three steps:

1. The chemometric model is calculated from the calibration spectra using a *PCA algorithm*. The calibration spectra originate from the calibration samples of all substances in this class.
2. In the *chemometric model* generated, the distances between all separable substances are checked for compliance with the specified safety distances.
3. All spectra are presented to the generated chemometric model for evaluation. In three runs, the reference spectra (*Type A*), spectra from independent samples (*Type B*), and spectra from the field (*Type C*) are presented successively. Here, no single *false positive* result is permitted.

Finally, a report is generated from the validation run results and is archived together with the parameters of the model generation in audit-compliant manner.

Calibration samples

Only spectra which have been recorded by *HiperScan GmbH* from traceable samples are used for the generation of the [chemometric models](#). The following samples have been obtained from the substance/substance group *Ylang-ylang oil*:

Supplier	Substance	Batch	Sample ID	Spectra	Certificate
Taoasis	Ylang-ylang oil	140114-115612	31506	40	not required
Taoasis	Ylang-ylang oil	07122-119828	31653	40	not required
Taoasis	Ylang-ylang oil	41813-124965	32808	40	not required

Validation samples

A total of 44 773 spectra were provided for validation. The results were evaluated separately according to the following sample categories:

Type A All calibration spectra.

- 120 spectra of 3 reference samples from the substance/substance group *Ylang-ylang oil*. These samples are listed above in the section [calibration samples](#). The reference samples originate from 3 different batches.
- 18 629 spectra from a total of 436 batches from further 150 substances. These spectra were recorded by *HiperScan GmbH*. *Type A* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type A* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix A](#). The samples in [Appendix A](#) were added for model generation not with the intention to have the model identify these substances, but with the intention to make the model reject them despite their similarity to substances identifiable.

Type B Spectra recorded under the control of *HiperScan GmbH* and not used for the generation of the database.

Samples from batches, from which no spectra have been used for the generation of the database are considered as independent samples. The number of batches from which independent samples have delivered *Type B* spectra for the validation is given below and thus represents the number of independent *Type B* samples. Samples from which a number of the spectra has been used for the generation of the database and other spectra have been incorporated into the validation, are marked with a †. The following applies to the remaining unmarked samples: There was at least one other sample from the same batch (other sales containers, other sample ID), of which reference spectra (*Type A*) were used for the generation of the database.

- 120 spectra from 4 reference samples of the substance/substance group *Ylang-ylang oil*.
- These include spectra of independent samples from 2 batches, of which no spectra have been used for the generation of the database.

Supplier	Substance	Batch	Sample ID	Spectra
Taoasis	Ylang-ylang oil	140114-115612	31506 [†]	20
Taoasis	Ylang-ylang oil	07122-119828	31653 [†]	20
Taoasis	Ylang-ylang oil	43197-126348	33049	40
Taoasis	Ylang-ylang oil	B966015-128199	33648	40

- 22 711 spectra from a total of 545 batches from further 202 substances. These spectra were recorded by *HiperScan GmbH*. *Type B* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type B* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix B](#).

[†]These spectra were recorded from material of the same package as the corresponding calibration spectra (*Type A*). Hence, these spectra are not independent and are not of *Type B*.

Type C Spectra from the field, which were not recorded under the control of *Hiperscan GmbH* and have not been used for the generation of the database. *Hiperscan GmbH* accepts the details provided by the customer regarding the manufacturer and batch number mainly unchecked.

- 0 spectra from 0 *Apo-Ident* customers from 0 batches from the substance/substance group *Ylang-ylang oil*.
- These include spectra of independent samples from 0 batches, of which no spectra have been used for the generation of the database.
- 3193 spectra from 304 *Apo-Ident* customers from a total of 1659 batches from a further 122 substances. These spectra were recorded by *Apo-Ident* customers. *Type C* samples are documented within this entire report. If the substance can be identified with this [chemometric model](#), the samples are listed in the *Type C* section of each substance page. If the substance cannot be identified by this model, the samples are listed in [Appendix C](#).

Validation results

The validation runs checked whether the substance/substance group *Ylang-ylang oil* can clearly be distinguished from all other substances using NIR spectroscopy with *Apo-Ident*. For this purpose, all relevant spectra of the various substances were compared with *Ylang-ylang oil* and it was evaluated how many matches (positive) and rejections (negative) were correct or incorrect. The following table breaks down the numbers of correct and incorrect results according to the expected result (*positive/negative*) and the validation spectrum type (*A/B/C*).

	False positive	True positive	False negative	True negative
Type A	0	120	0	18 629
Type B	0	75	45	22 711
Type C	0	0	0	3193

The substance/substance group *Ylang-ylang oil* can be clearly distinguished from all other substances. In order to make these figures comparable, the weighted *true negative rate (specificity)* and the weighted *true positive rate (recognition rate)* are determined:

	Specificity	Recognition rate
Type A	100.0000 % (> 99.9536 %)	100.0000 % (> 95.0000 %)
Type B	100.0000 % (> 99.9530 %)	62.5000 % (> 60.0000 %)
Type C	100.0000 % (> 98.3567 %)	n/a (n/a)

In order to ensure that each substance is entered with the same weight irrespective of the number of spectra available, each spectrum is weighted with the reciprocal value of its number. (If several new spectra of a substance with a very distinctive spectrum are added, the values for *specificity* and *recognition rate* are safer, but are by no means idealised.)

In order to illustrate the impact of the number of spectra, a comparison is made in parentheses as to how the *specificity* or *recognition rate* would worsen if three additional incorrect results were to exist amongst the spectra (*Rule of Three* [10, 11]). The larger the number of spectra, the lower the deterioration if the three hypothetical *incorrect results* were added.

Where the number of spectra does not exceed 20, no *recognition rate* is provided.

Identity according to identity test using NIR spectroscopy

For each substance, a certificate of the correct identity is obtained from an independent GMP-certified test laboratory. If the identity of the sample can be provided using NIR on an independently tested reference sample, in the following table the *Mahalanobis distance* to the reference sample is specified as well as the *Mahalanobis distance* to the next non-identical substance:

Sample ID	Reference sample ID	Distance to reference sample	Distance to next foreign sample
31506	31506	0.00	92.89
31653	31653	0.00	40.17
32808	32808	0.00	81.71

Appendix A: Additional calibration samples (Type A)

Not required.

Appendix B: Additional validation samples (*Type B*)

It is necessary that spectra also enter the validation which cannot be identified with this model. In this manner, it is verified that the model also rejects unknown substances. The spectra for these samples were recorded by *HiperScan GmbH*. They are allocated to the *Type B*. They also include the calibration spectra for other models.

The samples originate from 101 batches. From these 4331 spectra were recorded. The spectra recorded on independent samples of substances which can be identified with the model are listed respectively in the section *Type B* for the individual substances and do not appear again elsewhere in this list.

Supplier	Substance	Batch	Spectra	Certificate
Actavis	Abitima [®] clinic ...	159997	40	not required
Actavis	Abitima [®] clinic ...	145054	40	not required
Actavis	Abitima [®] clinic ...	159997	40	not required
Allergika	Dermifant [®] kids ...	1301020	80	not required
Allergika	Dermifant [®] kids ...	1306020	40	not required
Allergika	Dermifant [®] kids ...	1705081	40	not required
Allergika	Dermifant [®] kids ...	1607080	40	not required
Apotheker Bau...	Common wormwood ...	10815310556	40	not required
ASAV Apotheke...	Common wormwood ...	10116310555	60	not required
ASAV Apotheke...	Common wormwood ...	10815310556	40	not required
ASAV Apotheke...	Common wormwood ...	10815310556	40	not required
ASAV Apotheke...	Common wormwood ...	10815310556	40	not required
Audor	Lipoid S 100	zu erfassen	40	not required
Bayer	Ultrabas [®]	33851A	40	not required
Bayer	Ultrapril [®]	22100A	40	not required
Bayer	Ultrasicc [®]	33614A	40	not required
Bene	Thrombocid [®] gel	403I143	40	not required
Bene	Thrombocid [®] gel	401B101	40	not required
Bene	Thrombocid [®] gel	403I143	40	not required
Bene	Thrombocid [®] gel	401B101	40	not required
Bene	Thrombocid [®] gel	403I143	40	not required
Caelo	Base cream, dilu...	14242803	60	not required
Caelo	Milking grease S...	162049	40	20160711*
Caelo	Pine oil rubbing...	162091	40	20160713*
Caelo	Salmon oil	170345	40	20170213*
Caelo	Ultrasound conta...	12031301	40	not required
Caelo	Vario base cream	12077902	50	1405329
Caelo	Vario base cream	13257102	40	1405330
Dr. Wolff	Anefug [®] Simplex	237590	40	1407216
Dr. Wolff	Anefug [®] Simplex	234280	40	not required
Dr. Wolff	Linola [®] acute 0.5%	334620	40	not required
Dr. Wolff	Linola [®] sun prot...	134530	40	not required
Dyckerhoff	Regeneresen	150032Y-Na-5	40	not required
Dyckerhoff	Regeneresen	211111Rk-2	40	not required
Dyckerhoff	Regeneresen	061211Art-2	40	not required
Dyckerhoff	Regeneresen	zu erfassen	40	not required
Einhorn-Apoth...	Rizol raw material	AL0413-1	60	not required
Einhorn-Apoth...	Rizol raw material	200812174102290W	30	not required
Euro OTC	Capsaicin liquid...	2908036-01	41	not required
Fagron	Fitalite TM	1503259	40	not required
Fagron	Fitalite TM	1503259	40	not required
Fagron	Fitalite TM	14B18-T01-010352	60	not required
Fagron	Fitalite TM	1503259	40	not required
Fagron	Fitalite TM	14B18-T01-010352	60	not required
Fagron	Ora-Plus [®]	3334081	60	not required
Fagron	Ora-Sweet [®]	3062252	40	not required
Fagron	Versatile TM	1503256	40	not required

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra	Certificate
Fagron	Versatile TM	14B19-T02-010391	60	not required
Fagron	Versatile TM	1503256	40	not required
Fagron	Versatile TM	14B19-T02-010391	60	not required
Fagron	Versatile TM	1503256	40	not required
Fagron	Versatile TM Rich	1503257	40	not required
Fagron	Versatile TM Rich	14B19-T08-010395	60	not required
Fagron	Versatile TM Rich	14B19-T04-010395	60	not required
Fagron	Versatile TM Rich	1503257	40	not required
Fagron	Versatile TM Rich	1503257	40	not required
Ichthyol	Milk Cordes [®] O/W...	10D032	40	not required
Koko	dermaviduals [®] ba...	L003L13	60	not required
Paddock	Ora-Blend [®]	2280130	40	not required
Paddock	Ora-Blend [®] SF	2038178	40	not required
Perrigo	Ora-Blend [®] SF	3112578	40	not required
Perrigo	Ora-Plus [®]	3334081	40	not required
Perrigo	Ora-Sweet [®]	3062252	40	not required
Pharmapol	Glycerol trinitr...	228718	40	not required
Pharmapol	Glycerol trinitr...	211528	50	not required
Pharmapol	Glycerol trinitr...	232961	60	not required
Phylak Sachse...	Skin cream, high...	zu erfassen	40	not required
Phylak Sachse...	Skin cream, ligh...	zu erfassen	40	not required
Primavera	Blood orange oil	00413G25	40	not required
Primavera	Cardamom oil	00080M25	40	not required
Primavera	Cardamom oil	00897G25	40	not required
Primavera	Melissa oil	00720H25	40	not required
Primavera	Melissa oil 30%	01029J25	40	not required
Primavera	Melissa oil 30%	00344A26	40	not required
Primavera	Rose oil 10%	00721716	40	not required
Primavera	Rosemary camphor...	00559B26	40	not required
Primavera	Rosemary camphor...	00536K25	40	not required
Primavera	Tolu Resinoid	00115A25	40	not required
Primavera	Wild rose oil	94930-95763	40	not required
Salderman Gmb...	Salderman base e...	180123	40	not required
Spirig Pharma	Kerasal [®] base oi...	N011	40	not required
Spirig Pharma	Kerasal [®] base oi...	M031	40	not required
Stiefel	Physiogel [®] hypoa...	6940	40	not required
Stiefel	Physiogel [®] hypoa...	343P	40	not required
Stoko	Praecutan [®] lotio...	011657101571	40	not required
Stoko	Praecutan [®] lotio...	0915777806	40	not required
Taoasis	Angelica root ex...	21439-67783	40	not required
Taoasis	Black cumin oil	504511-115611	40	not required
Taoasis	Blood orange oil	494J09	40	not required
Taoasis	Common wormwood ...	7.9363	40	not required
Taoasis	Melissa oil	640	40	not required
Taoasis	Pine silvestris ...	1790-125357BAG90451	40	not required
Taoasis	Pine silvestris ...	1790-129074	40	not required
Taoasis	Rose oil, pure	12014-120330H08	40	not required
Taoasis	Savory herb	1569-129318	40	not required
Taoasis	Savory herb	498	40	not required
Taoasis	Thuja oil	28508-121881	40	not required
Taoasis	Thuja oil	66255-969	40	not required
Taoasis	Tolu res. 50% oil	zu erfassen	40	not required
Taoasis	Tolu res. 50% oil	493887-105383BAC90451	40	not required
Taoasis	Vanilla extract ...	789171-114788	40	not required

*The spectra were taken from a part of the same sample which the supplier used for its analysis (identity, purity, content). In this case the manufacturer certificates proof the identity and quality of the sample from those the spectra descend.

Appendix C: Additional validation samples (*Type C*)

Spectra for substances from the field which cannot be identified with this model also enter the validation. In this manner, it is verified that the model also rejects unknown substances. The spectra for these samples were recorded by *Apo-Ident* customers. They belong to *Type C*. The information provided by the customer regarding the manufacturer and batch number is taken over by *HiperScan GmbH* to a large extent unchecked.

The samples originate from 94 batches. From these, 162 spectra were recorded. The spectra recorded on independent samples of substances from the field which can be identified with the model are listed respectively in the section *Type C* for the individual substances and do not appear again elsewhere in this list.

Supplier	Substance	Batch	Spectra
Allergika/Kehr	Dermifant [®] kids lotion	1604060	1
Anzag/Galderma	Kerasal [®] base ointment	S075	1
Arkeo Immun UG	Rizol raw material	1192HJ	2
Bombastus	Vario base cream	12375801	1
Caelo	Common wormwood oil	79363	2
Caelo	Milk Cordes [®] O/W Emul...	13D002	1
Caelo	Milk Cordes [®] O/W Emul...	13D001	2
Caelo	Rizol raw material	15144304	1
Caelo	Vario base cream	16032402	1
Caelo	Vario base cream	14331902	4
Caelo	Vario base cream	13257102	2
Caelo	Vario base cream	13143401	1
Caelo	Vario base cream	12077902	1
Dr. Wolff	Anefug [®] Simplex	136840	1
Dyckerhoff	Regeneresen	120313Kn-2	1
Dyckerhoff	Regeneresen	60611dü-2	1
Dyckerhoff	Regeneresen	120023gh-2	1
Dyckerhoff	Regeneresen	120053NNR-2	1
Dyckerhoff	Regeneresen	120054Lu-2	1
Dyckerhoff	Regeneresen	120063ytfm-2	1
Dyckerhoff	Regeneresen	120073sy-2	1
Dyckerhoff	Regeneresen	120243p1-2	1
Dyckerhoff	Regeneresen	1202920v-2	1
Dyckerhoff	Regeneresen	120303Dü-2	1
Dyckerhoff	Regeneresen	120323Lu-2	1
Dyckerhoff	Regeneresen	120333NN-2	1
Dyckerhoff	Regeneresen	120502Ni-2	1
Dyckerhoff	Regeneresen	120453P1-2	1
Dyckerhoff	Regeneresen	250511Ad-2	1
Dyckerhoff	Regeneresen	1204620Le-2	1
Dyckerhoff	Regeneresen	120463GH-2	1
Dyckerhoff	Regeneresen	150423Ni-2	1
Dyckerhoff	Regeneresen	130111AU-2	1
Dyckerhoff	Regeneresen	150173th-2	1
Dyckerhoff	Regeneresen	150403HZ-2	1
Dyckerhoff	Regeneresen	1503839-Na	1
Euro OTC	Capsaicin liquid extr...	6061204	1
Euro OTC	Capsaicin liquid extr...	1201010-01	1
Euro OTC	Capsaicin liquid extr...	12111205	1
Euro OTC	Capsaicin liquid extr...	1205038-01	2
Euro OTC	Capsaicin liquid extr...	1205038-02	2
Fagron	Ora-Blend [®] SF	4136370	1
Fagron	Ora-Blend [®] SF	3745164	1
Fagron	Ora-Sweet [®]	3414679	1
Fagron	Versatile [™]	14B19-T02-010391	1
Galderma/Gehe	Kerasal [®] base ointment	S061	1

continued on the next page

continued from previous page

Supplier	Substance	Batch	Spectra
Galderma/Anzag	Kerasal [®] base ointment	S041	4
Gehe	Glycerol trinitrate 5%	214895	1
Gehe	Ora-Blend [®] SF	3112578	1
Gehe	Ora-Blend [®] SF	2038178	1
GSK Stiefel	Physiogel [®] hypoallerg...	859R	1
GSK/Stiefel	Physiogel [®] hypoallerg...	538T	1
Ichthyol	Milk Cordes [®] O/W Emul...	60613	1
Ichthyol - Gesellschaft	Milk Cordes [®] O/W Emul...	11D037	1
Ichthyol / Noweda	Milk Cordes [®] O/W Emul...	12D008	6
IchthyolGmbH/Anzag	Milk Cordes [®] O/W Emul...	12D003	2
Ichtyol, Noweda	Milk Cordes [®] O/W Emul...	13d007	1
Perrigo	Ora-Blend [®] SF	2088609	1
Perrigo	Ora-Sweet [®]	4247427	2
Pharmapohl	Glycerol trinitrate 5%	18102013A	1
Pharmapol	Glycerol trinitrate 5%	232961	3
Pharmapol	Glycerol trinitrate 5%	234783	2
Pharmapol	Glycerol trinitrate 5%	AKD81	1
Pharmapol	Glycerol trinitrate 5%	221414	4
Pharmapol	Glycerol trinitrate 5%	224310(ex)224658(in)	2
Pharmapol	Glycerol trinitrate 5%	25041301	1
Pharmapol Arzneimitte...	Glycerol trinitrate 5%	219108	4
Pharmapol Arzneimitte...	Glycerol trinitrate 5%	232961	7
Pharmapol/Direkt	Glycerol trinitrate 5%	224310	7
Pharmapol/Direkt	Glycerol trinitrate 5%	228718	13
Pharmapol/Direkt	Glycerol trinitrate 5%	226578	12
Pharmapol/Zentrum Apo	Glycerol trinitrate 5%	2913A-07002	1
PHOENIX 21.10.2013	Physiogel [®] hypoallerg...	1142N	1
Phoenix/Stiefel	Physiogel [®] hypoallerg...	399P	1
Phönix	Ora-Blend [®] SF	3475164	1
Phönix	Ora-Blend [®] SF	3434856	2
phönix, 09.11.2012	Capsaicin liquid extr...	120503801	2
Spirig Pharma	Kerasal [®] base ointment	N021	1
Spirig/Gehe	Kerasal [®] base ointment	S076	1
Spirig/Noweda	Kerasal [®] base ointment	N032	2
Spirig/Sanacorp	Kerasal [®] base ointment	N011	1
Steidl/Phytolab	Rizol raw material	200812174102290	1
Stiefel	Physiogel [®] hypoallerg...	926Q	1
Stiefel	Physiogel [®] hypoallerg...	270T	1
Stiefel	Physiogel [®] hypoallerg...	537T	1
Stiefel/Noweda	Physiogel [®] hypoallerg...	163R	1
Stiefel/Sanacorp	Physiogel [®] hypoallerg...	10000	1
Taoasis	Tolu res. 50% oil	546022-122636	1
Taoasis	Tolu res. 50% oil	546022-114802	2
Taoasis	Tolu res. 50% oil	23772-119167	1
Taoasis	Tolu res. 50% oil	23772-118646	1
Taoasis	Tolu res. 50% oil	493887-105384	1
Taoasis	Wild rose oil	106177-106175	1
Taoasis	Wild rose oil	99149-99474	1

Appendix D: Requirements of validation

In order to ensure adherence to the safe scientific status, the individual methods for manufacturing and testing must be validated under certain circumstances (compare § 34 para. 1 no. 3, § 35 para. 1 no. 4 and para. 4 sentence 1 no. 2 b, para. 6 sentence 3 *ApoBetrO* [Pharmacies Rules and Regulations]). The *ApoBetrO* [Pharmacies Rules and Regulations] incorporates a legal definition in § 1 a para. 16 (quotation translated):

“Validation is the provision of documented proof which with a high degree of safety documents that, via a specific process or standard work process, a medicinal product is manufactured and tested, which is in accordance with previously determined quality features.”

Validation documentation can be used to prove that methods or devices which are not described in the Pharmacopoeia within the meaning of § 6 para. 1 sentence 3 *ApBetrO* [Pharmacies Rules and Regulations] achieve the same results as those in the Pharmacopoeia. On the other hand, with the requirements of the demanded validation it must be observed whether the respective testing method is already incorporated in the Pharmacopoeia.

NIR spectroscopy as a general testing method need not be validated in accordance with the express ruling in the *Ph. Eur. Section 1.1* [3], as it is already described in *Section 2.2.40* of the *Ph. Eur.* as an area of application for the identification of raw materials.

However, a special validation requirement exists for the reference database. This requirement is met with the existing document. Further requirements or rules as to how this proof must be furnished do not exist. It is required that the processes guarantee the same results as the methods and devices in the Pharmacopoeia [17].

Carrying out identity tests with *Apo-Ident* is therefore also possible if the NIR spectroscopy process is not required in the Pharmacopoeia monograph of the substance for identity testing. All NIR analyses with *Apo-Ident* prove several, often all molecule groups and are therefore comparable with a series of individual, targeted chemical proofs [4]. Therefore, the identity proof with *Apo-Ident* replaces the monograph test series (with two or more test combinations).




Figure 2: The combination of tests of the monograph is replaced by the alternative method NIR spectroscopy using *Apo-Ident*. This is permissible because both test procedures result in the establishment of the identity of the raw material.

With this validation documentation, proof is furnished that identical results are achieved with *Apo-Ident* and Pharmacopoeia methods, i.e. confirmation of the identity of the raw material [2].

Appendix E: Conformity of Apo-Ident with the European pharmacopeia

According to *Ph. Eur. Section 2.2.40*, NIR spectroscopy is basically suitable for: “Identification of agents, excipients, dosage forms, intermediate manufacturing products, chemical raw materials and packaging materials” ([3], quotation translated).

The fact that *Apo-Ident* meets the further criteria of the European Pharmacopoeia under the headings in *Section 2.2.40*

- Apparatus
- Measurement methods
- Sample preparation and presentation
- Testing the functionality of the instrument
- Identification and characterisation (qualitative analysis)
- Quantitative analysis
- Ongoing model evaluation
- Transfer of databases
- Data storage

can be proven based on the *HiperScan GmbH* documentation of “Meeting *2.2.40 Ph. Eur.* by *Apo-Ident*” [4].

Appendix F: Addition to the two-stage models

As described in the general part of the *Apo-Ident* validation documentation, in the section [Model creation procedure and validation runs](#), under 2. *Generating the chemometric models (calibration) in f)*, it is possible to create a further chemometric model for certain substances (second-stage model) and to undertake the assessment in several steps.

Substances in stage-two models

In the *Apo-Ident Update 2018-04*, a distinction is made between the following substances in the listed stage-two models:

Submodel 1

Bacitracin
Betamethasone, micronized
Betamethasone valerate
Budesonide, micronized
Capsaicin, natural
Dexamethasone
Erythromycin
Hydrocortisone butyrate
Norethisterone acetate
Prednicarbate, micronized
Prednisolone, micronized
Triamcinolone acetonide

Submodel 2

Beclometasone dipropionate, anhydrous
Betamethasone dipropionate, micronized
Clobetasol propionate
Diphenylcyclopropenone
Gentamicin sulfate
Sodium benzoate
Sodium citrate
Oxybutynin hydrochloride

Submodel 3

Alfatradiol
Estriol
Pregnenolone
Progesterone, micronized
Quinine hydrochloride
Quinine sulfate dihydrate
Spironolactone
Testosterone propionate

Liquid submodel

2-ethylhexyl laurate
Base cream DAC
Lanette® ointment (conserved)
Non-ionic hydrophil ointment

Squalane

Wool alcohol cream DAB/SR

References

- [1] ABDA – BUNDESVEREINIGUNG DEUTSCHER APOTHEKERVERBÄNDE: Verordnung über den Betrieb von Apotheken (Apothekenbetriebsordnung – ApBetrO), 2012
- [2] REIMANN, B. ; REGIERUNGSPRÄSIDIUM DARMSTADT: Hinweise zur ordnungsgemäßen Prüfung von Arzneimitteln und Ausgangsstoffen (§§ 6 und 11 *ApBetrO*), 2007
- [3] *Europäisches Arzneibuch, Grundwerk 2014 einschließlich 1. bis 4. Nachtrag.* 8. Ausgabe. Deutscher Apotheker Verlag (978-3-7692-6508-8)
- [4] HIPERSCAN GMBH: Erfüllung von 2.2.40 Ph. Eur. durch Apo-Ident, 2013
- [5] ARBEITSGEMEINSCHAFT DER PHARMAZIERÄTE DEUTSCHLANDS (APD): Resolution in Verbindung mit §§ 6 und 11 *ApBetrO* Verwendung eines Nah-Infrarot-Spektrometers (NIR) zur Identitätsprüfung, 16. 10. 2013, DAZ 48, November 2013
- [6] ARBEITSGEMEINSCHAFT DER PHARMAZIERÄTE DEUTSCHLANDS (APD): Resolution 2014, Arbeitsgemeinschaft der Pharmazieräte Deutschlands (APD), Oktober 2014
- [7] *DAC/NRF.* Govi-Verlag (978-3-7741-0044-2)
- [8] KESSLER, W.: *Multivariate Datenanalyse.* WILEY-VCH Verlag, 2007 (978-3-527-31262-7)
- [9] NÆS, T. ; ISAKSSON, T. ; FEARN, T. ; DAVIES, T.: *Multivariate Calibration and Classification.* NIR Publications, 2002 (978 0 9528666 2 6)
- [10] HANLEY, J. A. ; LIPPMAN-HAND, A.: If nothing goes wrong, is everything all right? In: *Journal of the American Medical Association* 249 (1983), S. 1743–1745
- [11] JOVANOVIĆ, B. D. ; LEVY, P. S.: A Look at the Rule of Three. In: *The American Statistician* 51 (1997), S. 137–139
- [12] BRONSTEIN, I. N. ; SEMENDJAJEW, K. A. ; MUSIOL, G. ; MÜHLIG, H.: *Taschenbuch der Mathematik.* 5. überarbeitete und erweiterte Auflage. Verlag Harri Deutsch, 2000 (3-8171-2015-2)
- [13] MAHALANOBIS, P.: On the generalized distance in statistics. In: *Proc. Nat. Inst. Sci. India (Calcutta)* 2 (1936), S. 49–55
- [14] YAMBOR, B. ; DRAPER, W. ; BEVERIDGE, R.: Analyzing PCA-based face recognition algorithms: Eigenvector selection and distance measures. In: *Second Workshop Empirical Evaluation in Computer Vision* (2000)
- [15] HIPERSCAN GMBH: Identifikationsmethodik Apo-Ident, 2012
- [16] HIPERSCAN GMBH: Datenvorbehandlung des Identifikationssystems Apo-Ident, 2012
- [17] CYRAN ; RÖTTA: Apothekenbetriebsordnung, Kommentar § 6 Rn. 10, 2010

Index

- Abitima® clinic body cream, *see* [Hydrophilic ointments](#)
- Abitima® clinic face cream, *see* [Hydrophilic ointments](#)
- Acidum dichloroaceticum, *see* [Dichloroacetic acid](#)
- Acidum salicylicum cum vaselino albo 10 %, *see* [Salicylic white paraffin 10%](#)
- Alfason Basis Cresa®, 21
- Alfason® Repair, *see* [Linola® fat creme / Linola® H fat N / Alfason® Repair](#)
- Allergika evening primrose oil cream 20%, *see* [Allergika evening primrose oil cream 20% / Dermifant® kids cream](#)
- Allergika evening primrose oil cream 20% / Dermifant® kids cream, 25
- Allergika® Base cream, *see* [Hydrophilic ointments](#)
- Aloe vera gel, 10x concentrated, *see* [Water and aqueous solutions](#)
- Aloe Vera gel, 1:1, *see* [Water and aqueous solutions](#)
- alpha-bisabolol (racemic) at least 85%, 29
- Amciderm® Base cream, *see* [Liniment ointments](#)
- Angelica archangelica oleum, *see* [Angelica root oil](#)
- Angelica root oil, 33
- Anise oil, organic, 37
- Aqua, *see* [Water and aqueous solutions](#)
- Aqueous ointments, 41
- Argan oil, 47
- Asche Basis® cream, *see* [Hydrophilic ointments](#)
- Asche Basis® fat ointment, *see* [Fat ointments](#)
- Asche Basis® lotion, *see* [Hydrophilic ointments](#)
- Asche Basis® ointment, *see* [Ointment ash basis / Neribas®](#)
- Balm Bio Nature, 51
- Banana aroma, 55
- Banana flavour, *see* [Banana aroma](#)
- Base cream Taoasis, *see* [Hydrophilic ointments](#)
- Basil oil, 59
- Benzoin siam oil 20%, 63
- Bepanthen® solution, *see* [Water and aqueous solutions](#)
- Bepanthen® Wund- und Heilsalbe, 67
- Bergamot oil, 71
- Cajeput, *see* [Cajeput / Niaouli oil](#)
- Cajeput / Niaouli oil, 75
- Camphor oil, 79
- Carrot seed oil, 83
- Caryophylli floris aetheroleum, *see* [Clove oil](#)
- Cedar wood oil, 87
- Chamomile oil, *see* [Matricaria chamomilla oil](#)
- Cinnamon bark oil, Ceylon, 91
- Citric oil, 95
- Citronella oil, 99
- Citrus sinensis oleum, *see* [Grapefruit oil organic / Orange oil](#)
- Citrus oil, *see* [Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil](#)
- Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil, 103
- Clary sage oil, *see* [Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil](#)
- Clove oil, 109
- Contramarum aroma, 113
- Contramarum flavour, *see* [Contramarum aroma](#)
- Coriander oil, *see* [Thyme oil, white / Coriander oil](#)
- Cottonwood ointment, *see* [Fat ointments](#)
- Cumin oil, 117
- Cymbopogonis aetheroleum, *see* [Lemon grass oil](#)
- Cypress oil, 121
- Dermatest base ointment, *see* [Aqueous ointments](#)
- Dermatop® base cream, *see* [Hydrophilic ointments](#)
- Dermatop® base ointment, 125
- Dermifant® kids cream, *see* [Allergika evening primrose oil cream 20% / Dermifant® kids cream](#)
- Dexeryl®, 129
- Dichloroacetic acid, 133
- Dimeticone ointment 10% SR, *see* [Dimeticone ointment 10% SR / Optiderm® lotion](#)
- Dimeticone ointment 10% SR / Optiderm® lotion, 137
- DMS® base cream classic, *see* [Hydrophilic ointments](#)
- DMS® base cream high classic, *see* [Hydrophilic ointments](#)
- DMS® base cream high classic plus, *see* [Hydrophilic ointments](#)
- Dwarf pine oil, 141
- Ethanol 70% pure / denatured, 145
- Ethanol 70% rein / vergällt, *see* [Ethanol 70% pure / denatured](#)

- Eucalypti aetheroleum, *see* [Eucalyptus oil](#) / [Rosemary oil](#)
- Eucalyptus oil, *see* [Eucalyptus oil](#) / [Rosemary oil](#)
- Eucalyptus oil / Rosemary oil, 153
- Eucerinum O/W basis, *see* [Hydrophilic ointments](#)
- Eucerinum W/O basis, *see* [Hydrophilic ointments](#)
- Excipial® almond oil ointment, 157
- Excipial® hydro cream, *see* [Hydrophilic ointments](#)
- Excipial® lipo cream, *see* [Aqueous ointments](#)
- Excipial® U Hydrolotio, *see* [Liniment ointments](#)
- Excipial® U Lipolotio, *see* [Hydrophilic ointments](#)
- Excipial® U10 Lipolotio, *see* [Hydrophilic ointments](#)
- Fabitol® base cream, *see* [Aqueous ointments](#)
- Fat ointments, 161
- Fenistil® Gel, 169
- Fennel oil, 173
- Frankincense oil, *see* [Spruce needle oil](#) / [Frankincense oil](#) / [Silver fir oil](#)
- German chamomile oil, *see* [Matricaria chamomilla oil](#)
- Ginger oil, 177
- Grapefruit oil organic / Orange oil, 181
- Grapefruit oil, organic, *see* [Grapefruit oil organic](#) / [Orange oil](#)
- Hans Karrer Hydrocream MicroSilver, *see* [Sebexol® formula basis](#) / [Sebexol® cream lotion](#) / [Wofacutan washing lotion](#) / [Hans Karrer Hydrocream MicroSilver](#)
- Hans Karrer Lipocream MicroSilver, *see* [Hydrophilic ointments](#)
- Hans Karrer Lipolotion MicroSilver, *see* [Hydrophilic ointments](#)
- Hydrophilic ointments, 185
- Hyssop oil, *see* [Citrus oil](#) / [Lavender oil](#) / [Clary sage oil](#) / [Petitgrain oil](#) / [Hyssop oil](#)
- Immortelle oil, 201
- Juniper oil, 205
- Juniperi aetheroleum, *see* [Juniper oil](#)
- La Roche-Posay Cold Cream Naturel, 209
- La Roche-Posay Toleriane, 213
- Lavender oil, *see* [Citrus oil](#) / [Lavender oil](#) / [Clary sage oil](#) / [Petitgrain oil](#) / [Hyssop oil](#)
- Lavendula officinalis oleum, *see* [Citrus oil](#) / [Lavender oil](#) / [Clary sage oil](#) / [Petitgrain oil](#) / [Hyssop oil](#)
- Lemon grass oil, 217
- Lime oil, 221
- Limonis aetheroleum, *see* [Citric oil](#)
- Liniment ointments, 225
- Linola® body milk, *see* [Hydrophilic ointments](#)
- Linola® fat cream, *see* [Linola® fat creme](#) / [Linola® H fat N](#) / [Alfason® Repair](#)
- Linola® fat creme / Linola® H fat N / Alfason® Repair, 231
- Linola® H fat N, *see* [Linola® fat creme](#) / [Linola® H fat N](#) / [Alfason® Repair](#)
- Linola® O/W cream, *see* [Liniment ointments](#)
- Linola® Sept, *see* [Liniment ointments](#)
- Lipoderm® lotion, *see* [Hydrophilic ointments](#)
- Lipolotio urea 5% F body lotion, 235
- Lygal® head ointment N 3%, 239
- Lygal® ointment base, 243
- Majoranae aetheroleum, *see* [Marjoram oil](#)
- Mandarin oil, green, 247
- Manuka oil, 251
- Marjoram oil, 255
- Marjoram ointment, *see* [Fat ointments](#)
- Matricaria chamomilla oil, 259
- Matricaria oil, morrocan, 263
- Matricaria oil, roman, 267
- Melaleuca alternifolia oleum, *see* [Tea tree oil](#)
- Menthae piperitae aetheroleum, *see* [Peppermint oil](#)
- Myrrh oil, 271
- Myrtle oil, 275
- Neribas® cream, *see* [Hydrophilic ointments](#)
- Neribas® fat ointment, *see* [Fat ointments](#)
- Neribas® ointment, *see* [Ointment ash basis](#) / [Neribas®](#)
- Neroli oil, 279
- Neuroderm® care lotion, *see* [Neuroderm® care lotion](#) / [moisturising cream](#)
- Neuroderm® care lotion / moisturising cream, 283
- Neuroderm® moisturising cream, *see* [Neuroderm® care lotion](#) / [moisturising cream](#)
- Neuroderm® moisturising cream lipo, 287
- Niaouli oil, *see* [Cajeput](#) / [Niaouli oil](#)
- Nicotine, 291
- Nourivan™ Antiox, *see* [Liniment ointments](#)
- Ocimum basilicum oleum, *see* [Basil oil](#)
- Ointment ash basis / Neribas®, 295
- Oleum abies alba, *see* [Spruce needle oil](#) / [Frankincense oil](#) / [Silver fir oil](#)

- Oleum angelicae (e radice), *see* [Angelica root oil](#)
- Oleum aniba roseodora, *see* [Rosewood oil](#)
- Oleum anthemis nobilis, *see* [Matricaria oil, roman](#)
- Oleum Arganiae, *see* [Argan oil](#)
- Oleum bergamottae, *see* [Bergamot oil](#)
- Oleum boswellia serrata, *see* [Spruce needle oil / Frankincense oil / Silvire fir oil](#)
- Oleum Cajeputi artificiale, *see* [Cajeput / Niaouli oil](#)
- Oleum cananga odorata, *see* [Ylang-ylang oil](#)
- Oleum caryophylli, *see* [Clove oil](#)
- Oleum cedrus deodara, *see* [Cedar wood oil](#)
- Oleum cinnamomi ceylanici, *see* [Cinnamon bark oil, Ceylon](#)
- Oleum citri, *see* [Citric oil](#)
- Oleum citrus aurantifolia, *see* [Lime oil](#)
- Oleum citrus aurantium, *see* [Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil](#)
- Oleum citrus aurantium var. amara, *see* [Neroli oil](#)
- Oleum citrus bergamia, *see* [Bergamot oil](#)
- Oleum citrus paradisi, *see* [Grapefruit oil organic / Orange oil](#)
- Oleum citrus sinensis, *see* [Grapefruit oil organic / Orange oil](#)
- Oleum coriandri, *see* [Thyme oil, white / Coriander oil](#)
- Oleum cuminum cyminum, *see* [Cumin oil](#)
- Oleum cupressus sempervirens, *see* [Cypress oil](#)
- Oleum cymbopogon martinii, *see* [Palmarosa oil, organic](#)
- Oleum cymbopogon winterianus, *see* [Citronella oil](#)
- Oleum daucus carota, *see* [Carrot seed oil](#)
- Oleum eucalypti, *see* [Eucalyptus oil / Rosemary oil](#)
- Oleum foeniculum vulgare var. dulce, *see* [Fennel oil](#)
- Oleum gaultheria procumbens, *see* [Wintergreen oil](#)
- Oleum helichrysum italicum, *see* [Immortelle oil](#)
- Oleum hyssopus officinalis, *see* [Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil](#)
- Oleum juniperi e baccarum, *see* [Juniper oil](#)
- Oleum lavendula officinalis, *see* [Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil](#)
- Oleum leptospermum scoparium, *see* [Manuka oil](#)
- Oleum Majoranae, *see* [Marjoram oil](#)
- Oleum melaleuca alternifolia, *see* [Tea tree oil](#)
- Oleum melaleuca viridiflora, *see* [Cajeput / Niaouli oil](#)
- Oleum mentha spicata, *see* [Spearment oil](#)
- Oleum menthae piperitae dopp. rectific., *see* [Peppermint oil](#)
- Oleum myrrha, *see* [Myrrh oil](#)
- Oleum neroli, *see* [Neroli oil](#)
- Oleum niaouli, *see* [Cajeput / Niaouli oil](#)
- Oleum ocimum basilicum, *see* [Basil oil](#)
- Oleum olibanum, *see* [Spruce needle oil / Frankincense oil / Silvire fir oil](#)
- Oleum origani cretici, *see* [Oregano oil](#)
- Oleum ormenis multicaulis, *see* [Matricaria oil, morrocan](#)
- Oleum palmarosae, *see* [Palmarosa oil, organic](#)
- Oleum pelargonium graveolens, *see* [Rose-geranium oil, organic](#)
- Oleum pini pumilionis, *see* [Dwarf pine oil](#)
- Oleum pini sibiricum, *see* [Spruce needle oil / Frankincense oil / Silvire fir oil](#)
- Oleum pinus cembra, *see* [Swiss pine oil](#)
- Oleum pogostemon cablin, *see* [Patchouli oil](#)
- Oleum rosmarini, *see* [Eucalyptus oil / Rosemary oil](#)
- Oleum salviae, *see* [Sage oil](#)
- Oleum salviae sclarea, *see* [Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil](#)
- Oleum styrax tonkinensis, *see* [Benzoin siam oil 20%](#)
- Oleum thymus vulgaris, *see* [Thyme oil, red](#)
- Oleum vanillae planifoliae, *see* [Perfume oil vanilla](#)
- Oleum Vaselini album, *see* [White paraffin oil](#)
- Oleum vetiveria zizanioides, *see* [Vetiver bourbon oil](#)
- Oleum ylang-ylang, *see* [Ylang-ylang oil](#)
- Oleum zingiber officinalis, *see* [Ginger oil](#)
- Optiderm® lotion, *see* [Dimeticone ointment 10% SR / Optiderm® lotion](#)
- Orange aroma, [301](#)
- Orange flavour, *see* [Orange aroma](#)
- Orange oil, *see* [Grapefruit oil organic / Orange oil](#)
- Oregano oil, [305](#)
- Ormenis multicaulis oleum, *see* [Matricaria oil, morrocan](#)
- Palmarosa oil, organic, [309](#)
- Panthenol ointment Lichtenstein, [313](#)
- Patchouli oil, [317](#)
- Peppermint oil, [321](#)
- Perfume oil vanilla, [325](#)
- Petitgrain oil, *see* [Citrus oil / Lavender oil / Clary sage oil / Petitgrain oil / Hyssop oil](#)

- Piceae aetheroleum, *see* [Spruce needle oil / Frankincense oil / Silvire fir oil](#)
- Pini pumilionis aetherolum, *see* [Dwarf pine oil](#)
- Protegin® XN, *see* [Fat ointments](#)
- Raspberry aroma, [329](#)
- Raspberry flavour, *see* [Raspberry aroma](#)
- Ravensara, [333](#)
- Retterspitz external, *see* [Water and aqueous solutions](#)
- Rose-geranium oil, organic, [337](#)
- Rosemary oil, *see* [Eucalyptus oil / Rosemary oil](#)
- Rosewood oil, [341](#)
- Rosmarini aetheroleum, *see* [Eucalyptus oil / Rosemary oil](#)
- Sage oil, [345](#)
- Salicylic white paraffin 10%, [349](#)
- Salvia officinalis oleum, *see* [Sage oil](#)
- Sandalwood oil, [353](#)
- Sebexol® basic pH 5 formula basis, *see* [Sebexol® formula basis / Sebexol® cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver](#)
- Sebexol® cream lotion pH 5, *see* [Sebexol® formula basis / Sebexol® cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver](#)
- Sebexol® formula basis / Sebexol® cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver, [357](#)
- Siam benzoin, [365](#)
- Silvire fir oil, *see* [Spruce needle oil / Frankincense oil / Silvire fir oil](#)
- Solutio Cordes®, [369](#)
- Spearmint oil, [373](#)
- Spruce needle oil, *see* [Spruce needle oil / Frankincense oil / Silvire fir oil](#)
- Spruce needle oil / Frankincense oil / Silvire fir oil, [377](#)
- Stomahesive® adhesive paste, [383](#)
- Strawberry aroma, [387](#)
- Strawberry flavour, *see* [Strawberry aroma](#)
- Swiss pine oil, [391](#)
- SyrSpend® SF pH4 cherry aroma, *see* [Water and aqueous solutions](#)
- SyrSpend® SF pH4 aroma free, *see* [Water and aqueous solutions](#)
- Tea tree oil, [395](#)
- Thrombocid® ointment, *see* [Hydrophilic ointments](#)
- Thyme oil, red, [399](#)
- Thyme oil, white, *see* [Thyme oil, white / Coriander oil](#)
- Thyme oil, white / Coriander oil, [403](#)
- Unguentum cereum, *see* [Wax ointment \(stabilised\)](#)
- Unguentum dimeticoni 10% SR, *see* [Dimeticone ointment 10% SR / Optiderm® lotion](#)
- Unguentum majoranae, *see* [Fat ointments](#)
- Unguentum Populi, *see* [Fat ointments](#)
- Vetiver bourbon oil, [407](#)
- Wart ointment InfectoPharm® (NRF 11.31), [411](#)
- Water, *see* [Water and aqueous solutions](#)
- Water and aqueous solutions, [415](#)
- Wax ointment (stabilised), [421](#)
- White paraffin oil, [427](#)
- Wintergreen oil, [431](#)
- Wofacutan wash lotion, *see* [Sebexol® formula basis / Sebexol® cream lotion / Wofacutan washing lotion / Hans Karrer Hydrocream MicroSilver](#)
- Yarrow oil, [435](#)
- Ylang-ylang oil, [439](#)
- Zingiber officinalis oleum, *see* [Ginger oil](#)